

ISSN 0377- 9335

ENTOMON

Volume 48

December 2023

Number 4

47 YEARS OF EXCELLENCE



ASSOCIATION FOR ADVANCEMENT OF ENTOMOLOGY

ENTOMON

ENTOMON is a quarterly journal published by the Association for Advancement of Entomology devoted to the publication of original research in all facets of insects and related branches of Entomology.

EDITORIAL BOARD (2022 – 2025)

Chief Editor:

Palaniswami, M.S., ARS-ICAR (Retd), Thiruvananthapuram, India

Associate Editor:

Prathapan, K. D., Kerala Agricultural University, Thiruvananthapuram, India

Members:

Colvin John, University of Greenwich, London, United Kingdom

David, B.V., International Institute of Biotech & Toxicology, Padappai, India

David, K. J., National Bureau of Agricultural Insect Resources (NBAIR), Bengaluru, India

Debjani Dey, Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi

Jorge Ari Noriega, National Museum of Natural Science (CSIC), Madrid, Spain

Mohandas, N., KAU, Thiruvananthapuram, India

Phan Quoc Toan, Center for Entomology Research, Duy Tan University, Da Nang, Vietnam

Priyadarsanan, D.R., ATREE, Bengaluru, India

Raman, A., Charles Sturt University, NSW, Australia

Susanta Kumar Chakraborty, Vidyasagar University, Midnapore, West Bengal, India

Viraktamath, C.A., UAS, Bengaluru, India

Winston M.O. Thomson, NARI, Guyana

Address all manuscripts to the Chief Editor, ENTOMON, E mail: editor.entomon@kau.in. Submission of a manuscript to ENTOMON implies that the content has neither been published earlier nor will be sent to any other publisher without intimation. At least one of the authors need to be a member of AAE. A fee will be charged for publication.

AAE MEMBERSHIP/ SUBSCRIPTION RATES:

Admission Fee: Rs 100/- (\$10/- for foreigners)

Life Membership Fee for individuals: Rs 6000/- (or US \$ 600/- for foreigners)

Annual Membership Fee for individuals: Rs 1000/- (US \$ 150/- for foreigners)

Annual subscription for Institutions: Rs 6000/- (in India); US\$ 600/- (outside India)

© 2022 by the Association for Advancement of Entomology. All rights reserved

All remittance to the Journal or Association for Advancement of Entomology should be sent to the Secretary, Association for Advancement of Entomology, Vellayani, Thiruvananthapuram 695 522, Kerala. The amount can be transferred directly to the account of Association for Advancement of Entomology in the State Bank of India, Vellayani (Branch) [Account No. 67267412481; IFS Code: SBIN0070019; Swift code for outside India- SBININBBT44] or by cheques / drafts. Request for copies of ENTOMON should be addressed to the Secretary. Email: aae@kau.in; Web: www.entomon.in

ENTOMON is covered in the following abstracting/ indexing journals:

CABI, cabdirect.org, CAB abstracts, Review of Applied Entomology, Science citation index, Zoobank, New Entomological Taxa, Referativny Zhurnal, Zoological Records.

The NAAS rating of the journal is 5.24. The University Grants Commission has recognized ENTOMON in the official list of scientific journals (UGC-Care list Group I). ENTOMON is included in CABI's full text repository, CAB Direct and other related databases. ENTOMON is indexed by the SCOPUS. The journal is partnering with EBSCO for dissemination of papers published. ENTOMON is a member of the Committee on Publication Ethics (COPE).

ENTOMON

Vol. 48

December 2023

No. 4

Contents

	Page
<p>https://doi.org/10.33307/entomon/v48i4/982 Bees of the genera <i>Hoplonomia</i> and <i>Crociaspidia</i> (Halictidae, Nomiinae) of India, with key to species <i>K.T. Manjusha and T. Jobiraj</i></p>	485
<p>https://doi.org/10.33307/entomon/v48i4/983 Taxonomical studies of dragonfly nymphs (Odonata, Libellulidae) using their exuviae <i>Shaun Paul Adambukulam and Francy K. Kakkassery</i></p>	497
<p>https://doi.org/10.33307/entomon/v48i4/984 Screening of wild <i>Ipomoea</i> genotypes for resistance against sweet potato weevil <i>Cylas formicarius</i> F. based on multiple choice bioassay and phytochemical constituents <i>B.G. Sangeetha, C.A. Jayaprakas, T. Makesh kumar,</i> <i>T. Santhosh kumar, Shirly Raichal Anil and L.S. Rajeshwari</i></p>	511
<p>https://doi.org/10.33307/entomon/v48i4/986 Synthesis and characterization of silver nanoparticles using <i>Datura</i> <i>metel</i> L. (Solanaceae) leaf extract and its larvicidal activity on <i>Epilachna vigintioctopunctata</i> F. <i>C. Sundareswari and D.N.P. Sudarmani</i></p>	519

- <https://doi.org/10.33307/entomon/v48i4/987> 525
 Redescription of *Carbula indica* (Westwood, 1837)
 (Hemiptera, Heteroptera, Pentatomidae) from West Bengal, India
 with a key to the Indian species of the genus
Amartya Pal, Swetapadma Dash, Devanshu Gupta and P.C. Saha
- <https://doi.org/10.33307/entomon/v48i4/988> 531
 Evaluation of the insecticidal action of polyphenolic compounds
 from *Streblus asper* (Lour.) on the red cotton bug,
Dysdercus cingulatus (Fab.) (Hemiptera, Pyrrhocoridae)
L. Anila and M.S. Hashim
- <https://doi.org/10.33307/entomon/v48i4/989> 537
 Bioefficacy of *Tagetes minuta* L. against
Aphis craccivora Koch (Hemiptera, Aphididae)
V.V. Ashish, Haseena Bhaskar, U. Sreelatha and K.B. Deepthy
- <https://doi.org/10.33307/entomon/v48i4/990> 545
 Exploration of natural enemy fauna of aphids and associated ant
 species from eastern dry zone of Karnataka, India
V.S. Chinnu, Vidya Mulimani, S. Shanas and N. Sumithramma
- <https://doi.org/10.33307/entomon/v48i4/991> 553
 A new species of *Cigaritis* Donzel, 1847
 (Lycaenidae, Aphnaeinae) from the southern Western Ghats of
 Peninsular India
S. Ramasamy Kamaya Naicker, Sujitha Prabhakaran Chandrika, Baiju Kochunarayanan, Jebine Jose, Manoj Kripakaran, Rajkumar Chidambaram Palaniappan, Vinayan Padmanabhan Nair and Kalesh Sadasivan
- <https://doi.org/10.33307/entomon/v48i4/992> 567
 Redescription of *Cyphochilus niveosquamosus* (Blanchard, 1851)
 (Coleoptera, Scarabaeidae, Melolonthinae) with notes on two
 species of the genus from India
K.V. Prakash and H.M. Yeshwanth

<https://doi.org/10.33307/entomon/v48i4/993> 575

Evaluation of pongamia oil soap against red spider mites, *Tetranychus urticae* Koch and its effect on mite predators in brinjal ecosystem

P. Jeevitha, K.M. Sreekumar and Mukul Kumar Deka

Short Commuination

<https://doi.org/10.33307/entomon/v48i4/994> 583

Eri silkworm pupae as fish meal replacement in common carp (*Cyprinus carpio* L.) feed in indoor aquaria

*R.K. Gokulakrishnaa and Selvamuthukumaran
Thirunavukkarasu*

<https://doi.org/10.33307/entomon/v48i4/995> 591

Record of *Anoplocnemis phasianus* (Fabricius, 1781) (Hemiptera, Heteroptera, Coreidae) from Goa, India

*Vijaykumar S. Gadekar, Ayesha E. Shetkar and
Aishwarya S. Naik*

<https://doi.org/10.33307/entomon/v48i4/996> 595

Efficacy of insecticides against cashew leaf miner, *Acrocercops syngamma* Meyrick (Lepidoptera, Gracillariidae)

*O. Chethana, Jayalaxmi Narayan Hegde,
K. Rajashekharappa, T. Basavaraj Naik and B.V. Champa*

<https://doi.org/10.33307/entomon/v48i4/997> 599

Biology of shoot and fruit borer, *Earias vittella* Fabricius (Noctuidae, Lepidoptera) on okra

Bikash Subba, Nilanjana Chudhuri and S.K. Senapati

<https://doi.org/10.33307/entomon/v48i4/998> 605

Lysinibacillus fusiformis: A novel mosquitocidal bacterium isolated from Western Ghats, Kerala, India

*K. Aneha, Bhagyashree Bora, P. Hemaladkshmi,
S. Manikandan, Jibi Lukose, V. Abhisubesh,
Kakhuangailiu Gangmei, Sahadiya Mandodan,
A. Mathivanan, K. Vijayalakshmi and S. Poopathi*

https://doi.org/10.33307/entomon/v48i4/999 New faunistic records of two species of pygmy backswimmers (Hemiptera, Heteroptera, Pleidae) from Kerala, India <i>K. Jyothylakshmi and S. Nandakumar</i>	611
https://doi.org/10.33307/entomon/v48i4/1000 <i>Calamus tenuis</i> Roxb. a new host for <i>Chilo suppressalis</i> (Walker) (Lepidoptera, Crambidae) <i>Selim Mehmud, Sasanka Shekhar Ghosh, Himu Roy,</i> <i>Dip Jyoti Haloi and Pranab Bujarbarua</i>	617
Contents of volume 48, issues 1 - 4	619
Acknowledgements	630
Author index Volume 48, issues 1 - 4	638

Bees of the genera *Hoplonomia* and *Crocisaspidia* (Halictidae, Nomiinae) of India, with key to species

K.T. Manjusha* and T. Jobiraj

Department of Zoology, Government College Kodanchery (Affiliated to University of Calicut),
Kodanchery, Thamarassery, Kozhikode 673580, Kerala, India.
Email: manjugck23@gmail.com

ABSTRACT: The nomiine bees of the genera *Hoplonomia* Ashmead and *Crocisaspidia* Ashmead occurring in India are reviewed. Specimens were collected from Kerala. Biology and distribution of five species are summarized. An identification key to *Hoplonomia* and *Crocisaspidia* of India is provided.
© 2023 Association for Advancement of Entomology

KEY WORDS: Hymenoptera, Apoidea, taxonomy, distribution

INTRODUCTION

Hoplonomia Ashmead, 1904 is mainly a south Asian genus occurring from Afghanistan to Australia. It is not known from Africa (Michener, 2007). This genus is recognized by the banded abdomen and basally fused lamelliform projections on the metanotum. *Hoplonomia* is considered intermediate between *Acunomia* and *Crocisaspidia*. *Hoplonomia* is represented by 16 species worldwide. Five species viz, *Hoplonomia elliotii* (Smith, 1875), *H. kulliensis* (Tomar & Tomar, 2005), *H. westwoodi* (Gribodo, 1894), *H. callichora* (Cockerell, 1911) and *H. incerta* (Gribodo, 1894) are known from India. *Crocisaspidia* Ashmead, 1899 is characterized by the presence of double projection of scutellum and metanotum. The body is more robust than in most nomiine genera and is the only group of Nomiinae in which the coloured tergal bands are sometimes broken medially and these broken bands resemble *Thyreus*. Currently a single species is

known from India. Eleven species were revised by Pauly (1990). In this paper four species of *Hoplonomia* viz., *H. elliotii*, *H. incerta*, *H. westwoodi*, and *H. callichora* and one species of *Crocisaspidia* viz., *C. buddha* (Westwood, 1875) are diagnosed. *Hoplonomia kulliensis* probably a synonym of a species already described and being a doubtful species is not included here for diagnosis Pauly (2009).

MATERIALS AND METHODS

Literature was surveyed for published reports of *Hoplonomia* in India (Bingham, 1897; Pauly, 2009; Saini and Rathore, 2012; Pannure and Belvadi, 2017). Data regarding specimens from the six major collections for India viz., American museum of natural history (AMNH), Regional Museum of Natural History (RMNH), United States National Museum (USNM), Smithsonian Institution, Washington, Statens Museum for Kunst (SMUK) Denmark, Natural History Museum (NHMUK),

* Author for correspondence

Institut royal des Sciences naturelles de Belgium (IRSNB); and Zoological Museum, University of Copenhagen (ZMUC) were obtained. Materials were also collected from the state of Kerala, India using sweep nets. These materials were examined and preserved in the Department of Zoology, Government College Kodanchery, Kozhikode, Kerala, India (DZGCK). Distribution maps were prepared using all localities from India from where *Hoplonomia* and *Crocisaspidia* have been collected and was prepared using SimpleMapp (Shorthouse, 2010).

RESULTS AND DISCUSSION

Systematics

Halictidae, Nomiinae

Genus *Hoplonomia* Ashmead

Description: Length 6.5 to 11.5mm. Metanotum with two broad, basally fused lamelliform projections as in *Crocisaspidia*. Scutellum lacks projections or has only small ones in some males. Body is generally smaller and slender when compared to *Crocisaspidia*. Intermediate tibia with strong sub apical teeth. Propodeum sub vertical, defined, triangular propodeal area and tegulae normal. Clypeus in the form of plateau, flanked on the front with a median carina more or less marked.

Presence of two lamelliform projections on the metanotum is a characteristic feature which distinguishes *Hoplonomia* from other Nomiinae. This genus is allied to *Curvinomia* Michener on account of the abdomen being banded. Post scutellum in both sexes is armed with two straight spines. Scutellum provided with median depression and the hind angles ending in a small tubercle.

Distribution: *Hoplonomia* is known from Japan, China, India, South East Asia, Philippines, Indonesia, New Guinea, Bismarck, Solomon Islands, Australia, south to southern Queensland and Madagascar. It is not known from Africa. In India it is known from Maharashtra, Himachal Pradesh, Kerala, Karnataka, West Bengal, Andhra Pradesh, Pondicherry, Tamil Nadu and Rajasthan.

1. *Hoplonomia callichlora* (Cockerell, 1911)

Nomia callichlora Cockerell, 1911: 219, ♀. Holotype ♀: "N.W. India" [Pakistan], Karachi, leg. E. Comber, NHMUK (examined).

Distribution: North India (Fig. 26)

Materials Examined: Rajasthan, Abu, 24.64N; 72.77E, 1 ♀, leg. C.G. Nurse (NHMUK).

Diagnosis: *Length*, 9mm. Black, abdomen with broad bright green bands (the first three flushed with vermillion) on the first four segments; post scutellum with two strong black teeth; mandibles dark red in middle; flagellum ferruginous beneath; hind legs almost entirely amber colored; middle of face and clypeus with a delicate keel; tergite band covers the entire apical depression. *Male*. Femur less developed. *Female*. scutum with more patchy punctuation.

Shows much resemblance to *Hoplonomia elliotii* Smith, it is however readily separated as follows. Rather smaller, face narrower, covered with white hair; mesothorax smaller, more shining, with smaller punctures, a band of grayish-white hair running along lateral and hind margins, and the disc with much pale hair, hair of outer side of middle tibiae entirely white, green abdominal bands twice as broad.

2. *Hoplonomia elliotii* Smith, 1875

Nomia elliotii Smith, 1853:89, ♂, India Nomen nudum.

Nomia elliotii Smith, 1875:44, pl. 1, fig. 7, ♀ ♂

Types: Madras, Barrackpore and Nischindipore. (not examined)

=? *Nomia simplicipes* Friese 1897: 73, ♂, nec. ♀ Lectotype ♂: China, Kaulun near Hong Kong designated by Cockerell, 1919: 3 (not examined) Syn.of Pauly (2009)

Distribution: The species is known from India to Indochina and Southern China. In India it has been reported from Maharashtra, Karnataka, Kerala, Tamil Nadu, West Bengal, Goa and Assam (Fig. 25)




Fig. 1 *Crocisaspidia buddha*, male, scutellum and metanotum, Fig. 2 *Hoplonomia westwoodi*, male, scutellum and metanotum, Fig. 3 *Hoplonomia westwoodi*, scutellum and metanotum, Fig. 4 Hind leg of *Crocisaspidia buddha*, Fig. 5 Enlarged metabasitarsus of *Crocisaspidia*, Fig. 6 Narrow metabasitarsus of *Hoplonomia*

Materials Examined: Kerala, Malappuram, Calicut University, botanical garden, 11.13N; 75.89E, 27. ix.1986, 2 ♂, leg. E.E. Grissell (USNM); Trivandrum, Ponmudi Range, 8.48N; 76.94E, 3000ft, v.1972, 2 ♀, leg. T.R.S. Nathan (SMUK); Palakkad, Walayar Forest, 10.84N; 76.84E, 700ft, x.1959, 4 ♂, leg. P.S. Nathan (RMNH); Palakkad, Walayar Forest, 10.84N; 76.84E, 1000 ft, ix.1952, 1 ♀, leg. P.S. Nathan (IRSNB); Kasaragod, Neeleswaram, 12.25N; 75.12E, 24.vi.2018, 1 ♀, leg. Asha (DZGCK); Wayanad, Pulpally, 11.79N 76.16E, 13.v.2018, 1 ♀, leg. Manjusha (DZGCK); Kasargod, Periyar, 12.40N; 75.09E, 10.i.2015, 1 ♀, leg. Sangeetha (DZGCK); Kozhikode, Balussery, 11.45N; 75.82E, 23. v. 2018, 1 ♀ leg. Adersh (DZGCK); Kannur railway, 11.87N; 75.36E, 6. vii. 2014, 1 ♀, leg. Asha (DZGCK); Kasaragod,

Padannakkad, 12.26N; 75.36E, 2. v. 2018, 1 ♀ leg. Adarsh (DZGCK); Kannur, Madayipara, 38m, 12.16N; 75.33E, 1 ♂, 13. viii. 2015, coll. Prashantha, C, 1 ♀, 13. viii. 2015, coll. Pradeepa, S.D.

Maharashtra, Salsette Island, 19.11N 72.89E, 1910, 2 ♀, (NHMUK).

Tamil Nadu, Coimbatore, 11.00N 76.96E, xi.1958, 1 ♀ leg. P.S. Nathan (RMNH); Annamalai Hills, 11.11N 77.335E Cinchona, 3500ft, v.1960, 1 ♀, leg. P.S. Nathan (RMNH); Nilgiris Hills, Devala, 11.47N; 76.38E, 3200 ft, v.1961, 1 ♀ leg. P.S. Nathan (AMNH).

West Bengal, Calcutta, 22.57N; 88.36E 1 ♂, 1 ♀ (IRSNB).



Fig. 7



Fig. 8



Fig. 9



Fig. 10



Fig. 11



Fig. 12

Fig. 7 *Hoplonomia incerta*, male, metasoma, Fig. 8 *Hoplonomia elliotii*, male, metasoma, Fig. 9 *Hoplonomia westwoodi*, male, metasoma, Fig. 10 *Hoplonomia elliotii*, male, hind leg, Fig. 11 *Hoplonomia westwoodi*, male, hind leg, Fig. 12 *Crocispidia buddha*, female, scutellum and metanotum

Diagnosis: Female. Colouration. Head and mesosoma black, bright emerald green transverse fasciae in the apical margin of the basal four abdominal segments, post scutellar spine dark chestnut brown. **Sculpture.** Head and thorax closely and finely punctured, punctures more sparsely and coarsely on scutellum, smooth and opaque abdomen, punctures on the base of the segments, clypeus flat with carina in the mid line, scutellum with deep notch, presence of spines in the middle of post scutellum posteriorly, fine punctures in the space at the base of the median segment, transverse impressed lines across the middle of the basal four abdominal segments. **Pubescence-** white somewhat griseous thin pubescence, legs with pale glittering pubescence.

Male. Tubercles at the lateral angles of the scutellum and medial spines on the post scutellum

more prominent, posterior femur greatly swollen, tibia flattened, the apex on the inner side produced into blunt testaceous process.

3. *Hoplonomia incerta* (Gribodo, 1894)

Nomia incerta Gribodo 1894: 129, ♂, ♀, Lectotype ♀: Java, collection Gribodo, MCSN (examine), designated by Pauly, 2009

=*Nomia punctata* Westwood 1875: 213, ♂, ♀, nec Smith 1858. Type: China, BMNH (nonexamine). Syn. of Pauly (2009)

=*Nomia pilosella* Cameron 1904: 211, ♂ Holotype ♂: India, Khasia Hills, OUMNH (examined) Syn of Pauly (2009)

=*Nomia maturans* Cockerell 1912: 10. Holotype: Formose, Takao, 10. XI. 1907(not examined, sec description), Syn. of Pauly (2009)



Fig. 13



Fig. 14



Fig. 15



Fig. 16



Fig. 17



Fig. 18

Fig. 13 *Hoplonomia elliotii*, female, scutellum and metanotum, Fig. 14 *H. incerta*, female, metasoma, Fig. 15 *H. callichlora*, female, metasoma, Fig. 16 *H. elliotii*, female, metasoma, Fig. 17 *H. westwoodi*, female, metasoma, Fig. 18 *H. elliotii*, female, metasoma

Distribution: Nepal, India, China, Korea, Peninsula, Japan, Taiwan, Indonesia. In India it has been reported from Arunachal Pradesh and Meghalaya (Fig. 26)

Materials Examined: Sikkim, 27.60N; 88.45E, 1: ♂ leg. F.A. Noller (UZMK).

Diagnosis: Metanotum with double projection; Integumentary bands blue coloured, opalescent, band absent on the first abdominal segment; Female has the mesothorax with punctures of unequal size and the lobes of the post scutellum with obtuse process.

4. *Hoplonomia westwoodi* Gribido, 1894

Nomia simillima Smith 1875: 44, pl. II, fig. 4, ♂ = *Nomia carinata* Smith 1875: 57, ♀. Holotype ♀: Ceylon, NHMUK. **Syn. Nov.**

= *Nomia westwoodi* Gribodo 1894: 128, nom. nov. pour *Nomia simillima* SMITH 1875, nec 1863

= *Nomia erythrogaster* Cameron 1898: 61, pl. 4, fig. 10, ♂. Lectotype: India, Poona, OUMNH, designated by Baker 1993: 258

Distribution: This species is found in India, Afghanistan, Pakistan and Srilanka. In India it has been reported from Maharashtra, Andhra Pradesh,



Fig. 19



Fig. 20



Fig. 21

Fig. 19 *Hoplonomia westwoodi*, female, mesosoma,
 Fig. 20 *H. elliotii*, female hind leg,
 Fig. 21 *H. westwoodi*, female hind leg

Karnataka, Pondicherry, Rajasthan, Tamil Nadu and West Bengal (Fig. 27)

Materials Examined: Andhra Pradesh, Hyderabad, Utrap, 17.38N 78.45E, 14-20. x.1997, 1 ♂, leg. Dr. Olejnicek (OOL): Bapatla, 15.93N; 80.38E, 12. xii. 2006, 1 ♀, coll., David, K.J.

Goa, Mormugao, 15.37N; 73.87E, vi. 1925, 1 ♂, ix. 1925, 1 ♀, leg. J.C. Bridwell (USNM).

Gujarat, Banas kantha, Deesa, 24.26N; 72.18E, iv. 1898, 1 ♂, x.1898, 1 ♂, iii. 1899, 2 males, 3 females, leg. C.G. Nurse (BMNH).

Jharkhand, Ranchi, 23.37N; 85.32E, iv. 1957, 1 ♂, leg. G. Angalet (USNM).

Karnataka, Mangalore, 12.86N; 74.84E xi.1926, 3 ♀, leg. J.C. Bridwell (USNM); Bangalore, GKVK, 930m, 12.76N; 77.74E, 1 ♀, 5. vi. 2013, coll. Girish; 1 ♀, 23. iv. 2012, coll. Arun B.C; 1 ♀, 7. i. 2014, 1 ♀, 20. i. 2014, coll. Arati Pannure; 1 ♂, 10. ii. 2008, coll. Nayana, E. D; 4 ♀ and 1 ♂, 4, 5, 6, ix. 2014, coll. Pradeep; GKVK, 934m, 12.76N; 77.74E, 1 ♀ and 3 ♂, 31. i. 1982, coll. B. Mallik; Hebbal, 900m, 13.03N 77.59E, 2 ♀, 31. x. 2014, 1 ♀, 11. xi. 2014, 1 ♀, 16. xi. 2014, 1 ♀, 21. ii. 2015, 1 ♂, 6. i. 2015, 1 ♂, 20. xi. 2015, coll. Zameeruddin; Sadahalli, 906m, 13.21N; 77.64E, 1 ♀, 6. ii. 2015, 1 ♀, 26. ii. 2015, coll. Zameeruddin; Belgaum, Arabhavi, 582m, 16.22N; 74.82E, 2 ♀, 20. ix. 2014, coll. Revansidda; Bellary, 15.14N; 76.92E 1 ♀, 21. ix. 2011, coll. M. Srinivasa; Chikkamagaluru, Kadur, 758m, 13.49N; 75.73E, 1 ♀, 24. xi. 2014, coll. Prashantha, C; Mudigere (20 km SW), 2 ♂, 15. iii. 2008, coll. Nayana, E; Hassan, karekere, 934m, 12.96N; 76.25E, 2 ♀, 23. vi. 2014, coll. Zameeruddin; Dakshina Kannada, Kankandi, 20m, 12.86N; 74.85 E, 1 ♀, 5. iii. 2015, coll. Prashantha, C; Kodagu, Ponnampet, 85m, 12.14N; 75.94E, 1 ♀, 21. vii. 2015, coll. Prashantha, C; Kolar, Horticulture college, 830m, 13.13N; 78.16E, 1 ♀, 16. xii. 2014, coll. Zameeruddin; Koppal, Munirabad, 466m, 15.27N; 76.32E, 1 ♀, 5. xii. 2012, coll. Najeer; Mandya (Sasalu), 841m, 12.52N; 76.89E, 2 ♀, 6. x. 2014, coll. Pradeep; VC farm, 727m, 12.52N; 76.89E, 2 ♂, 10. viii. 1982, coll. B. Mallik; Mysore, Chinnamballi, 716m, 12.09N; 76.83E, 2 ♀, 19. vii. 2015, coll. Prashantha, C; COH, 824m, 13.13N; 78.16E, 1 ♂, 20. vii. 2015, coll. Prashantha, C; Hunsur, 12.30N; 76.29E, 1 ♀, 18. iv. 2009, Dhanyavathi, P.N; Nanjangud, 24.64N; 72.77E 1 ♂, 24. i. 2009, coll. Dhanyavathi, P. N; Banur, 13.32N; 75.77E 1 ♀, 23. iv. 2009, Dhanyavathi, P.N. Udupi, Brahmavar, 13.43N; 74.74E, 1 ♂, 12. iv. 1985, coll. A.R.V Kumar.

Kerala, Malappuram, Calicut University, botanical garden, 11.13N; 75.89E 27. ix. 1986, 1 ♂, 1 ♀, leg. E.E. Grissell (USNM); Palakkad, Walayar Forest, 10.84N; 76.84E 700ft, ix.1959, 1 ♀x. 1959, 5 ♂ 9 ♀, leg. P.S. Nathan (RMNH), ix. 1953, 1 ♀ leg. P.S. Nathan (IRSNB).

Fig. 22. *Crocisaspidia buddha*, female

Fig. 23

Fig. 22 *Crocisaspidia buddha* female; Fig. 23 *C. buddha*, male

Maharashtra, Amraoti Dist., Melghat Tiger res, Dhakna, 21.15N; 77.64E, 1300ft, 21. ii. 1976, 1 ♂, leg. M.L. Ripley (USNM); Raigad, Matheran, 18.98N; 73.26E iii. 1899, 1 ♀ (BMNH), 3 ♂ 3 ♀ iv. 1899, 1 ♂, leg. C.G. Nurse (BMNH); Bombay Presidency, 19.04N; 72.90E, 1 ♀ (BMNH).

Orissa, Teypone, 20.5N; 84.4E, 1775ft, ix. 1958, 5 ♀ x. 1958, 6 ♀ leg. P.S. Nathan (RMNH).

Pondicherry, Karaikal, 11.92N; 79.83E, 22. ii. 1946, 1 ♂ leg. P.S. Nathan (USNM); i. 1964, 1 ♂, iii. 1964, 1 ♂, vii. 1964, ♀ leg. P.S. Nathan (SMUK); xii. 1958, 1 ♀, .1959, 2 ♂, iii. 1962, 2 ♂, 1 ♀ iii. 1962, 25 ♂, iv. 1962, 58 ♂, 2 ♀ vi. 1962, Ashmead, 5 ♀ vii. 1962, 2 ♂, 4 ♀, viii. 1962, 3 ♀, iii. 1964, 1 ♀ leg. P.S. Nathan (RMNH); Nettiapackam, 11.86N; 79.63E, x. 1963, 1 ♂, leg. P.S. Nathan (SMUK); 10km N.

Auroville, Discipline Farm, 12.00N; 79.80E, 2. iii. 2010, 1 ♀, leg. & col. F. Burger: Karaikal Territory, Kurumbagaran, 10.92N; 79.83E, 12. viii. 1954, 1 ♀, ix. 1954, 1 ♀, leg. P.S. Nathan (IRSNB).

Rajasthan, Abu, 24.64N; 72.77E 1 ♀, leg. C.G. Nurse (NHMUK).

Sikkim, 27.60N; 88.45E 1 ♂, leg. Fedschlotter (UZMK).

Tamilnadu, Coimbatore, 11.00N; 76.96E 1400ft, vi. 1965, 1 ♂, 2 ♀, ix. 1964, 1 ♀, leg. P. S. Nathan (SMUK), 1. vii. 1953, 2 ♀, ix. 1953, 2 ♀, xi. 1953, 3 ♂ leg. P.S. Nathan (IRSNB); Palani Hills, 10.49N; 77.50E Educu Mtn, 19. xi. 2005, *Mentha* sp., 1 ♂, leg. et col. F. Burger, ix. 1951, 1 ♂, leg. P.S. Nathan (USNM); Tuticorin, 8.80N; 78.14E, 18. x. 1938, 1 ♀




Fig. 24.

Fig. 24 A- *Hoplonomia westwoodi* male, B- *H. westwoodi* female, C- *Hoplonomia elliotii* male, D- *H. elliotii* female, E.- *H. incerta* male, F- *H. incerta* female, G- *H. callichlora* female

(BMNH), vii. 1958, 1 ♂, viii. 1958, 1 ♀, ix. 1959, 1 ♂, iv. 1962, 12 ♀, leg. P.S. Nathan (RMNH); Annamalai Hills, 11.11N; 77.35E, Cinchona, 3500ft, v. 1964, 1 ♂, leg. P.S. Nathan (RMNH); 1.vii.1953, 2 ♀ Tranquebar, 11.02N; 79.85E, vi. 1951, 6 ♀ 9. xii. 1951, 1 ♂, vii. 1953, 3 ♀ xii. 1953, 4 ♀, leg. P.S. Nathan (IRSNB); Nilgiris Hills, Moyar camp, 11.41N; 76.69E, 2100ft, iv. 1954, 13 ♂, leg. P.S. Nathan (IRSNB); Nilgiris hills, Singara, 11.34N;

76.81E, 3400ft, v. 1954, 1 ♀, 1 ♂, leg. P.S. Nathan (IRSNB).

Diagnosis: Similar to *H. elliotii* with posterior lateral angles of the scutellum tuberculate and post scutellum with two teeth; but smaller; posterior femur and tibia not so swollen and thick; inner angle of the apex of tibia is produced and rounded and not forming flat truncate process as in *H. elliotii*; abdomen beneath and posterior legs pale rufotestaceous




Fig: 25




Fig:26

Fig. 25 Distributional map of *Hoplonomia elliotii* recorded so far from India; Fig. 26 Distributional map of *Crocisaspidia buddha*, *H. callichlora*, *H. incerta* recorded so far from India

Subfamily Nomiinae,
Genus: *Crocisaspidia* Ashmead

Diagnosis: Scutellum is characteristic as compared to other Nomiinae for its lateral projections are flat with raised carinate edges and extend straight back not upward from the rest of scutellum. All species are larger with 9 -16mm length. Enamel bands on tergites are on apical margins. In females basal plate of hind tibia is completely rounded. The propodeal area 'V' shaped ending on the posterior face vertically, horizontal only in its lateral parts. Tegulae is in the form of ears.

Distribution: This genus is basically Afrotropical (9 species), one species in Madagascar, and three in the Arabian Peninsula. Currently only a single species is known from India and it is known from West Bengal and Gujarat.

5. *Crocisaspidia buddha* (Westwood, 1875)

Nomia budha Westwood, 1875: 209, pl. IV, fig. 1, ♂. Lectotype ♂: India, OUMNH, designated by Pauly 1990: 48 (reviewed)

= *Nomia bahadur* Nurse, 1904: 568, ♀, ♂.




Fig:27

Fig. 27 Distributional map of *Hoplonomia westwoodi* recorded so far from India

Lectotype ♀: Deesa, BMNH (examined)
Distribution: West Bengal, Gujarat (Fig. 26)

Materials Examined: West Bengal, 23.55N 88.57
Chapra (Mackenzie), 1 ♂ (NHMUK).

Gujarat, Deesa, 24.25N; 72.18E ix.1901, 1 ♀ leg.
C.G. Nurse (NHMUK)

Diagnosis: Males are with characteristically shaped posterior tibias. In females integumentary bands are pale blue green in color; all bristles of the tibia and posterior tarsi with whitish grey bristles; first tergite with sparse punctuations.

Key to Genera and Species of *Hoplonomia* and *Crocisaspidia* of India

Males: antenna of 13 segments; hind leg without scopal hair and femora often thickened; no sting.

Females: antenna of 12 segments; hind legs with scopal hairs; sting.

Males (male of *H. callichlora* unknown)

1. Scutellum with a pair of lateral ear-shaped lamella, metanotum with a pair of large projections (Fig.1) basitarsus of hind leg is large*Crocisaspidia buddha*

- Scutellum mutic (Fig. 2) or with a pair of much reduced teeth (Fig. 3), metanotum with a pair of narrow projections (Figs. 2, 3); basitarsus of hind leg narrow..... *Hoplonomia*

2. First tergum without enamel-like band, but with lateral tuft of white hairs; terga 2 to 4 with yellow, green or blue enamel-like apical bands (Fig.7).....*H. incerta*

- First to fourth terga with yellow, green or blue enamel-like apical bands (Figs. 8, 3)

3. Hind leg nearly completely black, only the apical lobe of the tibia amber, hind femur very thick (Fig.10).....*H. elliotii*

- Hind leg nearly completely amber, hind femora moderately thick (Fig.11).....*H. westwoodi*

Females

1. Scutellum with lateral ear-shaped lamella, metanotum with a pair of large projections (Fig.12)*Crocisaspidia buddha*

- Scutellum mutic, metanotum with a pair of small and narrow projections (Fig.13)Genus *Hoplonomia*

2. First tergum without enamel-like band, but with lateral tuft of white hairs; terga 2 to 4 with yellow, green or blue enamel-like apical bands (Fig.14).....*H. incerta*

- First to fourth terga with yellow, green or blue enamel-like apical bands.....3

3. First tergum with enamel-like apical band about twice as broad, covering the totality of the apical depression (Fig.15).....*H. callichlora*

- First tergum with enamel-like band not exceeding half of the apical depression.....4

4. Scutum with dense punctuation (Fig.18); hind legs largely amber (Fig. 20).....*H. elliotii*

-Scutum with more spacer punctuation (Fig. 19); hind legs largely black (Fig. 11).....*H. westwoodi*

ACKNOWLEDGEMENTS

The authors are immensely thankful to Alain Pauly, Royal Belgian Institute of Natural Sciences, Brussels, Belgium for providing related literature, photographs and immense support throughout the preparation of the article. The authors thank Ranjith A.P. Research Associate at Ashoka Trust for Research in Ecology and Environment for the help rendered in generating the distributional map. The authors also thank Principal and staff of Government College, Kodanchery for providing infrastructural facilities and immense support for this study.

REFERENCES

- Ashmead W.H. (1899) Classification of the bees of the superfamily Apoidea Transactions of the American Entomological Society 26: 49–100.
- Ashmead W.H. (1904) A list of the Hymenoptera of the Philippine islands, with description of new species. Journal of the New York Entomological Society 12: 1–22.
- Baker D.B. (1993) The type material of the nominal species of exotic bees described by Frederick Smith vi.312 pp. Oxford Thesis, Oxford University. Entomologis 154: 1–16.
- Bingham C.T. (1897) The Fauna of British India including Ceylon and Burma Hymenoptera. Vol. 1, Taylor and Francis, London. 579pp.
- Cameron P. (1898) Hymenoptera Orientalia or contributions to knowledge of the Hymenoptera of the Oriental Zoological region. Part VII. Memoirs of the Manchester Literary and Philosophical Society 42 (11): 1–84, pl. IV.
- Cameron P. (1904) Descriptions of new species of aculeata and parasitic Hymenoptera from Northern India. Annals and Magazine of Natural History 13: 211–233.
- Cockerell T.D.A. (1911) Descriptions and records of bees - XXXV. Annals and Magazine of Natural History 7(8): 310–319.
- Cockerell T.D.A. (1912) Descriptions and records of Bees - XLVII. Annals and Magazine of Natural History 10(59): 484–494.
- Cockerell T.D.A. (1919) The metallic coloured halictine bees of the Philippine islands. Philippine Journal of Science 15: 9–13.
- Friese H. (1897) Monographie der Bienengattung *Nomia* (Latr.) (Palaeartische Formen). In: Festschrift zur Feier des fünfzigjährigen Bestehens des Vereins für schlesische Insektenkunde in Breslau. Breslau: Verein für Schlesische Insektenkunde pp45–84.
- Gribodo G. (1894) Note Imenotterologiche, Not II. Bollettino della Societa Entomologia Italiana [Firenze]. pp 76–135, 262–314.
- Michener C.D. (2007) The Bees of the world, John Hopkin University Press, Baltimore.
- Maryland USA. Notes on the Apidae (Hymenoptera) in the collection of the British Museum with Desce. 953pp.
- Pannure A. and Belvadi V.V. (2017) An updated distributional checklist of Bees of the subfamily Nomiinae (Hymenoptera: Apoidea: Halictidae) with new records from south India. ENTOMON 42(4): 311–328.
- Pauly A. (1990) Classification of African Nomiinae (Hymenoptera, Apoidea, Halictidae). Royal Museum Central Africa, Tervuren. Annales Zoological Sciences 261: 1–206.
- Pauly A. (2009) Classification des Nomiinae de la Region Orientale, de Nouvelle-Guinee et des iles de l'Ocean Pacifique (Hymenoptera: Apoidea: Halictidae) Entomologie 79:151–229.

- Saini M.S. and Rathore V.S. (2012) Species checklist of family Halictidae (Hymenoptera: Apoidea) along with keys to its subfamilies, genera & subgenera from India. *International journal of Environmental Sciences* 3(1): 134–166.
- Shorthouse D.P. (2010) Simple Mappr, an online tool to produce publication-quality point maps. (Retrieved from <https://www.simplemappr.net>. Accessed July 25, 2020).
- Smith F. (1875) Descriptions of new species of Bees belonging to the genus *Nomia* Latreille. *Transactions of the entomological society of London* 5: 53–70.
- Tomar S. and Tomar M. (2005) Male genitalia of two Indian halictine bees (Halictidae: Apoidea: Hymenoptera). *Flora and fauna, Jhansi* 11 (2): 148–150.
- Westwood J.O. (1875) Descriptions of some new species of short tongued bees belonging to the genus *Nomia* of Latreille. *Transactions of the Entomological Society of London* 23(3): 207–222.

(Received March 13, 2023; revised ms accepted September 20, 2023; published December 31, 2023)

Taxonomical studies of dragonfly nymphs (Odonata, Libellulidae) using their exuviae

Shaun Paul Adambukulam* and Francy K. Kakkassery

*Research and Postgraduate Department of Zoology, St. Thomas College (Autonomous)
(Affiliated to University of Calicut), Thrissur 680001, Kerala, India.
Email: shaunpaultr@gmail.com, kakkassery@yahoo.com*

ABSTRACT: Taxonomic studies of dragonfly nymphs were done up to species level using their exuviae. Exuviae, being the last instar larval skin, possess all larval features. These features can be used for the identification of odonate nymphs up to species level. Seven species belonging to the family Libellulidae were identified using the features present on exuviae. This is a non-invasive method that can be used for the taxonomic studies of dragonfly nymphs without rearing them in the laboratory. The study describes the morphological features of seven species of dragonfly nymphs belonging to family Libellulidae using exuviae. A taxonomic key for the identified exuviae were also provided.

© 2023 Association for Advancement of Entomology

KEYWORDS: Species identification, non-invasive method, taxonomic key

INTRODUCTION

Odonata comprises dragonflies and damselflies, one of the most ancient insect groups. In Kerala, a total of 174 species of odonates have been reported of which 65 are endemic to the Western Ghats, and 10 to India (Gopalan *et al.*, 2022). The kole wetlands of central Kerala reported 44 species of odonates (Chandran *et al.*, 2021). Being amphibiotic and hemi-metabolous insects, adults are terrestrial, and nymphs have aquatic modes of life. Adult females, after mating, lay their eggs in different aquatic ecosystems. Egg hatch into nymphs. Dragonfly nymphs are strictly predacious, and by using their extensible labium, they capture aquatic insects, crustaceans, small molluscs, larvae of other insects, and oligochaetes. At the end of the aquatic life, the nymph finds a support, such as a rock or plant stem, where it can tear up its skin and

enter the aerial life as an adult. The skin that remains on the rock or plant stem is called exuviae, and they are nothing more than the dried skin of the last larval instar and are therefore vulnerable. Since the exuvia is the moulted skin of the penultimate instar, it possesses all the larval features (Adambukulam and Kakkassery, 2013a, b). The main features that can be used for the taxonomic identification larva includes shape of the labium, number and shape of antennal segments, number of premental and palpal setae, presence or absence of mid dorsal spines, and position of lateral abdominal spines, shape of caudal appendages etc. These features show variation within the family, genus and species level and hence can be used to study the larval taxonomy. Taxonomical studies of dragonfly nymphs were usually conducted either by collecting nymphs from the field or eggs from an ovipositing female in the field and rearing them

* Author for correspondence

in the laboratory. In Kerala, Nirmalakumari and Balakrishnan (1981, 1983) studied the life history of the nymphs of *Rhodothemis rufa* (Rambur, 1842) and *Urothemis signata* (Rambur, 1842) by rearing the larvae in the laboratory. No further studies on Odonate nymphs were reported from Kerala till 2013. Adambukulam and Kakkassery (2013a, b) conducted the larval description studies of dragonfly nymphs by using their exuviae. A typical libellulidae larvae can be identified by their spoon shaped labium and well-developed premental and palpal setae. Interspecific variations were seen in the number of setae. The presence or absence of mid dorsal abdominal and lateral abdominal spines can also be taken as a taxonomic feature. The present study describes the morphological features of seven species of dragonfly nymphs belonging to family Libellulidae using their exuviae.

MATERIALS AND METHODS

During the exuvial survey in Thrissur district, exuvia were collected from a pond (10°27'50"N; 76°11'31"E) located in a paddy field and also from a man-made cement tank located in Ammadam, a small village, 8 km away from Thrissur. Collections were done during the early morning hours, because the process starts in late midnight to avoid predation pressure and hence the collector had the opportunity to see the adult just emerging from the exuviae, which helped to identify the species at the primary level in the field itself. The specimens (exuviae) were collected in plastic bottles of the photographic film and brought into the laboratory. Wet specimens were dried by placing them under an incandescent lamp and then they were pinned and preserved as dry specimens for analysis. The exuviae were dissected by observing through the Stereo Dissection Microscope (CZM 4, LABOMED) for analyzing the larval features present on exuviae. The description of the general morphology follows Tennessen (2019). The SEM images of the main features present on the exuviae (Figs. 1 A-F) were taken using JEOL 6390LA. Exuvia sample was smeared over the multisided adhesive carbon tape fixed on specimen stubs and over-coated with gold using JFC 1600. This ion-sputtering device performs rapid and efficient gold coating on the microscopic

specimen, allowing surface visualization. The SEM measurements were performed at 15 kV accelerating voltage. Different magnifications were used, as indicated on the images. The main taxonomic features present on exuviae includes shape of the labium, the number of premental and palpal setae, setaceous margin of palpus, crenations on palpus, presence or absence of mid dorsal protuberances, their shape (spine like or hook like), lateral spines, extension of wing pads, presence of hairs at base of the abdomen and length and size of caudal appendages. The features observed on exuviae were compared with original larval descriptions by Kumar (1984, 1988, 1990) and Hussain and Riaz (1999a, b). The final plates were prepared using Leica S80AP0 stereo microscope (Leica Microsystems, Germany) with an in-built camera (Leica MC 170 HD) and LAS auto-imaging software. The specimens were retained in the Entomology museum of Department of Zoology, St. Thomas College Thrissur.

RESULTS AND DISCUSSION

A total of 105 exuviae were collected during the study. They include *Acisoma panorpoides* Rambur, 1842 (12 nos.), *Bradinopyga geminata* (Rambur, 1842) (22 nos.), *Neurothemis tulia* (Drury, 1773) (9 nos.), *Orthetrum sabina* (Drury, 1773) (26 nos.), *Rhyothemis variegata* (Linnaeus, 1763) (12 nos.), *Tholymis tillarga* (Fabricius, 1798) (11 nos.) and *Zyxomma petiolatum* Rambur, 1842 (13 nos.). All belong to the family Libellulidae.

1. *Acisoma panorpoides* (Figs. 2 A-F)

A small exuvia having dark brown colour; body length ranges between 16.2–18mm; head small, triangular with compound eyes projecting antero-laterally; antennae seven segmented and filiform; labium is spoon shaped; extends up to the coxae of mid leg at rest; distal margin of prementum is triangular, and projecting; premental setae 11+11; palpal setae 8+8; distal margin of palpus formed into crenations with claviform setae; thorax bears long legs; hind leg extends beyond the body; wing pads diverging; hind wing pad extends up to the middle of the 7th segment; abdomen is more or less oval shaped; mid dorsal protuberances absent;




Fig. 1 Taxonomic features on Exuviae. A-Premental setae in *Rhyothemis variegata*; B-Premental setae in *Bradinopyga geminata*; C- Palpal crenations in *Pantala flavesecens*; D- Palpal spines in *B. geminata* E- Mid dorsal protuberances in *Orthetrum sabina*; F- Abdominal hairs in *O. sabina*




Fig. 2 *Rhyothemis variegata*. A-Dorsal view; B-Ventral view ; C-Premental setae D-Palpal setae; E- Mid dorsal spines F- Lateral spines and Wing pads

lateral spines on S8 and S9; S10 smaller than anal appendages; epiproct triangular, broad at the base; cerci shorter than epiproct and are pointed; paraproct equal length of epiproct. These features were compared with the larval features given (Kumar, 1984) by rearing the larvae in the laboratory.

Remarks: *A. panorpoides* is widely distributed in India, Nepal, China, Sri Lanka, Burma, Thailand, Malaysia, Indonesia, Japan, Taiwan, Philippines, Africa and Madagascar. In Kerala they usually associated with ponds and canals in paddy fields and also near to other stagnant water bodies and their oviposition occurs just after first monsoon rain. The emergence occurs at the end of monsoon season. The exuviae found to be attached to lower side of aquatic plants at a height of 10-20cm above water level.

2. *Bradinopyga geminata* (Figs. 3A-F)

A medium sized transparent exuvia; body length ranges between 19–22mm; head small, triangular with compound eyes projecting antero-laterally; antennae seven segmented and filiform; labium is spoon shaped; extends up to the coxae of first pair of leg at rest; distal margin of prementum triangular, and projecting; premental setae 19+19; palpal setae 16+16; distal margin of palpus crenated with claviform setae; long slender legs with dark bands on femur and tibia; wing pads diverging; hind wing pad extends up to the middle of the 6th segment; abdomen is broader than head; dorsally convex; ventrally flattened; irregular dark spots present; abdominal segments S2–S8 almost equal sized; S9–S10 tapering; mid dorsal protuberances absent; well-developed lateral spines on S8 and S9; S10 smaller than anal appendages; epiproct triangular, acuminate broad at the base; cerci shorter than epiproct and acuminate; paraproct equal length of epiproct. The exuvial features were compared with the larval features Sangal and Kumar (1970).

Remarks: This species is widely distributed throughout the oriental region. They usually breed in overhead tanks and garden ponds. The exuviae were usually found attached to manmade garden ponds. But in garden ponds in which ornamental

fishes are growing their emergence is found to be less. Sometimes they emerge in large numbers. It is a multivoltine species and emerges throughout the year.

3. *Neurothemis tulia* (Figs. 4 A-F)

A medium sized exuvia with brown colour; body length ranges between 11–14mm; head small, pentagonal in outline; with compound eyes projecting antero-laterally; antennae seven segmented and filiform; labium is spoon shaped; extends up to the coxae of first pair of leg at rest; distal margin of prementum triangular, and projecting; premental setae 13+13; palpal setae 8+8; distal margin of palpus crenated; long slender legs with dark bands on femur and tibia; wing pads diverging; hind wing pad extends up to the middle of the 5th segment; abdomen is broader than head; dorsally convex; ventrally flattened; more or less oval shaped; mid dorsal protuberances absent; lateral spines on S8 and S9; S10 smaller than anal appendages; epiproct triangular, acuminate broad at the base; cerci shorter than epiproct and acuminate; paraproct equal length of epiproct. The features were compared with studies of Kumar (1988) and Begum *et al.* (1990).

Remarks: *N. tulia* is an extremely widespread common species, found throughout mainland tropical and subtropical Asia. In Kerala the species is distributed everywhere. They are also multivoltine and emergence occurs all the year around. They breed in small ponds, streams and rivers etc. A variation in the number of premental and palpal setae is observed in Indian species where it is 13 and 8 respectively (Kumar, 1988), for the Bangladesh specimen it is 10 and 9 from Bangladesh (Begum *et al.*, 1990) The difference in the number may be due to the geographical intra specific variations. The collected exuviae show features of Indian representative.

4. *Orthetrum sabina* (Figs. 5 A-F)

A medium sized exuvia with muddy brown colour; body length ranges between 19–22mm; head small, pentagonal in outline; with compound eyes projecting antero-laterally; antennae seven

segmented and filiform; labium is spoon shaped; extends up to the coxae of first pair of leg at rest; distal margin of prementum triangular, and projecting; premental setae 7+7; palpal setae 7+7; distal margin of palpus crenated; legs robust and covered with hairs; wing pads diverging; hind wing pad extends up to the middle of the 6th segment; abdomen is broader than head; dorso-ventrally flattened; more or less oval shaped; spine like mid dorsal protuberances present from S4–S9; setae along posterior margin of all abdominal terga; lateral spines absent; sternum of S9 bears a bunch of hairs; S10 smaller than anal appendages; anal appendages are hairy; epiproct triangular, acuminate broad at the base shorter than paraprocts; cerci less than half length of epiproct and acuminate; paraproct acuminate equal length of epiproct.

Remarks: *O. sabina* is widely distributed in Ethiopian, Oriental and Australian region. Kumar (1973) studied the life history of *O. sabina*, and in his studies that the final instar has 11+11 premental setae and 7+7 palpal setae respectively. But the exuvia collected by the authors show variation in the number of premental and palpal setae 7+7 premental and 7+7 palpal setae and this may be due to geographical variation. In Kerala it is multivoltine species and emerges in large number during pre and post monsoon seasons.

4. *Rhyothemis variegata* (Figs. 6 A-F)

A small sized exuvia with grayish black colour; body length ranges between 11–13mm; head small, pentagonal in outline; with compound eyes projecting antero-laterally; antennae seven segmented and filiform; labium is spoon shaped; extends up to the coxae of first pair of leg at rest; distal margin of prementum triangular, and projecting; premental setae 1+1; palpal setae 5+5; distal margin of palpus crenated; legs long and slender; hind leg extending beyond the abdomen; wing pads diverging; hind wing pad extends up to the middle of the 5th segment; abdomen is broader than head; dorso-ventrally flattened; oval shaped; spine like mid dorsal protuberances present from S3–S9; lateral spines present on S8–9; S10 smaller than anal appendages; anal appendages are hairy;

epiproct triangular, acuminate broad at the base shorter than paraprocts; cerci less than half length of epiproct and acuminate; paraproct acuminate equal length of epiproct.

Remarks: This species is distributed throughout the Oriental region. The larvae have been described by Chowdhury and Akhteruzzaman (1981). In their description the larvae possess 2+2 premental setae but in the collected exuviae it is 1+1. In Kerala it usually emerge during post monsoon season in large numbers.

6. *Tholymis tillarga* (Figs. 7A-F)

A small sized exuvia with grayish black colour; body length ranges between 11–13mm; head small, pentagonal in outline; with compound eyes projecting antero-laterally; antennae seven segmented and filiform; labium is spoon shaped; extends up to the coxae of first pair of leg at rest; distal margin of prementum triangular, and projecting; premental setae 7+7; palpal setae 5+5; distal margin of palpus crenated; long, slender legs with dark bands; hind leg extending beyond the abdomen; wing pads diverging; hind wing pad extends up to the middle of the 5th segment; abdomen is broader than head; dorso-ventrally biconvex; tapering towards the distal end; hook like mid dorsal protuberances present from S3–S9; lateral spines present on S8–9; S10 smaller than anal appendages; anal appendages are hairy; epiproct triangular, acuminate broad at the base; cerci shorter than epiproct and acuminate; paraproct acuminate equal length of epiproct.

Remarks: This species has a wide distribution in all parts of the world except Europe and the Americas. The larvae are sluggish and bottom dwellers and were described (Kumar, 1973; Chowdhury and Akhteruzzaman, 1981). In Kerala the emergence occurs during post monsoon season in large numbers.

7. *Zyxomma petiolatum* (Figs. 8 A-F)

A medium sized exuvia with blackish grey colour; body length ranges between 17–20mm; head small, pentagonal in outline; with compound eyes




Fig.3 *Orthetrum sabina*. A-Dorsal view; B-Ventral view; C-Premental setae; D-Palpal setae; E- Mid dorsal spines F- Hairs on the abdomen




Fig. 4 *Acisoma panorpoides*. A-Dorsal view; B-Ventral view; C-Premental setae; D-Palpal setae; E- Wing pads; F- Lateral spines on the abdomen




Fig 5: *Zyxomma petiolatum*. A-Dorsal view; B-Ventral view; C-Premental setae; D-Palpal setae; E- Mid dorsal protuberances; F- Lateral spines on the abdomen




Fig. 6 *Neurothemis tulia*. A-Dorsal view; B-Ventral view; C-Premental setae; D-Palpal setae; E- Mid dorsal protuberances; F- Lateral spines on the abdomen




Fig. 7 *Bradinopyga geminata*. A-Dorsal view; B-Ventral view; C-Premental setae; D-Palpal setae; E- Palpus with crenations; F- Lateral spines on the abdomen




Fig. 8 *Tholymis tillarga*. A-Dorsal view; B-Ventral view; C-Premental setae; D-Palpal setae; E- Mid dorsal protuberances; F- Lateral spines on the abdomen

projecting antero-laterally; antennae seven segmented and filiform; labium is spoon shaped; extends up to the coxae of first pair of leg at rest; distal margin of prementum triangular, and projecting; premental setae 12+12; palpal setae 8+8; distal margin of palpus crenated; legs long and slender; hind leg extending beyond the abdomen; wing pads diverging; hind wing pad extends up to the middle of the 5th segment; abdomen is broader than head; dorso-ventrally biconvex; tapering towards the distal end; hook like mid-dorsal spines present from S1–S10, spines on S1–S3 and S10 were very small, S4–S9 prominent, sharp and pointed; postero lateral spines on S8–9, spine on S9 is longest and reaching up to the length of epiproct; S10 smaller than anal appendages; anal appendages are hairy; epiproct triangular, acuminate broad at the base; cerci shorter than epiproct and acuminate; paraproct acuminate equal length of epiproct.

Remarks: It is a common species in India, Myanmar and Sri Lanka and is a crepuscular dragonfly appearing shortly before dusk and coming often to house lights. Breed in small stagnant pools and even in domestic wells. The larvae was studied by (Kumar, 1973; Chowdhury and Akhteruzzaman, 1981). The breed in natural water bodies or man-made water bodies. They are multivoltine species and emerge in all seasons throughout the year.

Key to the 7 species of Libellulidae

1. Mid dorsal abdominal spines absent.....2
 Mid dorsal abdominal spines present.....5
2. Body length <15mm; palpal setae <10.....3
 Body length >15mm; palpal setae >10.....4
3. Wing pads extending up to middle of S5; premental setae 13+13; palpal setae 8+8; lateral spines on S8–9;.....*Neurothemis tulia*
 Premental setae 11+11; palpal setae 7+7; lateral spines on S9–10; wing pads extending up to middle of S7.....*Acisoma panorpoides*
4. Wing pads extending to S6; premental setae

17+17; palpal setae 15+15; lateral spines on S8–9;*Bradinopyga geminata*

5. Abdominal lateral spines absent on S8–9.....6

 Abdominal lateral spines present on S8–9.....7

6. Premental setae 7+7; palpal setae 7+7; long tapering abdomen with tuft of hairs at S9.....*Orthetrum sabina*

7. Abdomen with dark mid dorsal stripe from S3–9.....8

 Abdomen without mid dorsal stripe from S3–9...9

8. Premental setae 7+7; palpal setae 5+5; mid dorsal protuberances from S4–9.....*Tholymis tillarga*

9. Mid dorsal protuberance from S3–9 are spine like; premental setae 1+1; palpal setae 5+5;*Ryothemis variegata*

Mid dorsal protuberances from S1–10 are hook like; premental setae 12+12; palpal setae 8+8;*Zyxomma petiolatum*

This present work was focused on morphological features of exuvia (final instar exoskeleton) for identifying dragonfly nymphs up to species level. Most of the nymphal studies of dragonflies were based on the rearing of larvae that directly collected from the field or by collecting the eggs from ovipositing females from the field. Both methods are time consuming. Thus studies, using exuviae can be considered as a noninvasive method for nymphal diversity. This methodology of exuvia collection from the field is suggested for those species which are difficult to observe as adults and also for identifying larval features endemic and endangered dragon fly species. The presence of exuviae also indicates the successful breeding sites of dragonflies and thus health of the aquatic ecosystem can be assessed. The counts of exuviae may give a clear picture of real dragonfly abundance and diversity than do male biased counts, and that numbers emerging from a small or medium sized water body can usually be monitored by a single researcher (Corbet and Hoess, 1998).

ACKNOWLEDGEMENT

Authors thank to Principal of St. Thomas' College, (Autonomous) Thrissur for providing the facilities.

REFERENCES

- Adambukulam P.S. and Kakkassery F. (2013a) Taxonomic and Diversity studies on Odonate nymphs by using their Exuviae. *Journal of Entomology and Zoology studies* 1(4): 47–53.
- Adambukulam P.S. and Kakkassery F. (2013b) Taxonomic studies of the last instar nymph of *Lathrecista asiatica asiatica* (Fabricius 1798) (Family: Libellulidae, Order: Odonata) using its Exuvia, *Journal of Entomology and Zoology studies* 1(5): 103–109.
- Begum A., Bashir M.A. and Biswas V. (1990) On the life history of *Neurothemis tullia tullia* (Drury) from Dhaka, Bangladesh (Anisoptera: Libellulidae). *Indian Odonatology* 3: 11–20.
- Corbet P.S. and Hoess R. (1998) Sex ratio of odonata at emergence. *International Journal of Odonatology* 1(2): 99–118.
- Chandran A.V., Jose S.K. and Gopalan S.V. (2021) Dragonflies and damselflies (Insecta: Odonata) of the Kole Wetlands, central Kerala, India. *Journal of Threatened Taxa* 13(3): 17963–17971.
- Chowdhury S.H. and Akhteruzzaman M.D. (1981) Dragonfly (Odonata: Anisoptera) larvae from Chittagonog. *Bangladesh Journal of Zoology* 9 (2):133–144.
- Gopalan S.V., Sherif M. and Chandran A.V. (2022) A checklist of dragonflies & damselflies (Insecta: Odonata) of Kerala, India. *Journal of Threatened Taxa* 14(2): 20654–20665. doi: 10.11609/jott.7504.14.2.20654-20665
- Hussain R. and Riaz M. (1999a) Description of last instar naiads of *Rhyothemis variegata variegata* (Linnaeus) and *Pantala flavescens* (Fabricius) (Libellulidae: Odonata), *International Journal of Agricultural and Biological Engineering* 1: 145–146.
- Hussain R. and Riaz M. (1999b) The naiads of *Acisoma panorpoides panorpoides* Rambur and *Brachythemis contaminata* Fabricius) (Libellulidae: Odonata). *International Journal of Agricultural and Biological Engineering* 1: 147–148.
- Kumar A. (1973) Description of the last instar larvae of Odonata from Dehra Dun Valley India with notes on biology II (Suborder: Anisoptera). *Oriental Insects* 7: 291–331.
- Kumar A. (1984) Studies on the life history of Indian dragonflies, *Acisoma panorpoides panorpoides* Rambur (Libellulidae: Odonata). *Records of Zoological. Survey of India* 81: 203–213.
- Kumar A. (1988) Studies on the life history of Indian dragonflies *Neurothemis tullia tullia* Drury from Dehra Dun, India (Odonata: Libellulidae). *Indian Odonatology* 1: 35–44.
- Kumar A. (1990) Studies on the life history of Indian dragonflies *Orthetrum sabina sabina* Drury (Odonata: Libellulidae) *Records of Zoological. Survey of India* 88(4): 1–6.
- Kumar A. and Sangal S.K. (1970) Studies on the taxonomy of larvae of Doon Valley Odonata II. *Journal of Natural History* 4: 306–313.
- Nirmala Kumari K.R. and Balakrishnan Nair R. (1981) On the predaceous naiad of *Rhodothemis rufa* (Rambur) (Insecta: Odonata). *ENTOMON* 6(1): 57–60.
- Nirmala Kumari K.R. and Balakrishnan Nair R. (1983) Satiation time and predatory behaviour of the dragonfly nymph *Urothemis signata signata* (Rambur) *Proceedings of the Indian National Science Academy. Part B Biological Sciences* 3(49): 210–216.
- Tennessen K.J. (2019) *Dragonfly Nymphs of North America. An Identification Guide*. Springer, Ebook. ISBN : 978-3-319-97775-1

Screening of wild *Ipomoea* genotypes for resistance against sweet potato weevil *Cylas formicarius* F. based on multiple choice bioassay and phytochemical constituents

B.G. Sangeetha^{*}, C.A. Jayaprakas, T. Makesh kumar, T. Santhosh kumar[#], Shirly Raichal Anil and L.S. Rajeshwari

ICAR-Central Tuber Crops Research Institute, Sreekariyam, Thiruvananthapuram 695011, Kerala, India.

[#]Department of Entomology, College of Agriculture, Vellayani, Thiruvananthapuram 695522, Kerala, India.

Email: Sangeetha.G@icar.gov.in

ABSTRACT: Screening of wild *Ipomoea* spp. and identification of new sources of resistance to the sweet potato weevil (*Cylas formicarius* Fabricius) with *I. palmata*, *I. mauritiana*, *I. obscura*, *I. triloba* were carried out. The leaves, vines and tubers of the different *Ipomoea* sp. were screened using multiple choice bioassay. The insect feeding holes on *I. mauritiana* leaves (1.67 ± 1.528), vines (7.67 ± 2.96) and tubers (12.67 ± 2.309) was significantly less compared with other *Ipomoea* sp. Further, the two-choice bioassay was done, using *I. batatas* and *I. mauritiana* for comparison. Based on the morphological screening different phytochemical constituents was identified using GC-MS analysis of the methanolic extract of roots of selected *Ipomoea* spp. (*I. mauritiana*, *I. palmata* and *I. batatas*). The results indicated that the phytochemical constituent of *I. mauritiana* viz., undecane, quinic acid which is to have insecticidal activity. The major constituent of *I. batatas* comprises of melezitose (38.53%) and alpha-l-rhamnopyranose (21.26%). It can be concluded that the phytochemical constituents of *I. mauritiana* was responsible for the antibiosis. © 2023 Association for Advancement of Entomology

KEYWORDS: Antibiosis, bioactive, insecticidal, bioassay, undecane

INTRODUCTION

The weevil *Cylas formicarius* F. belonging to Coleoptera, Brentidae, is a destructive pest of sweet potato and is widely spread throughout the tropical regions of the world, but the methods of control are the significant problem faced by growers in most countries producing sweet potato. Generally, weevils cause severe feeding destruction

to sweet potato roots, vines, stems and leaves through their life cycle, beginning from the egg stage to adult stage. Weevil infested tubers are bitter due to the production of a terpene compound and the infested tubers are unfit for consumption or convert to livestock, resulting major economic losses (Uritaini *et al.*, 1975; Palaniswami and Mohandas, 1993; Korada *et al.*, 2010a; Kyereko *et al.*, 2019). Although *C. formicarius* prefers sweet potato,

* Author for correspondence

more than 30 species of *Ipomoea* and other genera have been recorded as its host plants (Sutherland, 1986; McConnell and Hossner, 1991). About 500-600 species were included in the genus *Ipomoea* sp. within the Family Convolvulaceae (Austin and Huáman, 1996). Studies have proved that the management sweet potato weevil (SPW) can be done by integrated pest management *viz.*, removal and destruction of hosts, cultural methods, biological control, botanicals, chemical pesticides, tolerant varieties and use of semiochemicals (Palaniswami *et al.*, 1992; Pillai *et al.*, 1993; Palaniswami and Chattopadhyay, 2006; Korada *et al.*, 2010a).

Earlier studies on the identification of the resistant sweet potato genotypes to the weevil indicated only relatively tolerant ones. Studies conducted at AVRDC and Penghu Island has reported screening of the population *I. trifida* x *I. batatas* hybrids with high yield and low weevil infestation (Talekar, 1987). An indigenous cultivar Selopia was identified moderately resistant to the weevil by screening based on crown damage grade index (DGI), percentage tuber damage, tuber DGI, adult emerged per kg infested tuber (Palaniswami and Mohandas 1992). Korada *et al.* (2010b) reported that among the sweet potato genotypes, *viz.*, Goutam, Sourin, Gouri and CIP-6 evaluated for SPW resistance, CIP-6 was the most susceptible. Further in their electroantennogram studies identified the electrophysiological response of female antenna to the volatile extracts of aerial plant parts and roots was higher than the male antenna of the weevil. In olfactometer studies, the headspace volatiles of genotype CIP-6 attracted more number of female *C. formicarius* weevils than volatiles of Gouri, Goutam and Sourin. Variation in the preference of sweet potato genotypes to *C. formicarius* is attributed to differential emission of volatiles from the aerial parts and roots. Reddy *et al.* (2015) reported that the weevil, developed faster on *Ipomoea batatas* than on *I. triloba*.

Anyanga *et al.* (2013) found that hydroxycinnamic acid esters on the exterior and the root latex, decreases weevil's nourishment and oviposition providing resistance to SPW. Okada *et al.* (2019)

identified genetic regions associated with weevil resistance in 90IDN-47 and PSL sweet potato genotypes by genome wide association studies (GWAS) in Japan. In their experiment on the degree of weevil damage to the genotypes, no single nucleotide polymorphisms (SNPs) were identified above the significance thresholds. However, one relatively high peak was found in the 90IDN-47 genotype, which showed resistance to weevils. On the other hand, one relatively high peak was also detected in the PSL genotype, which showed susceptibility to weevils. These results suggest that two regions could affect weevil resistance and may contain the gene(s) controlling weevil resistance. SPW can survive on average longer than four months on sweet potato as well as *I. triloba* (Reddy and Chi, 2015). Hence identification of host plant resistance source against weevil is one of the alternative strategies for the pest management. In the present study, genotypes from different species of *Ipomoea* were selected based on the reports (Reddy and Chi, 2015) on host preference by weevils and experiments were conducted to screen wild *Ipomoea* spp. for resistance against weevil based on the nature of feeding by sweet potato weevils and their phytochemical constituents.

MATERIALS AND METHODS

Multiple choice bioassay: Multiple choice bioassays (Vos and Jander, 2008) were carried out using leaves, vines and roots of plant species *viz* *Ipomoea batatas*, *I. mauritiana*, *I. palmata*, *I. obscura* and *I. triloba*. Five plant samples were placed in large Petridish (180x30mm) and 20 weevils (@1male: 5females) were released in the centre of the Petridish. The sweet potato weevils were reared in Entomology laboratory and were used for bioassay. The position and readings of feeding holes by weevils (cumulative values) were recorded for three consecutive days. Experiment using leaves, vines and roots were conducted separately in three replications and each experiment was repeated thrice. Control samples (leaves, vines, roots) for each experiment were kept separately.

Two - choice bioassay: No-choice bioassays (Vos

and Jander 2008) were carried out using fresh leaves, vines and roots of sweet potato and *I. mauritiana*. The plant samples were placed in large Petridish (180x30mm) and 18 sweet potato weevils (1male: 5 females) were introduced to the centre of the Petridish. The sweet potato weevils were reared in Entomology laboratory and were used for bioassay. The position and readings of feeding holes by weevils (cumulative values) were recorded for three consecutive days. Experiment using leaves, vines and roots were conducted separately in three replications and each experiment was repeated thrice. Control samples (leaves, vines, roots) for each experiment were kept separately.

Data were subjected to analysis of variance using IBM SPSS version 21. The differences between the treatments was measured by tuckey's test at $P_{0.05}$, and the treatment means were compared using the least significant difference at 5 per cent. Data for no choice assay were subjected to t-test at $P_{0.05}$.

Gas chromatography-Mass spectrum analysis:

Further for GC-MS analysis one tuberous wild *I. mauritiana*, one non-tuberous wild *I. palmata* and *I. batatas* were selected for the analysis. The required quantity of the whole plant tubers/roots was washed, air dried and weighed. It was transferred to a flask, treated with methanol of 500ml until the tubers was fully immersed, incubated overnight and filtered through a Whatmann No. 41 filter paper. Before filtering, the filter paper along was wetted with methanol. The filtrate is then concentrated to 5 ml using flash evaporator. The GC-MS analysis was done at Sophisticated Analytical Instruments Facility (SAIF), IIT, Chennai. GC-MS analysis of the methanol extract was performed using an Agilent-Technologies 8890 Network GC system equipped with an Agilent-Technologies 5977 mass selective detector (Agilent-Technologies, Little Falls, CA, USA). For MS detection, the electron ionization mode with ionization energy of 70 eV was used, with a mass range at m/z 50–600. An HP-5MS capillary column (30 m \times 250 μ m, film thickness 0.25 μ m) was used for GC/MS. The column temperature was programmed from 180 to 300 °C

at a rate of 5 °C/min with the lower and upper temperature being held for 3 and 5 min, respectively. GC was performed in the split mode. Helium was used as carrier gas at a flow rate of 1.2 ml/min. An injection 1 μ l was used for each diluted extract. Essential compounds were identified by their retention times and mass fragmentation patterns using data of standards at NIST library

RESULTS AND DISCUSSION

Multiple choice bioassay: The weevil feeding holes on *I. mauritiana* was significantly less compared with *I. batatas*, *I. triloba*, *I. palmata* and *I. obscura*. The insect feeding holes on *I. mauritiana* leaves (1.67 ± 1.52) was significantly low, when compared to other *Ipomoea* species (Table 1). Similarly the same pattern was observed for the three consecutive days and mortality of insects was also observed. The insect feeding holes on *I. mauritiana* vines was less (7.67 ± 2.96), compared to other *Ipomoea* species. The same pattern was observed for the three consecutive days given (Table 2). The insect feeding holes on *I. mauritiana* tubers was significantly low (12.67 ± 2.30), when compared to other *Ipomoea* species (Table 3).

Two-choice bio-assay: The weevil feeding holes on leaf, vines and tubers of *I. mauritiana* and *I. batatas* indicated great variation between them. *I. mauritiana* showed resistance to the weevil (Table 4).

Table 1. Leaf feeding (no. of holes) by the weevils on *Ipomoea* species in multiple choice bioassay

Species	1 st Day	2 nd Day	3 rd Day
<i>I. mauritiana</i>	1.67 ± 1.52^a	4.67 ± 1.15^a	7.33 ± 0.57^a
<i>I. triloba</i>	4.00 ± 1.00^a	7.00 ± 1.00^a	9.67 ± 0.57^a
<i>I. palmata</i>	10.00 ± 2.64^b	15.33 ± 2.30^b	18.33 ± 2.88^b
<i>I. obscura</i>	10.33 ± 2.08^b	14.67 ± 0.57^b	17.67 ± 1.15^b
<i>I. batatas</i>	7.20 ± 4.10^b	17.00 ± 1.73^b	19.00 ± 1.00^b

Mean values (mean \pm standard error) represent error of feeding holes (cumulative) by sweet potato weevil on different *Ipomoea* sp. leaves

Table 2. Vine feeding (no. of holes) by the weevils on *Ipomoea* species in the multiple choice bioassay

Species	1 st Day	2 nd Day	3 rd Day
<i>I. mauritiana</i>	7.67± 2.96 ^{ab}	10.56± 3.37 ^{ab}	14.89± 2.14 ^b
<i>I. triloba</i>	4.22 ± 0.50 ^b	9.56 ± 1.38 ^b	12.00 ± 1.19 ^b
<i>I. palmata</i>	12.67 ± 3.18 ^a	18.00 ± 3.46 ^a	26.44±1.01 ^a
<i>I. obscura</i>	8.78 ± 1.50 ^{ab}	14.22± 3.65 ^{ab}	16.44 ± 2.14 ^b
<i>I. batatas</i>	11.22± 1.16 ^a	17.78± 2.41 ^a	24.22± 2.79 ^a

Mean values (mean±standard, $p_{0.05}$) represent the feeding holes (cumulative) by sweet potato weevil on different *Ipomoea* sp vines

Table 3. Tuber feeding (no. of holes) by the weevils on *Ipomoea* species in the multiple choice bioassay

Species	1 st Day	2 nd Day	3 rd Day
<i>I. mauritiana</i>	12.67 ± 2.30 ^a	22.67± 6.02 ^a	30.33 ± 2.51 ^a
<i>I. triloba</i>	10.67 ± 3.05 ^a	22.67 ± 8.73 ^a	34.33± 4.93 ^a
<i>I. palmata</i>	12.00 ± 0.00 ^a	26.67± 10.40 ^a	31.33± 5.50 ^a
<i>I. obscura</i>	12.67 ± 1.15 ^a	27.33± 3.51 ^a	30.67 ± 3.21 ^a
<i>I. batatas</i>	19.00 ± 2.64 ^b	31.00± 1.00 ^a	37.33± 2.08 ^a

Mean values (mean±standard, $p_{0.05}$) represent the feeding holes (cumulative) by sweet potato weevil on different *Ipomoea* sp roots

Table 4. Weevil feeding (no. of holes) on *Ipomoea* species in the two choice bioassay

Species	1 st Day	2 nd Day	3 rd Day
Leaves			
<i>I. mauritiana</i>	5.67±2.51	8.67±1.15	13.00±1.00
<i>I. batatas</i>	16.00±2.00	18.33±1.52	26.67±2.88
Vines			
<i>I. mauritiana</i>	11.00±1.00	13.00±1.73	14.67±2.51
<i>I. batatas</i>	16.33±0.57	19.00±2.64	25.67±1.15
Tubers			
<i>I. mauritiana</i>	6.33±0.57	12.33±2.51	17.67±1.52
<i>I. batatas</i>	28.33±7.63	35.00±5.00	43.00±4.58

Mean values (mean±standard, $p_{0.05}$) represent the feeding holes (cumulative) by the weevil

Gas chromatography-Mass spectrum analysis:

In all the multiple choice as well as two choice bioassay the feeding of weevils was significantly less in *I. mauritiana* which may be due to the presence of various phytochemical constituents. This shows the non-preference of the weevils always depends on the nature of host plant. GC-MS analysis of methanol extract of samples revealed phytochemical compounds, its retention time (RT) and peak area (%). The bioactivity of the identified compounds reported are presented along with its reference (Table 5). The phytochemical constituent of *I. mauritiana* include compounds undecane and quinic acid which are reported to have insecticidal activity whereas sucrose reported to enhance insecticidal activity. The most prevailing compounds identified in *I. mauritiana* were sucrose (77.01%), quinic acid (20.93%) whereas in *I. batatas* they were melezitose (38.53%) and alpha-I-rhamnopyranose (21.26%).

Higher levels of octadecyl and hexadecyl esters of hydroxycinnamic acids were identified in the root surface and root latex of sub-saharan sweetpotato variety, New Kawogo, contributing resistance to sweet potato weevil (Stevenson *et al.*, 2009). Anyanga *et al.* (2013) reported that the these compounds in high concentrations on root surfaces was strongly associated with resistance against adult oviposition and feeding. They reduce the development of sweet potato weevil larvae and suggested that differences in the concentration of these compounds between varieties explain differences in resistance. Among the five *Ipomoea* species the weevil infestation was significantly less in *I. mauritiana*. Phytochemical screening of methanolic extract revealed the presence of various compounds which are reported to have insecticidal activity. These components might be responsible for the low weevil infestation in *I. mauritiana*.

ACKNOWLEDGEMENTS

The authors are thankful to the Director, ICAR-CTCRI for providing the facilities and support for the research work.

Table 5: List of the phytochemical compounds detected from the methanol extract of *Ipomoea* species through GC–MS analysis

No	R/T	Peak %	Compound	Bioactivity	Reference
<i>I. mauritiana</i>					
1.	6.15	2.06	Undecane <i>Ludwigia stolonifera</i>	Constituent of	Baky <i>et al.</i> , 2021
2.	14.08	77.01	Sucrose	Insecticide activity	Ezhilan and Neelamegam, 2012
3.	17.97	20.93	Quinic acid	Insecticidal activity	Li <i>et al.</i> , 2021
<i>I. batatas</i>					
1.	5.68	3.48	D-Alanine, N-proparglyoxy carbonyl-decyl ester	Constituent of <i>Averrhoa bilimbi</i>	Suluvoy and Grace <i>et al.</i> , 2017
2.	6.85	2.26	DL-Arabinose	Antimicrobial activity	Mohammed <i>et al.</i> , 2018
3.	6.92	2.04	2-Deoxy-2-fluoro-1,6-anhydro- β -d-glucopyranose	Constituent of <i>Alternaria alternata</i>	Kamal <i>et al.</i> , 2015
4.	7.13	2.03	4H-Pyran-4-one,2,3-dihydro-3,5-dihydroxy-6-methyl	Antifungal activity	Teoh and Don, 2015
5.	9.07	3.28	5-Hydroxymethylfurfural	Insecticidal activity	Chuang <i>et al.</i> , 2018
6.	9.66	7.03	5-O-Methyl-d-gluconic acid dimethylamide	Antimicrobial, antioxidant	Kazi and Gude, 2022
7.	10.24	2.19	Octanamide, N-(2-mercaptoethyl)	Secondary metabolite of <i>Vitis vinifera</i>	Kadhim <i>et al.</i> , 2017
8.	12.40	2.77	Methyl 4-nitrohexanoate	Constituent of <i>Hugonia mystax</i>	Vasuki <i>et al.</i> , 2022
9.	14.60	38.53	Melezitose	Insecticidal activity	Gore and Schal <i>et al.</i> , 2004
10.	18.45	4.04	Desulphosinigrin	Antibacterial activity	Olajuyigbe <i>et al.</i> , 2018
11.	26.38	1.97	1H-Benzocyclohepten-7-ol, 2,3,4,4a, 5,6,7,8-octahydro-1,1,4a	Floral volatile constituents of <i>Crataeva religiosa</i>	Sharma <i>et al.</i> , 2018
12.	27.04	1.83	Spiro[4,5]decan-7-one,1,8-dimethyl-8,9-epoxy-4-isopropyl	Anti-inflammatory	Subin and Jagathy 2017
13.	28.22	0.74	Santamarine	Natural antioxidant with anti-photoaging	Oh <i>et al.</i> , 2021
<i>I. palmata</i>					
14.	6.48	1.28	Maltol	Mosquito larvicidal activity	Rajamanikyam <i>et al.</i> , 2017
15.	9.41	1.28	4-Methylmannitol	Constituent of khat leaves	Alsanosy <i>et al.</i> , 2020
16.	10.53	1.77	2H-Pyran-2-onetetrahydro-6-propyl-	Fatty acid composition of <i>Trichosanthes cucumerina</i> bio-oil	Manimaran <i>et al.</i> , 2020

17.	13.88	3.29	Panaxydol	first isolated from roots of <i>P. ginseng</i> induces apoptosis in cancer cells	Takahashi <i>et al.</i> , 1964; Kim <i>et al.</i> , 2016
18.	14.00	2.56	Melezitose	Insecticidal activity	Gore and Schal <i>et al.</i> , 2004
19.	18.07	7.65	Quinic acid	Insecticidal activity	Li <i>et al.</i> , 2021
20.	23.39	2.33	3-(6,6-Dimethyl-5-oxohept-2-enyl)-cycloheptanone	Constituent of <i>Myoporum bontioides</i>	Minh <i>et al.</i> , 2020
21.	26.56	5.22	Scopoletin	Antitermite activity	Adfa <i>et al.</i> , 2010
22.	27.04	2.59	1,8-Naphthalenedione, 8a-ethylperhydro	Constituent of <i>Plectranthus hadiensis</i>	Sripathi <i>et al.</i> , 2017

REFERENCES

- Adfa M., Yoshimura T., Komura K. and Koketsu M. (2010) Antitermite Activities of Coumarin Derivatives and Scopoletin from *Protium javanicum* Burm. *Journal of Chemical Ecology* 36: 720–726.
- Alsanosy R., Alhazmi A.H., Sultana S., Abdalla N.A., Ibrahim Y., Bratty A.M., Banji D., Khardali I. and Khalid A. (2020) Phytochemical screening and cytotoxic properties of ethanolic extract of young and mature khat leaves. *Journal of Chemistry* March 2020: 1-9. doi:10.1155/2020/7897435.
- Anyanga M.O., Muyinza H., Talwana H., Hall D.R., Farman D.I., Ssemakula G.N., Mwanga R.O.M. and Stevenson P.C. (2013) Resistance to the Weevils *Cylas puncticollis* and *Cylas brunneus* Conferred by Sweetpotato Root Surface Compounds. *Journal of Agricultural and Food Chemistry* 61: 8141–8147.
- Austin D.F. and Huamán Z. (1996) A synopsis of *Ipomoea* (Convolvulaceae) in the Americas. *Taxon* 45(1): 3–38.
- Baky H.M., Shawky M.E., Elgindi R.M. and Ibrahim A.H. (2021) Comparative volatile profiling of *Ludwigia stolonifera* aerial parts and roots using vgc-ms/ms and screening of antioxidant and metal chelation activities. *ACS Omega* 6: 24788–24794.
- Chuang J.K., Chen J.K., Cheng L.C. and Hong B.G. (2018) Investigation of the antioxidant capacity, insecticidal ability and oxidation stability of *Chenopodium formosanum* seed extract. *International Journal of Molecular Sciences* 19: 2726.
- Ezhilan B.P. and Neelamegam R. (2012) GC-MS analysis of phytochemicals in the ethanol extract of *Polygonum chinense* L. *Pharmacognosy Research* 4: 11–14.
- Gore J.C. and Schal C. (2004) Laboratory evaluation of boric acid-sugar solutions as baits for management of german cockroach infestations. *Journal of Economic Entomology* 97(2): 581–587.
- Kadhim J.M., Al-Rubaye F.A. and Hameed H.I. (2017) Determination of bioactive compounds of methanolic extract of *Vitis vinifera* using GC-MS. *International Journal of Toxicological and Pharmacological Research* 9(2): 113–126.
- Kamal A.S., Hamza F.L. and Hameed H.I. (2015) Antibacterial activity of secondary metabolites isolated from *Alternaria alternata*. *African journal of Biotechnology* 14(43): 2972–2994.
- Kazi M. and Gude A. (2022) Review on poisonous, pesticidal and medicinal attributes of *Cleistanthus collinus* (roxb.) Benth. *World journal of pharmaceutical and medical research* 8(4): 66–78.
- Kim S.H., Lim M.J., Kim Y.J., Kim Y., Park S. and Sohn J. (2016) Panaxydol, a component of *Panax ginseng*, induces apoptosis in cancer cells through EGFR activation and ER stress and inhibits tumour growth in mouse models. *International Journal of Cancer* 138: 1432–1441.
- Korada R.R., Naskar S.K., Palaniswami M.S. and Ray R.C. (2010a) Management of sweet potato weevil [*Cylas formicarius* (Fab)]: an overview. *Journal of Root Crops* 36: 14–26.
- Korada R.R., Naskar S.K., Prasad A.R., Prasanna A.L. and Jyothi K.N. (2010b) Differential volatile emission

- from sweet potato plant: mechanism of resistance in sweet potato for weevil *Cylas formicarius* (Fab.); Current Science 99: 1597–1601.
- Kyereko W.T., Hongbo Z., Amoanimaa-Dede H., Meiwei G. and Yeboah A. (2019) The major sweet potato weevils; Management and control: A review. Entomology, Ornithology & Herpatology 8: 218.
- Li X., Sivignon C., Silva D.P., Rahb_Y., Queneau Y. and Sanchez M.S. (2021) Design and synthesis of 3, 5- hetero diesters of 4-deoxy quinic acid and their aphicidal activity against *Acyrtosiphon pisum*. Tetrahedron 83: 1–10.
- Manimaran R., Kumar M.M.K. and Narayanan S.N. (2020) Synthesis of bio-oil from waste *Trichosanthes cucumerina* seeds: a substitute for conventional fuel. Scientific Reports 10: 17815.
- McConnell J.S. and Hossner L.R. (1991) pH-dependent adsorption isotherms of glyphosate Journal of Agricultural and Food Chemistry 39 (4): 824–824.
- Minh T.T., Ngoan T.T., Thuong M.T.N., Toan K.H., Truong X.N., Huong T.T., Ly P.T.G., Ai T.T.D. and Khan D.N. (2020) Chemical compositions and bioefficacy against *Spodoptera litura* of essential oil and ethyl acetate fraction from *Myoporum bontioides* leaves Vietnam. Journal of Chemistry 58(1): 57–62.
- Mohammed J.G., Hameed H.I. and Kamal A.S. (2018) Analysis of methanolic extract of *Fusarium chlamydosporum* using GC-MS technique and evaluation of its antimicrobial activity. Indian Journal of Public Health Research and Development 9(3): 229–234. doi:10.5958/0976-5506.2018.00214.0
- Oh H.J., Kim J., Karadeniz F., Kim R.H., Park Y.S., Seo Y. and Santamarine K.S.C. (2021) Shows Anti-Photoaging Properties via Inhibition of MAPK/AP-1 and Stimulation of TGF β /Smad Signaling in UVA-Irradiated HDFs. Molecules 26: 3585.
- Okada Y., Monden Y., Nokihara K., Shirasawa K., Isobe S. and Tahara M. (2019) Genome wide association studies (GWAS) for yield and weevil resistance in sweet potato (*Ipomoea batatas* (L.) Lam) Plant Cell Reports 38: 1383–1392.
- Olajuyigbe O.O., Onibudo E.T., Cooposamy M.R., Ashafa T.O.A. and Afolayan J.A. (2018) Bioactive compounds and *in vitro* antimicrobial activities of ethanol stem bark extract of *Trilepisium madagascariense* DC. International Journal of Pharmacology 14(7): 901–912.
- Palaniswami M.S. and Chattopadhyay S. (2006) Ecology based integrated management of sweet potato weevil in India. Acta Horticulture 703: 127–136.
- Palaniswami M.S. and Mohandas N. (1992) Search for resistance to *Cylas formicarius* F. in sweet potato. Journal of Root Crops 18: 41–52.
- Palaniswami M.S. and Mohandas N. 1993. Studies on attractants of adult weevils of *Cylas formicarius* F. Journal of Root crops 20(1): 44–52.
- Palaniswami M.S., Mohandas N. and Visalakshi A. (1992) An integrated package for sweet potato weevil (*Cylas formicarius* F.) management. Journal of Root Crops 18:113–119.
- Pillai K.S., Rajamma P and Palaniswami M.S. (1993) New technique in the control of sweet potato weevil using synthetic sex pheromone in India. International Journal of Pest Management 39: 84–89.
- Rajamanikyam M., Gadea S., Vadlapudia V., Parvathanenic P.S., Koudea D., Dommatib K. A., Tiwarib K.A., Misraa S., Sripadid P., Amanchya R. and Upadhyayula M.S. (2017) Biophysical and biochemical characterization of active secondary metabolites from *Aspergillus allahabadii*. Process Biochemistry 56: 45–56.
- Reddy V.P.G. and Chi H. (2015) Demographic comparison of sweetpotato weevil reared on a major host, *Ipomoea batatas*, and an alternative host, *I. triloba*. Scientific Reports 5(11871):1–9. doi:10.1038/srep11871.
- Sharma S., Mishra P. and Patni V. (2018) Analysis of volatile constituents in normal flower and insect induced flower gall of *Crataeva religiosa* Journal of Pharmacognosy and Phytochemistry 7(1): 2667–2673.
- Sripathi R., Jayagopal D. and Ravi S. (2017) A study on the seasonal variation of the essential oil composition from *Plectranthus hadiensis* and its antibacterial activity. Natural Product Research 32(7): 871–874. doi: 10.1080/14786419.2017.1363748.
- Stevenson P.C., Muyinza H.H., David R.P., Elaine A.F., Dudley I., Talwana, H. and Mwanga R.O.M. (2009) Chemical basis for resistance in sweetpotato *Ipomoea batatas* to the sweet potato weevil *Cylas puncticollis*. Pure and Applied Chemistry 81: 141–151. doi: 10.1351/PAC-CON-08-02-10.
- Subin M.P. and Jagathy K.V. (2017) Preliminary phytochemical screening and GC-MS analysis in the methanolic leaf extracts of *Polyalthia korinti* (Dunal) Benth. & Hook & Thomson.

- World Journal of Pharmaceutical Research 6 (6): 1225–1237.
- Suluvoy J.K and Grace B.V.M. (2017) Phytochemical profile and free radical nitric oxide (NO) scavenging activity of *Averrhoa bilimbi* L. fruit extract 3. Biotech 7: 85.
- Sutherland J.A. (1986) A review of the biology and control of the sweet potato weevil, *Cylas formicarius elegantulus* (F). Tropical Pest Management 32: 304–315.
- Takahashi M., Isoi K., Kimura Y. and Yoshikura M. (1964) Studies on the components of *Panax ginseng* C.A. Meyer. Journal of the Pharmaceutical Society of Japan 84: 757–759.
- Talekar NS. 1987. Resistance in sweet potato to sweet potato weevil. Insect Science and its Application 8: 819–823.
- Teoh Y.P. and Don M.M. (2015) Effect of Temperature on *Schizophyllum commune* Growth and 4H-pyran-4-one,2,3-dihydro-3, 5-dihydroxy-6-methyl- Production using a Bubble Column Bioreactor Chiang Mai Journal of Science 42(3): 539–548.
- Uritaini I., Saito T., Honda H. and Kim W.K. (1975) Induction of furano terpenoids in sweet potato roots by the larval components of the sweet potato weevils. Agricultural and Biological Chemistry 37:1875–1862.
- Vasuki B., Chitra P., Vijayabaskaran M., Mahadevan N. and Sambathkumar R. (2022) Gas chromatography-mass spectrometry analysis of *Hugonia mystax* leaves International Journal of Pharmaceutical Sciences and Research 13(1): 409–416.
- Vos M.D. and Jander G. (2008) Choice and no-choice assays for testing the resistance of *Arabidopsis thaliana* to chewing insects. Journal of Visualized Experiments 15: 683.

(Received August 24, 2023; revised ms accepted November 17, 2023; published December 31, 2023)

Synthesis and characterization of silver nanoparticles using *Datura metel* L. (Solanaceae) leaf extract and its larvicidal activity on *Epilachna vigintioctopunctata* F.

C. Sundareswari* and D.N.P. Sudarmani

Post-graduate and Research Department of Zoology, Ayya Nadar Janaki Ammal College
(Autonomous, affiliated to Madurai Kamaraj University), Sivakasi 626123, Tamil Nadu, India.
Email: sundareswaric@gmail.com

ABSTRACT: Insecticidal activities of synthesized silver nanoparticles of leaf extract of *Datura metel* L. (Solanaceae) (DM) against grubs *Epilachna vigintioctopunctata* F. (Coleoptera, Coccinellidae) at varying levels of concentrations was evaluated. DM leaf extract was used to create AgNPs, and nanoparticle production could be seen after six hours. UV-vis spectrophotometer, Particle size analyzer, FTIR and SEM analysis were used to confirm the synthesis of AgNPs. GCMS spectra of leaf extract of DM showed 20 substances, of which nine were known s phytochemicals and the others were unidentified. UV-visible spectra to analyse the Surface Plasmon Resonance for AgNPs revealed in the range of 366 - 374nm. LC₅₀ values for the AgNPs synthesized leaf extracts calculated 24 h after treatment against the fourth instar larvae of *E. vigintioctopunctata* using probit analysis revealed the LC₅₀ of the aqueous leaf extract as 252.31ppm and that of AgNPs synthesized leaf extract as 396.09ppm. When comparing the aqueous and AgNPs synthesized leaf extracts, AgNPs nanoparticles synthesized leaf extracts were more efficient as larvicide than aqueous leaf extract. © 2023 Association for Advancement of Entomology

KEY WORDS: AgNPs, aqueous leaf extract, larvicidal activity, LC₅₀ values

INTRODUCTION

A common and well-liked vegetable crop growing in the subtropics and tropics is eggplant, *Solanum melongena* L. (Sarkar *et al.*, 2006) and an important vegetable crop in India. According to their size, shape, and color, there are numerous varieties of brinjal available in India (Nisha *et al.*, 2009). The significant losses caused by damage to different agricultural crops are largely attributable to insect pests. According to estimates, insect pests

are responsible for 23 per cent of all agricultural losses (Agarwal, 2011). Pesticides are used often and to control these pests in the vegetable fields, which has led to widespread resistance development, unfavorable impacts on non-target organisms, the presence of hazardous residues in food, and environmental and health risks (Kranthi *et al.*, 2002). All these have highlighted the requirement to create unique, risk-free, and environmentally friendly pest management methods. *Epilachna vigintioctopunctata* F. (Coleoptera,

* Author for correspondence

Coccinellidae) is a serious pest of solanaceous crops, especially potatoes and aborigines, throughout its range. Adult and grubs feed on the surface of leaves by scraping away the surface cells between the main veins to leave irregular-shaped holes or strips. Heavy feeding damages the leaves, giving them a skeletonized or lace-like appearance. The damaged leaves turn brown and curl as they dry before falling off. *Datura metel* L. (Solanaceae) is a shrub-like, commonly found in India and has pesticide and medicinal properties. An attempt was made to investigate *D. metel* in the creation of silver nanoparticles from its leaf extract and its efficiency as larvicide at varying concentrations on *E. vigintioctopunctata*.

MATERIALS AND METHODS

Leaves of *D. metel* were gathered in and around Meenampatti, Sivakasi Taluk, between July 2021 and March 2022. The study region is located between latitudes of 9°27'2.6424" from the north and longitudes of 77°48'26.0496" from the east. *D. metel* leaves plucked were cleaned in tap water and allowed to air dry for 5-7 days in the shade. Using an electric blender, the air-dried plant components were powdered. For extraction, 10g of fine leaf powder was collected in a beaker with 100ml of double-distilled water that had been sterilized. After that, it was heated at 60°C for 3 number 1, and the extract was kept at -20°C and used within a week.

Phytochemical analysis: Agilent GC 7890A/MS 5975C and a capillary column Agilent DB5MS were used in the GCMS to evaluate the *D. metel* leaf extract. The mass spectrometer was tuned to 70 eV, and the computer library created by WILEY7, NIST05, and NIST05s was used to identify unknown substances using probability-based matching.

Synthesis of silver nanoparticles from leaf extract: The precursor used to create silver nanoparticles was silver nitrate. Analytical-grade silver nitrate (AgNO_3) in the amount of 16.961mg was measured out and mixed with 90ml of Milli-Q water. In a 1 Erlenmeyer flask, 90ml of produced 1mM aqueous AgNO_3 solution was added to 10ml

of aqueous leaf extract and incubated at room temperature in the dark. AgNPs were being produced as a result of the transition from light yellow to dark brown (Lingarao and Savithramma, 2013).

Characterization of nanoparticles: Utilizing UV-Visible spectroscopy, the initial characterization of AgNPs was completed. Synthesized silver nanoparticles were subjected to FTIR analysis, particle size analysis and SEM.

Epilachna beetle rearing: A mass culture of *E. vigintioctopunctata* was kept in a lab at the Department of Zoology, Ayya Nadar Janaki Ammal College, Sivakasi, Tamilnadu, India, using brinjal leaves as feed, in order to consistently obtain a large number of epilachna larvae for experimental usage. The *E. vigintioctopunctata* infested brinjal plant was grown in the nearby agrifarm of Sivakasi, where the adults of *E. vigintioctopunctata* were collected. The laboratory reared epilachna larvae were used for bioassays and the cultures were maintained throughout the study period. The adults were allowed to breed in the laboratory condition, when the larvae hatch from the egg they were fed with fresh brinjal leaves. They were permitted to develop till becoming third instar stages, and then were used to assess the larvicidal efficiency.

Larvicidal activity: The leaf extract's larvicidal activity was tested in the laboratory settings. By dilution with de-ionized water, various amounts of aqueous and synthetic silver nanoparticle leaf extract were created. Each extract had a concentration of 100, 50, 25, 12, 5 and 6.25 mg ml⁻¹. To examine leaf extract's larvicidal potential, a spray test was conducted. Different concentrations of extracts were made, sprayed on brinjal leaves, and given to epilachna beetle larvae as feed. To determine the larvicidal activity of each concentration of extracts, ten larvae per concentration were subjected and replicated thrice. Similar to this, three replicates of control group (tap water) for each test was done. After a 24h exposure period, the mortality rate was observed. For each concentration, the percentage of larval mortality from the replicates was calculated.

Statistical analysis: Larval mortality analysis assessed using standard deviation and mean separation. Using the program SPSS, 2007, the probit analysis for LC_{50} , the upper and lower 95 per cent confidence bounds were determined.

RESULTS AND DISCUSSION

Aqueous *D. metel* leaf extract was used to perform green production of silver nanoparticles. The initial confirmation of the biosynthesis of silver nanoparticles using aqueous leaf extract was the visible observation of colour change. The initial colour of the leaf extract suspension was pale yellow; but, after the addition of silver nitrate and an overnight incubation at room temperature, the colour changed to brown (Fig. 1).

GCMS analysis of leaf extract: As many as 20 compounds were detected in GCMS spectra of *D. metel* leaf extract, of which nine were recognized. The remaining 12 substances were unidentified. Retention times, mass spectra, and a library of typical compounds were compared to determine whether phytochemicals were present. Among the nine compounds, L-Arabinitol ($C_5H_{12}O_5$) had 40.57 per cent peak area, with a retention time of 4.887 minutes, having a molecular weight of $152.15 \text{ g mol}^{-1}$. The remaining compounds were found in the range of 1 – 7 per cent peak area (Table 1).

Characterization of leaf extract: The Surface Plasmon Resonance (SPR) is analyzed using an ultraviolet-visible spectrum. AgNPs UV-visible spectra fell between 366 and 374 nm (Fig. 2a).

Fourier Transform-Infrared (FTIR) analysis:

FTIR analysis showed the vibrational spectra of AgNPs synthesized leaf extract (Table 2). The alcohol molecule found in the extract was responsible for distinctive peak that occurred at 3453.31. The peak at 760.87 was assigned to the stretching vibration C-CL group. The alkenes group is present in the AgNPs produced extract as indicated by the 1385.76 and 1639.38. Presence of aromatic compound is established by the peak at 1512.09. Alkanes group presence is confirmed by the absorption peaks at 2063.69 and 2885.31 by C-H stretch and C-H stretch respectively. The peak at 2309.6 indicates the presence of alkynes in the AgNPs synthesized extract with C≡C stretch in AgNPs synthesized leaf extracts (Fig. 2b).

Particles size analysis: Dynamic light scattering (DLS) was used to establish the intensity-weighted mean diameter (Z-average) of the particle size of DM-AgNPs in the range of 0.5-1 nm (Fig. 2c). The particle size and shape of the biosynthesized DM-AgNPs are also determined using SEM examination. The SEM investigation revealed that the produced nanoparticles' mean particle diameters ranged from 20 to 27 nm (Fig. 2d).

Larvicidal activity: LC_{50} values for the AgNPs synthesized leaf extracts calculated 24 h after treatment against the fourth instar larvae of *E. vigintioctopunctata* using probit analysis revealed the LC_{50} of the aqueous leaf extract as 252.31 ppm and that of AgNPs synthesized leaf extract as 396.09 ppm. Larval mortality improved with

Table 1. GCMS spectra of identified phytochemicals from the extract of *Datura metel*

No	RT (Min)	Compound	Peak area (%)	Molecular formula	Molecular weight ($\text{g} \cdot \text{mol}^{-1}$)
1	4.887	L-Arabinitol	40.57	$C_5H_{12}O_5$	152.15
2	5.542	Propanamide	3.41	C_3H_7NO	73.095
3	5.709	Benzeneacetaldehyde	7.09	C_8H_8O	120.1485
4	7.398	Benzoic acid	1.22	$C_7H_6O_2$	122.12
5	8.464	Silane	4.77	H_4Si	32.117
6	11.264	o-Cyanobenzoic acid	1.98	$C_8H_5NO_2$	147.130
7	12.230	Pinacolyl ethyl phosphonofluoridate	2.31	$C_8H_{18}FO_2P$	196.199

Table 2. Functional groups detected in AgNPs synthesized leaf extract of *D. metel* as revealed by FTIR

No	Absorption (cm ⁻¹)	Class of compounds	Bond
1	743.51	Alkyl halide	C-CL stretch
2	1385.76	Alkenes	C-H bend
3	1512.09	Aromatic	C=C stretch
4	1639.38	Alkenes	C=C stretch
5	2063.69	Alkanes	C-H stretch
6	2309.60	Alkynes	Ca=C stretch
7	2885.31	Alkanes	C-H stretch
8	3453.31	Alcohol	O-H stretch


Fig. 1 Visible colour change of *Datura metel* leaf extract after silver nanoparticles synthesis

Fig. 2 Characterization of Silver Nanoparticles synthesized (a) UV-Visible Spectral analysis; (b) FTIR analysis; (c) Particle size analysis; (d) SEM analysis

increase in the concentration of silver nanoparticles synthesized leaf extract (Table 3). AgNPs synthesized leaf extract @100 concentration recorded 80 per cent mortality; at the same the aqueous leaf extract showed 10 per cent mortality. When comparing the aqueous and AgNPs synthesized leaf extracts, AgNPs nanoparticles synthesized leaf extracts were more efficient than aqueous leaf extract as larvicide.

Due to the pressing needs to create ecologically friendly technology, biosynthesis of nanoparticles has received a lot of attention. A significant method for creating many types of nanoparticles, such as

copper, iron, platinum, silver, and zinc, has been the biosynthesis of nanoparticles utilizing biological agents (Rasheed *et al.*, 2017; Sharon *et al.*, 2018). Bioresearch in the domain of explaining the mechanism of plant-mediated nanoparticle production holds great promise (Kumar and Yadav, 2009). The synthesis of AgNPs using *D. metel* was verified in the current work utilizing a variety of cutting-edge methods. When *D. metel* leaf extract is added to the silver nitrate solution, the mixture changes color from pale yellow to brown due to the reduction of the silver ion, indicating the creation of silver nanoparticles. The current work provides proof that the extract of *D. metel* has a potential to

Table 3. Silver nanoparticles synthesized leaf extract on the mortality of IVth instar larvae of *E. vigintioctopunctata* (@10 larvae/ conc)

Concentration	Mean mortality of IV th instar larvae in	
	aqueous	AgNPs
6.25	0	1
12.5	0	2
25	0	2
50	0	5
100	1	8

reduce silver ions (Ag⁺ into Ag⁰) and convert silver nitrate to silver nanoparticles while also having larvicidal effects on epilachna beetles. Similar research on *Polyalthia longifolia* samples, whose colour ranges from nearly colorless to brown, was published by Kaviya *et al.*, in 2011.

By comparing mass spectra, retention times, and a library of common compounds, it was possible to confirm that leaf extract contained phytochemical components. In GCMS spectra of *D. metel* leaf extract 20 compounds were detected, of which nine were recognized as phytochemicals. Mishra and Patnaik (2020) obtained comparable findings from a methanol extract of the complete *Withania somnifera* plant. According to the size and polydispersity of NPs, the distinctive peak of AgNPs is located about 430nm (Anandalakshmi *et al.*, 2016). According to Rajagopal *et al.* (2021), the UV-Vis spectra of the CuNPs produced using *Wrightia tinctoria* displayed absorption peak maxima at 357 nm. Plants create phytochemicals either through their primary or secondary metabolism. The majority of the time, they are biologically active in the plant host and aid in plant growth or defense against pests, diseases, or predators.

Measurements using the Fourier transform infrared spectroscopy technique are used to pinpoint the potential biomolecules in charge of the reduction, capping, and effective stability of silver nanoparticles (Padalia *et al.*, 2015). It was demonstrated that NPs were generated by the presence of functional groups like alcohol, halides,

alkanes, alcohol, and aromatic compounds (Prabha *et al.*, 2022).

The larvicidal activity of silver nanoparticles made from *D. metel*'s aqueous extract against the epilachna beetle was observed in this work. The LC₅₀ values calculated in the present study revealed that both aqueous and AgNPs have remarkable larvicidal effect on epilachna beetle. According to Islam *et al.* (2011), the LC₅₀ values for three medicinal plants were 18.40, 23.70, and 29.61 per cent, for the epilachna phytophagous pest. When bioefficacy of two indigenous plant products, namely seed extracts of *Strychnos nuxvomica* and *Pachyrrhizus erosus*, and two entomopathogenic fungi, *Beauveria bassiana* and *Metarrhizium anisopliae*, were tested against the epilachna beetle on bottle gourd, the population of the epilachna beetle was significantly reduced (Vishwakarma *et al.*, 2011). According to Ahmed (2007), spraying the castor plant *Ricinus communis* aqueous extract on sunflower, *Helianthus annulus* foliage and capitula, reduced epilachna attacks resulting improved the oil seed harvest.

The present findings clearly show that at certain levels, the suggested green silver nanoparticles have an effect on *E. vigintioctopunctata* that causes mortality. Ag⁺ ions are produced from the nanoparticle's surface by oxidation processes, and when they enter an insect's physiological processes, they interact with biological components (such as insect proteins) and induce toxicity (Park *et al.*, 2010). AgNPs are also known to bind with thiol groups in proteins and promote their denaturation, which causes the death of larvae (Johnston *et al.*, 2010). Hence, *D. metel*'s aqueous leaf extract and silver nanoparticles produced leaf extract can be effectively used as larvicide to *E. vigintioctopunctata*.

ACKNOWLEDGEMENT

The authors thank the Principal and Management of Ayya Nadar Janaki Ammal College (Autonomous), Sivakasi for the financial support of the project through the Ayya Nadar Janaki Ammal Research Fund and for providing research facilities in carrying out the work.

REFERENCES

- Agarwal R.G. (2011) Pesticides consumption pattern. In: *50 Years of Agrochemicals and India's Struggle Towards Food and National Security*. Dhanuka Agritech Limited. 1-94pp.
- Ahmed K.N. (2007) Effect of biopesticides in the suppression of major insect pests and yield of oilseed crops under different climatic conditions in Bangladesh. Proceedings of National Symposium on Plant Protection-Technology Interface, Association for Advancement of Plant Protection, Bidhan Chandra Krishi Viswavidyalaya, Nadia, West Bengal, India. 2007 December 28-29, pp81-82.
- Anandalakshmi K., Venugobal J. and Ramasamy V. (2016) Characterization of silver nanoparticles by green synthesis method using *Petalium murex* leaf extract and their antibacterial activity. Applied Nanoscience 6: 399–408
- Islam K., Islam M.S., Ferdousi Z. (2011) Control of *Epilachna vigintioctopunctata* Fab. (Coleoptera: Coccinellidae) using some indigenous plant extracts. Journal of Life and Earth Science 6: 75–80.
- Johnston H.J., Hutchison G., Christensen F.M., Peters S. and Hankin S. (2010) A review of the *in vivo* and *in vitro* toxicity of silver and gold particulates: particle attributes and biological mechanisms responsible for the observed toxicity. Critical Reviews in Toxicology 40 (4): 328–46.
- Kaviya K., Santhanalakshmi J. and Viswanathan B. (2011) Green synthesis of silver nanoparticles using *Polyalthia longifolia* leaf extract along with D-Sorbitol: Study of antibacterial activity. Journal of Nanotechnology 7: 84–91
- Kumar V. and Yadav S.K. (2009) Plant-mediated synthesis of silver and gold nanoparticles and their applications. Journal Chemical Technology and Biotechnology 84: 151–157.
- Lingarao M. and Savithramma N. (2013) Isolation and characterization of -Sitostrol from leaf methanol extract of *SvensoniaHyderabadensis* – A rare medicinal plant taxon. International Journal of Natural Product Science 3(1): 1–7.
- Mishra D. and Patnaik S. (2020) GC-MS Analysed Phyto-Chemicals and antibacterial activity of *Withania somnifera* (L.) dunal extract in the context of treatment to *Liver Cirrhosis*. Biomedical & Pharmacology Journal 13(1): 71–78.
- Nisha P., Nazar A.P. and Jayamurthy P. (2009) A comparative study on antioxidant activities of different varieties of *Solanum melongena*. Food and Chemical Toxicology 47: 2640–2644.
- Padalia H., Moteriya P. and Chanda S. (2015) Green synthesis of silver nanoparticles from marigold flower and its synergistic antimicrobial potential. Arabian Journal of Chemistry 8: 732–741
- Park E.J., Yi J., Kim Y., Choi K. and Park K. (2010) Silver nanoparticles induce cytotoxicity by a Trojan-horse type mechanism. Toxicology 24 (3): 872–878.
- Prabha I., Nivetha A. and Sakthivel C. (2022) Effective/ comparative investigation on green mediated nano copper oxide: fabrication, characterization and environmental applications. Materials today: Proceedings 51(4): 1690–1695. doi:10.1016/j.matpr.2020.10.024
- Rajagopal G., Nivetha A., Ilango S., muthudevi G.P., Prabha I. and Arthimanju R. (2021) Phytofabrication of selenium nanoparticles using *Azolla pinnata*: evaluation of catalytic properties in oxidation, antioxidant and antimicrobial activities, Journal of Environmental Chemical and Engineering 9: 105483.
- Rasheed T., Bilal M., Iqbal H.M.N. and Li C. (2017) Green biosynthesis of silver nanoparticles using leaves extract of *Artemisia vulgaris* and their potential biomedical applications. Colloids Surf B Biointerfaces 158: 408–415.
- Sarker R.H., Yesmin S. and Hoque M.I. (2006) Multiple shoot formation in eggplant (*Solanum melongena* L.). Plant Tissue Culture Biotech 16: 53–61.
- Sharon E.A., Velayutham K. and Ramanibai R. (2018) Biosynthesis of copper nanoparticles using *Artocarpus heterophyllus* against dengue vector *Aedes aegypti*. International Journal of Life Science and Scientific Research 4(4): 1873–1879. doi:10.21276/ijlssr.2018.4.4.4
- Vishwakarma R., Prasad P.H., Ghatak S.S. and Mondal S. (2011) Bio-efficacy of plant extracts and entomopathogenic fungi against *Epilachna* beetle, *Henosepilachna vigintioctopunctata* (Fabricius) infesting bottle gourd. Journal of Insect Science 24(1): 65–70.

Redescription of *Carbula indica* (Westwood, 1837) (Hemiptera, Heteroptera, Pentatomidae) from West Bengal, India with a key to the Indian species of the genus *Carbula* Stål, 1865

Amartya Pal*, Swetapadma Dash, Devanshu Gupta and P.C. Saha

Zoological Survey of India, M-Block, New Alipore, Kolkata 700053, West Bengal, India.

Email: amartyapal08@gmail.com, oasisnainital@gmail.com, devanshuguptagb4102@gmail.com, sahapradip1974@gmail.com

ABSTRACT: *Carbula indica* (Westwood, 1837) (Hemiptera, Heteroptera, Pentatomidae) re-described with illustrations based on the material collected from the Himalayan hills of Darjeeling, West Bengal, India. New characters are included, along with both male and female genitalia and morphometric measurements to facilitate easy species determination. A key to the Indian species of *Carbula* Stål, 1865 is also presented. © 2023 Association for Advancement of Entomology

KEYWORDS: Taxonomy, morphometrics, genitalia, species determination, key

INTRODUCTION

The genus *Carbula* Stål, 1865, is distributed in the Oriental and Afrotropical regions, represented by 76 species worldwide, of which ten species are known from India (Ravneet Kaur *et al.*, 2013). *Carbula* can be recognized by the following characters: body dorsally ovate and ventrally convex; head rounded or somewhat truncated at apex; mandibular plates equal in length and pronotum with anterior lateral angles obtuse; genital capsule quadrangular and paramere with bilobed crown. Westwood (1837) placed *Carbula indica* under the genus *Pentatoma*, Distant (1902) later transferred it to the genus *Carbula* Stål. Both described the species based on external coloration and a few morphological features, which are insufficient to identify the species correctly. This

species was reported from West Bengal (Darjeeling), Sikkim and Nepal by Distant (1902); later reported from China (Rider *et al.*, 2002). During a collection of Himalayan Pentatomidae, four specimens (2 males, 2 females) of *C. indica* from Darjeeling were found and this opportunity is taken to re-describe and illustrate the species, including male and female genitalia and measurements to facilitate easy species identification. A key to the Indian species of *Carbula* Stål, 1865 is also provided.

MATERIALS AND METHODS

The material for the present study includes dry-pinned specimens housed at the Hemiptera Section of the Zoological Survey of India, Kolkata, West Bengal. The male genitalia was dissected following

* Author for correspondence

the method given by Ahmad (1986). After boiling the whole abdomen in hot water for about 10–15 minutes, the female genitalia was dissected with potassium hydroxide (KOH - 10%). The internal contents were cleared after thoroughly washing it in distilled water 2–3 times. With the help of fine forceps, the terminalia and spermatheca were detached from the abdominal ventrites. The measurements are given in millimeter and presented as median, with minimum and maximum values given in bracket. The following dimensions were measured: Body length (from apex of mandibular plates to apex of membrane, dorsal view), head length (from apex of mandibular plates to anterior margin of pronotum, dorsal view), head width (width of head including compound eyes, dorsal view), interocular width (between inner margins of compound eyes, dorsal view), length of each antennal segment, length of each segment of rostrum, pronotum length (medially, from anterior to posterior margin of pronotum, dorsal view), pronotum width (maximum width between humeri in dorsal view), scutellum length (medially from base to apex) and scutellum width (maximum width at base between basal angles of scutellum). Morphological terminology used for male and female genitalia broadly follows Salini (2019). The photographs were taken under a Leica M205A stereomicroscope using a Leica DMC-4500 camera. The photographs were processed in LAS V4.12 software for morphometry. Photographs were edited using Adobe Photoshop CS (Version 8.0).

Abbreviations used: Vlf: Valvifers, Lt: Laterotergite, P: Paramere, PA: Processes of aedeagus, CP: Conjunctival Processes, PT: Phallosome.

RESULTS AND DISCUSSION

Carbula indica (Westwood, 1837) [Figs. 1-16]

Pentatoma indica Westwood, 1837: 42.

C. fusca Distant, 1887: 346.

C. indica: Distant, 1902: 171.

Type locality: INDIA: West Bengal: Darjeeling: Kurseong

Material examined: 2 male, 2 female, Lepchajagat, Darjeeling, West Bengal, 4.vi.1975, Coll. J. K. Jonathan.

Redescription:

Coloration: Dark brownish yellow (Figs. 1, 12); anterior lateral margins of pronotum yellowish-brown; first three segments of antennae yellowish-brown, segment- IV slightly darker brown and apical two-third portion of last segment blackish. Abdomen pale brown with a broad central black fascia, spiracles and small marginal spots, black; rostrum blackish-brown with apex black; legs yellowish-brown with black punctation; genital capsule, pale brown and blackly punctate.

Head: Wider (across eyes) than long (1.911 : 1.503 mm) (Fig. 3); antennae five-segmented, segment IV longest, segment V longer than II, III, segment I smallest; rostrum long, four-segmented, passing metacoxae reaching up to II abdominal segment (Fig. 15) but sometimes up to anterior margin of III abdominal segment; segment II longest.


Thorax: Pronotum hexagonal, wider than longer, its width is 2.5 times than its length and about 2.5 times wider than head, anterolateral margins slightly serrated, lateral margins smooth, humeral angles rounded (Fig. 3).

Scutellum: Slightly longer than broad at base (3.747: 3.714mm) (Figs. 1, 10), apex rounded; hemelytral membrane passing tip of abdomen (Fig. 2).

Legs: With fine hairs and blackly punctate. Tibiae longer and slender than respective femora. Tarsi hairy, second tarsal segment shortest.

Abdomen: Abdomen broader than longer (6.396: 4.946) and with a broad central black fascia (Fig. 4).

Male genitalia: *Pygophore*. Longer than broad (2.136: 1.804mm), medially, in dorsal view (Fig. 7); postero-lateral lobes (= caudal lobes) tumescent (Figs. 5, 6), medial area slightly elevated (Fig. 5), subpentagonal in shape (Figs. 6, 7), Dorsal rim widely excavated with a narrow median notch




Figs. 1–11. *Carbula indica* (Westwood, 1837), male. 1, dorsal; 2, ventral; 3, head; 4, abdomen; 5, apex of abdomen (ventral); 6, pygophore (ventral); 7, pygophore (dorsal); 8, phallus (dorsal); 9, phallus (ventral); 10, phallus (lateral); 11, paramere. P–Paramere, PA–Processes of aedeagus (=Penial lobes). CP–Conjunctival Processes, PT–Phallotheca.

which is associated with convex lobe-like outgrowth nearly resembling the 1+1 triangular tooth-like structure on the infoldings of dorsal rim (Fig. 7). There is a shallow roughly W-shaped notch at the medial portion of ventral rim (Fig. 6). **Phallus.** Phallotheca of phallus (Figs. 8, 9) is barrel-shaped, slightly sclerotized without any process; one pair of slightly sclerotized processes of aedeagus (=Penial lobes), encircled by transparent, membranous conjunctival processes (Fig. 10); aedeagus long, tube-like and much longer than processes of aedeagus (=Penial lobes).

Paramere. Moderately sclerotized, crown bilobed (Fig. 11), upper lobe slender; lower lobe broader, spoon-shaped with a small projection at proximal end, slightly curved upwards facing towards upper lobe.

Female genitalia: Valvifers VIII sub-triangular with inner lateral margin curved (Fig. 14); valvifers VIII ventrally not meeting each other; valvifers IX fused to single quadrate plate with posterior margin concave, laterotergite VIII posteriorly encompassing laterotergite IX (Fig. 14);



Figs. 12–16. *Carbula indica* (Westwood, 1837), female. 12, dorsal; 13, ventral; 14, Terminalia; 15, rostrum; 16, Spermatheca. Vifs–Valvifers, Lt–Laterotergite

spermatheca (Fig. 16) with proximal spermathecal duct longer than distal spermathecal duct, median dilation present but not balloon-like; intermediate part of spermatheca (=spermathecal pump) short, only single flange (proximal flange) visible; apical receptacle large or-bicular, with a finger-like process.

Measurements:

Males (n = 2); median (minimum–maximum). Body length 7.050 (6.798–7.303); head: length 1.503

(1.496–1.510), width (across eyes) 1.911 (1.897–1.925), interocular width 1.183 (1.177–1.189); lengths of antennal segments: I—0.280 (0.271–0.289), II—0.756 (0.747–0.765), III—0.728 (0.721–0.735), IV—0.931 (0.926–0.936), V—0.839 (0.833–0.845); lengths of segments of rostrum: I—1.549 (1.536–1.562), II—1.680 (1.600–1.761), III—0.800 (0.800–0.801), IV—0.668 (0.626–0.711); pronotum: length 1.942 (1.938–1.946), width: 4.879 (4.861–4.897); scutellum: length 3.276 (2.805–3.747), width (at basal angles) 3.247 (2.781–3.714).

Females (n = 2); median (minimum–maximum). Body length 7.367(7.359–7.375); head: length 1.693 (1.665–1.721), width (across eyes) 2.034 (1.956–2.112), interocular width 1.230 (1.155–1.251); lengths of antennal segments: I—0.358 (0.352–0.364), II—0.755 (0.745–0.765), III—0.767 (0.765–0.769), IV—0.936 (0.929–0.943), V—0.898 (0.885–0.911); lengths of segments of rostrum: I—1.585 (1.566–1.604), II—1.734 (1.707–1.762), III—0.796 (0.792–0.801), IV—0.642 (0.628–0.656); pronotum: length 2.107 (1.992–2.222), width: 5.420 (5.284–5.556); scutellum: length 3.304 (2.861–3.748), width (at basal angles) 3.297 (2.849–3.746).

Distribution: India: West Bengal (Darjeeling, Kurseong) and Sikkim. *Elsewhere:* Nepal (Distant, 1902) and China (Rider *et al.*, 2002).

Key to species of *Carbula* Stål, 1865 from India

1. Humeral angles spinously produced.....2
 - Humeral angles obtusely angulate.....5
2. Upwardly and forwardly directed humeral angles.....3
 - Slightly backwardly directed humeral angles with subacute spines...***biguttata* (Fabricius, 1794)**
3. Scutellum densely punctate, upper lobe of parameres slightly angulate apically ***aliena* Distant, 1918**
 - Scutellum sparingly and coarsely punctate, upper lobe of parameres apically nearly broadly rounded.....4
4. Scutellum with a Y-shaped luteous marking at middle portion and punctate except each basal angle and the apex..... ***scutellata* Distant, 1887**
 - Scutellum brownish-ochraceous without Y-shaped luteous marking and uniformly punctate except two basal angle and apex ***aspavia* Distant, 1908**
5. Mandibular plates equal to clypeus.....6
 - Mandibular plates longer than clypeus

.....***rugulosa* Distant, 1902**

6. Median continuous black stripe on each of the abdominal segments; rostrum passing metacoxae reaching up to II abdominal segment but sometimes up to anterior margin of III abdominal segment, humeral angle less produced with apices rounded

.....***indica* (Westwood, 1837)**

- Median black stripe only on two terminal abdominal segments; rostrum just passing metacoxae reaching up to II abdominal segment, humeral angle broadly produced with apices obtuse and slightly paler and levigate7

7. Scutellum brownish-ochraceous, uniformly punctate and without any luteous spots at each basal angle and apex ..***crassiventris* (Dallas, 1849)**

- Scutellum luteous, sparingly punctate with a large luteous spot at each basal angle and apex.....8

8. Antennae luteous.....***socia* (Walker, 1867)**

- Antenna with the first, second, and third joints ochraceous, fourth and fifth black with their bases ochraceous9

9. Antero-lateral margins of pronotum smooth, luteous levigate.....***producta* Distant, 1901**

- Antero-lateral margins of pronotum crenulated, luteous levigate***insocia* (Walker, 1868)**

The *C. indica* specimens were compared with the photographs of paratype of *C. indica* held by Natural History Museum, London, thus enabled us to redescribe the species in greater detail. *C. indica* is closely related to *C. crassiventris* on the basis of morphological data, during re-examination of the both species were found to have some minute morphological differences between these two species. Re-examination revealed that humeral angle of pronotum less protruded in *C. indica*; first three antennal segment of *C. indica* yellowish-brown, segment- IV slightly darker brown and apical two-third portion of last segment blackish, while in *C. crassiventris* whole segments of antennae reddish brown; in *C. indicia* adomen pale brown with a broad central black band, while in *C.*

crassiventris there was a more or less distinct central spot on each of the two terminal abdominal segments. It is important to note that this study will help to facilitate the taxonomic study of genus *Carbula* as the published information on this species is limited. Earlier work cited above only provided information on external coloration and a few morphological features, while this study has provided the details of morphology of this species, in the form of digital photographs of both male and female habitus and genitalia, for the first time.

ACKNOWLEDGEMENTS

The authors are grateful to Dr. Dhriti Banerjee, Director, Zoological Survey of India for encouragements and laboratory facilities and to Dr. C. Raghunathan, Divisional-in-charge, Entomology Division, Zoological Survey of India for his valuable suggestions and support. The authors are also thankful to all the officers and staffs of Hemiptera section for their helpful co-operation and to Ms. Debika Bhunia, Zoological Survey of India for helping in preparing the photo plates. The authors acknowledge Dr. Santhamma Salini, ICAR–National Bureau of Agricultural Insect Resources, Bengaluru, India, for the information regarding *Carbula* in the preparation of key.

REFERENCES

- Ahmad I. (1986) A fool-proof technique for inflation of male genitalia in Hemiptera (Insecta). *Pakistan Journal of Entomology* 1(2): 111–112.
- Dallas W.S. (1849) Notice of some hemipterous insects from Boutan (East Indies), with descriptions of the new species. *Transactions of the Entomological Society of London* 5: 186–194.
- Distant W.L. (1887) Contributions to a knowledge of Oriental Rhynchota. Part 1. Fam. Pentatomidae. *Transactions of the Entomological Society of London* 3: 341–359.
- Distant W.L. (1901) Undescribed Indian Rhynchota: Pentatomidae. *Entomologist* 34: 346–347.
- Distant W.L. (1902) The fauna of British India including Ceylon and Burma, Rhynchota. Vol. 1. Taylor and Francis, London. 438pp.
- Distant W.L. (1908) The fauna of British India including Ceylon and Burma, Rhynchota Vol. IV. Taylor and Francis, London. 15+501pp.
- Distant W. L. (1918) The fauna of British India including Ceylon and Burma, Rhynchota Vol.VII. Taylor and Francis, London. 7+210pp.
- Fabricius J.C. (1794) *Entomologia systematica emendata et aucta secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus*. C.G. Proft, Hafniae 4: 472. https://digitalcommons.usu.edu/bee_lab_er/220
- Kaur R., Singh D., and Kaur H. (2013) Taxonomic Significance of External Genitalia in differentiating four Species of Genus *Carbula* Stål (Heteroptera, Pentatomidae) from North India. *Journal of Entomology and Zoology Studies* 1: 33–42.
- Rider D.A., Zheng L.Y. and Kerzhner I.M. (2002) Checklist and nomenclatural notes on the Chinese Pentatomidae (Heteroptera). II. Pentatominae. *Zoosystematica Rossica* 11(1): 135–153.
- Salini S. (2019) Pentatomidae (Hemiptera, Heteroptera, Pentatomoidea) of India. In: *Indian Insects Diversity and Science* (Eds. Ramani S., Mohanraj P. and Yeshwanth H.M.), CRC Press, Taylor & Francis Group, Boca Raton. pp121–146.
- Stål C. (1865) *Hemiptera Africana*. Norstedtiana, Stockholm. iv + 256pp.
- Walker F. (1867) Catalogue of the specimens of heteropterous Hemiptera in the collection of the British Museum. Part II. E. Newman, London. pp241–417.
- Walker F. (1868) Catalogue of the specimens of Hemiptera, Heteroptera in the collection of the British Museum. Part III. E. Newman, London. pp418–599.
- Westwood J.O. (1837) A catalogue of Hemiptera in the collection of the Rev. F. W. Hope, M.A. with short Latin diagnoses of the new species. Printed by J. C. Bridgewater, London. Pt. 1, pp46.

Evaluation of the insecticidal action of polyphenolic compounds from *Streblus asper* (Lour.) on the red cotton bug, *Dysdercus cingulatus* (Fab.) (Hemiptera, Pyrrhocoridae)

L. Anila^{1*} and M.S. Hashim²

¹Department of Biochemistry, NSS College, Nilamel, Kollam 691535, Kerala, India.

²Dr.Hashim's Institute of Life Sciences, Karyavattom, Thiruvananthapuram 695581, Kerala, India.
Email: dranilaleelamma@gmail.com

ABSTRACT: The study aims to evaluate oxidative stress and the activity of acetylcholinesterase (AChE), upon applying polyphenolic bioinsecticide isolated from *Streblus asper* (PBSA) at a concentration of 0.595 µg/insect (LD₅₀) by topical application on *Dysdercus cingulatus* Fabricius (Red cotton bug- Hemiptera, Pyrrhocoridae). The results demonstrated that the active fraction exhibited significant inhibition in activities of AChE, antioxidant enzymes and Glutathione-S-transferase (GST) and a significant increase in the lipid peroxides (MDA/ TBARS) which led to the fact that *D. cingulatus* became more susceptible to the tested PBSA. The study has provided basic information on the mechanism of action of PBSA that will be promising to develop effective alternatives to synthetic insecticides.

© 2023 Association for Advancement of Entomology

KEYWORDS: Bioinsecticide, mechanism of action, acetylcholinesterase, antioxidant enzymes, lipid peroxides

INTRODUCTION

Cotton (*Gossypium hirsutum* L.) plant is a shrub and is widely cultivated of great economic importance in over 90 countries for its natural fiber and secondly for seeds (Chaudhry, 2010). The red cotton bug, *Dysdercus cingulatus* F. (Heteroptera, Pyrrhocoridae) is an important pest of cotton. Although synthetic chemical insecticides can control it, the side effects are enormous (Vennila *et al.*, 2000). Pollution of the environment by pesticides has been increasing due to their use to manage various pests. Bioinsecticides are highly effective, safe, and ecologically acceptable (Arya *et al.*, 2022). Recent emphasis is on the use of natural pesticides, which are usually of plant origin. Unlike

synthetic chemical pesticides, which leave harmful residues in the aquatic environment (Aktar *et al.*, 2009; Mahmood *et al.*, 2016) bioinsecticides are biodegradable, specific in action (harmless to non-target organisms), and also possess the ability to counter pest resistance issues caused by synthetic pesticides (Mishra *et al.*, 2020).

Streblus asper Lour (Family: Moraceae) is a small tree which is indigenous to tropical countries such as India, Sri Lanka, Malaysia, the Philippines and Thailand (Glasby, 1991). It is a well-known ethnomedicinal plant which is also used in Ayurveda (Singh and Singh, 1987; Singh and Ram, 1988). It finds place in the Ayurvedic Pharmacopoeia of India and an up-to-date and comprehensive review of

* Author for correspondence

S. asper that covers its traditional and folk medicinal uses, phytochemistry and pharmacology (Rastogi *et al.*, 2006). Preliminary study has reported that polyphenolic extracts from the stem bark of *S. asper* possess insecticidal activity against the fifth instar of *D. cingulatus* (Hashim and Devi, 2003) and its partially purified polyphenolic bioinsecticide from *S. asper* (PBSA) had significant effects in the mortality of newly emerged fifth instar *D. cingulatus* (Anila and Hashim, 2022). A study on the mechanism of insecticidal activity of the most active PBSA in *D. cingulatus* by analyzing oxidative stress and inhibition of acetylcholinesterase activity upon topical application using its LD₅₀ concentration was taken up.

MATERIALS AND METHODS

The red cotton bugs, were obtained from the laboratory maintained under controlled conditions (temp 28-30°C, RH 95 ± 2 % and a photoperiod of light 12 h; dark 12 h) by feeding soaked cottonseeds. Newly emerged fifth instar insects were used for the experiments. Each treatment contains three replications and fifteen insects were used for each replication.

The stem bark of the plant, *S. asper* was collected from Nagarcoil Forest (Tamil Nadu, India) and was authentically identified. Polyphenolic compounds were extracted from *S. asper* according to the procedure Hashim and Devi (2003). Two compounds were maximum insecticidal activity. The compound I with maximum insecticidal activity was identified as flavanone compound family (Anila and Hashim, 2022). PBSA was used for evaluation of mechanism of action of insecticidal activity by analyzing oxidative stress and inhibition of AChE activity upon its application on *D. cingulatus*.

Newly emerged fifth instar *D. cingulatus* (15 days after of molting) were used and insects were divided into two groups containing 12 each. Ethanol (40%) was topically applied to group I which served as control and PBSA @ 0.595 µg/insect (LC₅₀) dissolved in ethanol (40%) topically applied to the group II insects which served as test. After 12 and 24 hours exposure the haemolymph (40µl) was

collected directly into a polystyrene tube containing few crystals of phenylthiourea by making a tiny incision in the antennae of *D. cingulatus* from each group. Hemocytes were removed from the haemolymph by centrifugation at 10000 rpm for 15 min. Separated haemolymph was used for various analyses. The whole brain was minced separately and homogenized with normal saline and centrifuged for 10 minutes at 3000 rpm and the supernatant was used for various analyses. Homogenates (as described) were performed in pools of 12 insects and every enzyme activity determination was the average of 6 independent pools of 12 insects (Daffre and Faye, 1997).

AchE activity was done by the method of Ellman *et al.* (1961). The activity was expressed as Units per milligram protein where 1 Unit = nanomoles of thiocholine liberated per minute.

Catalase (CAT) activity was measured (Maehly and Chance, 1954). The estimation was done spectrophotometrically following the decrease in absorbance at 230 nm. The specific activity is expressed in terms of Units/ mg protein where 1 Unit = velocity constant per second.

The measurement of superoxide dismutase (SOD) involves generation of superoxide radical by photoreduction of riboflavin and its detection by nitrite formation from hydroxylamine hydrochloride at 543 nm (Das *et al.*, 2000). SOD activity was expressed in Units/mg protein where 1 Unit = enzyme concentration required to inhibit OD at 560 nm of chromogen produced by 50 per cent in 1 minute.

Glutathione-S-transferase (GST) activity was measured spectrophotometrically by the method of Habig *et al.* (1974) using S-2,4-dinitrophenyl glutathione (CDNB) as a substrate. The principle of the method is based on measurement of the conjugation of S-2,4-dinitrophenyl glutathione (CDNB) with reduced glutathione. The formation of adduct of CDBN, S-2,4-dinitrophenyl glutathione was monitored by measuring the net increase in absorbance at 340 nm against the blank. The activity of GST was expressed in terms of l mol/min/mg protein.

The glutathione content (GSH) was determined as described by the improved method of Benke *et al.* (1974). The quantity of reduced glutathione was expressed in mg/ g protein or mg/ dl haemolymph.

Thiobarbituric acid-reacting substances (TBARS) were estimated by the method of Niehaus and Samuelsson (1968).

Protein contents of supernatant were determined after TCA (trichloro acetic acid) precipitation (Lowry, 1951) using bovine serum albumin (BSA) as the standard protein. The protein was measured at 670 nm absorbance in a spectrophotometer. Statistical significance was determined by one way Analysis of Variance (ANOVA) in SPSS 20.0 package. The data given in figures are expressed as mean \pm SEM, for n = 6 experiments.

RESULTS AND DISCUSSION

Mechanism of insecticidal action of PBSA on AChE activity:

The control insects without any insecticide exposure exhibited an acetylcholinesterase activity of 28 units/mg protein in haemolymph. After 12 h of PBSA exposure the enzyme activity significantly reduced to 12.95 units/mg protein and after 24 h of exposure again the activity significantly decreased to 7.89 units/mg protein. Similar pattern of inhibition was observed in enzyme activity in brain of bioinsecticide treated insects. Activity of AChE was significantly inhibited in both haemolymph and brain of red cotton bugs treated with PBSA @0.595 μ g/insect (LD_{50}) after 12 and 24 h of topical application when compared to control insects (Table 1). The activity

Table 1. Activity of AChE in haemolymph and brain of *Dysdercus cingulatus* exposed to PBSA

Biochemical parameters studied	After 12 hours				After 24 hours			
	Haemolymph		Brain		Haemolymph		Brain	
	Control	Test	Control	Test	Control	Test	Control	Test
AChE (U ¹)	28.16 \pm 0.803	12.95 ^a \pm 0.584	42.36 \pm 1.75	27.96 ^a \pm 1.02	27.53 \pm 0.679	7.89 ^{ab} \pm 0.5511	42.84 \pm 1.27	20.33 ^{ab} \pm 0.694

¹Unit = nanomoles of thiocholine liberated per minute per milligram protein; Values expressed as mean \pm SEM, for n = 6 experiments. ^atest group is compared to control group at p = 0.05. ^btest group after 12 hours is compared to test group after 24 hours at p = 0.05

Table 2. Activity of antioxidant enzymes in haemolymph and brain of *Dysdercus cingulatus* exposed to PBSA

Biochemical parameters studied	After 12 hours				After 24 hours			
	Haemolymph		Brain		Haemolymph		Brain	
	Control	Test	Control	Test	Control	Test	Control	Test
SOD (U ²)	3.17 \pm 0.150	1.85 ^a \pm 0.072	5.39 \pm 0.179	3.17 ^a \pm 0.117	3.08 \pm 0.078	1.16 ^{ab} \pm 0.046	5.46 \pm 0.194	2.26 ^{ab} \pm 0.086
CAT (U ³)	16.58 \pm 0.358	8.96 ^a \pm 0.142	32.47 \pm 0.909	25.49 ^a \pm 0.610	16.44 \pm 0.252	6.32 ^{ab} \pm 0.157	33.54 \pm 0.708	18.67 ^{ab} \pm 0.424
GST (U ⁴)	3.46 \pm 0.096	1.84 ^a \pm 0.045	6.31 \pm 0.117	4.25 ^a \pm 0.080	3.22 \pm 0.061	0.924 ^{ab} \pm 0.029	6.43 \pm 0.104	3.09 ^{ab} \pm 0.070

Values expressed as mean \pm SEM, for n = 6 experiments; ^atest group is compared to control group at p = 0.05; ^btest group after 12 hours is compared to test group after 24 hours at p = 0.05. ² Unit = enzyme concentration required to inhibit OD at 560 nm of chromogen produced by 50 % in 1 minute. ³ Unit = velocity constant/ second; ⁴ Unit = m M of CDNB utilized /min/mg protein

was significantly inhibited after 24 hours of topical application when compared to 12 h exposure.

Activity of catalase and superoxide dismutase (SOD) was significantly inhibited in both 12 and 24 h time intervals after the topical application of PBSA @ 0.595 µg/insect (LD₅₀) and the inhibition was more pronounced in 24 h exposure in both haemolymph and brain. There was a significant reduction in GST activity in the haemolymph and brain of insects after the topical application of PBSA at 0.595 µg/insect (LD₅₀) for 12 and 24 h time intervals when compared to control insects (Table 2). The inhibition was more pronounced in the haemolymph and brain after 24 h exposure when compared to insects treated for 12 h.

Duration dependent decrease in GSH content was observed in both haemolymph and brain of PBSA at 0.595 µg/insect (LD₅₀) treated insects when compared to control insects. The decrease was higher in 24 h treated insects when compared to 12 h treated insects. TBARS content was significantly increased in haemolymph and brain after 12 and 24 h exposure of PBSA at 0.595 µg/insect (LD₅₀) treated insects when compared to control group (Table 3). There was significant increase in TBARS level in 24 h treated insects when compared to 12 h.

In the present study the polyphenolic bioinsecticide exposure have been demonstrated to reduce the activity of acetyl cholinesterase significantly in red cotton bugs. The results are in agreement with

Maazoun *et al.* (2017) who reported that *Urginea maritima* bulbs extract exhibited inhibitory AChE activity in *Sitophilus oryzae* (L.). Pesticide-induced oxidative stress is the final manifestation of a multi-step pathway, resulting in an imbalance between pro-oxidant and antioxidant defense mechanisms. Concomitantly, pesticide intoxication induces a derangement of certain antioxidant mechanisms in different tissues, including alterations in antioxidant enzymes and the glutathione redox system (Banerjee *et al.*, 2001). Therefore the attempt to analyze the antioxidant status and measure the activity of AChE in red cotton bugs exposed to PBSA which belongs to a flavanone family, showed a significant inhibition in the activity of AChE and antioxidant enzymes.

Reactive oxygen species (ROS), such as superoxide anions (O₂⁻) and H₂O₂ are produced throughout the cells during normal aerobic metabolism. The intracellular concentration of ROS is a consequence of both their production and their removal by various antioxidants. A major component of antioxidant system in mammalian cells consists of three enzymes, SOD, CAT and (glutathione peroxidase) GPX. These enzymes work in concert to detoxify O₂ and H₂O₂ in cells. It has been established that many pesticides are capable of inducing oxidative stress by overwhelming or modulating cellular drug metabolizing systems (Sule *et al.*, 2022).

Oxidative stress occurs when there is an imbalance between free radical generation and antioxidant

Table 3. Concentration of reduced glutathione (GSH) and TBARS content in haemolymph and brain of *Dysdercus cingulatus* exposed to PBSA

Biochemical parameters studied	After 12 hours				After 24 hours			
	Haemolymph		Brain		Haemolymph		Brain	
	Control	Test	Control	Test	Control	Test	Control	Test
MDA (TBARS)	0.925 ± 0.047	1.73 ^a ± 0.050	1.42 ± 0.053	2.13 ^a ± 0.072	0.930 ± 0.042	2.24 ^{ab} ± 0.075	1.42 ± 0.061	2.55 ^{ab} ± 0.068
GSH	16.58 ± 0.592	9.47 ^a ± 0.376	24.63 ± 0.581	12.38 ^a ± 0.314	17.16 ± 0.551	6.18 ^{ab} ± 0.330	25.24 ± 0.778	8.59 ^{ab} ± 0.424

Values expressed as mean ± SEM, for n = 6 experiments; ^atest group is compared to control group at p = 0.05; ^btest group after 12 hours is compared to test group after 24 hours at p = 0.05

defenses. It often results in severe pathological consequences, such as membrane disruption, DNA damage and protein damage and cytotoxicity (Saini *et al.*, 2023). The activities of the antioxidant enzymes like superoxide dismutase (SOD), glutathione peroxidase (GSH-Px) and glutathione-S-transferase (GST) were decreased by fenitrothion incubation. The same treatment reduced the level of antioxidant glutathione (GSH). The activities of glutathione-S-transferase (GST) and gamma-glutamyl transpeptidase (gamma-GT) were more affected by fenitrothion and endosulfan, respectively, indicating oxidative stress (El-Shenawy, 2010). These support the present findings that the significant decrease in antioxidant enzymes in haemolymph and brain of bioinsecticide-treated insects when compared to control.

Recent reports showed that there were significantly reduced GSH levels in all tissues after methiocarb administration in experimental animals (Ozden *et al.*, 2009) and methomyl decreased AChE, superoxide dismutase (SOD) and glutathione S-transferase (GST) activities and increased level of lipid peroxidation (LPO) (Mansour *et al.*, 2009). These reports are in support with the present findings that the phenolic compound isolated from *Streblus asper* (PBSA) exhibited the insecticidal activity as evidenced by its inhibitory effect on the activities of AChE and antioxidant enzymes and significant increase in the level of lipid peroxidation.

In conclusion, polyphenolic bioinsecticide from *Streblus asper* (PBSA) has significant effects on newly emerged fifth instar *D. cingulatus*, and they caused increased mortality in a duration-dependent manner. The AChE activity and antioxidant status are impaired after the exposure of PBSA to red cotton bugs. The abnormal change in antioxidant status and acetylcholinesterase activity in cotton bugs may be the reason for the insecticidal action. The compound PBSA may therefore serve as an effective alternative to conventional insecticides in controlling red cotton bugs.

REFERENCES

Aktar W., Dwaipayan S. and Ashim C. (2009) Impact of Pesticides Use in Agriculture: Their Benefits and

Hazards. *Interdisciplinary Toxicology* 2 (1): 1.

Anila L. and Hashim M.S. (2022) Insecticidal Activity of a Bioinsecticide from Stem Bark of *Streblus asper* on Red Cotton Bug (*Dysdercus cingulatus*). *Research Journal of Agricultural Sciences* 13 (2): 535–539.

Arya S., Ravendra Kumar, Om Prakash, Avneesh Rawat, and Pant A.K. (2022) Impact of Insecticides on Soil and Environment and Their Management Strategies. In: *Agrochemicals in Soil and Environment: Impacts and Remediation*, Springer. pp213–230.

Banerjee B.D., Vandana S. and Rafat S.A. (2001) Pesticide-Induced Oxidative Stress: Perspective and Trends. *Reviews on Environmental Health* 16 (1): 1–40.

Benke G.M., Cheever K.L., Miler F.E. and Murphy S.D. (1974) Comparative Toxicity, Anticholinesterase Action and Metabolism of Methyl Parathion and Parathion in Sunfish and Mice. *Toxicology and Applied Pharmacology* 28 (1): 97–109.

Chaudhry M.R. (2010) Cotton Production and Processing. In: *Industrial Applications of Natural Fibres: Structure, Properties and Technical Applications*. John Wiley and Sons Ltd. pp219–236.

Daffre S. and Faye I. (1997) Lipopolysaccharide Interaction with Hemolin, an Insect Member of the Ig-Superfamily. *FEBS Letters* 408 (2): 127–130.

Das K., Luna S. and Chainy G.B.N. (2000) A Modified Spectrophotometric Assay of Superoxide Dismutase Using Nitrite Formation by Superoxide Radicals. *Indian journal of biochemistry & biophysics* 37: 201–204

El-Shenawy N.S. (2010) Effects of Insecticides Fenitrothion, Endosulfan and Abamectin on Antioxidant Parameters of Isolated Rat Hepatocytes. *Toxicology in Vitro* 24 (4): 1148–1157.

Ellman G.L., Diane Courtney K., Valentino Andres Jr, and Robert M.F. (1961) A New and Rapid Colorimetric Determination of Acetylcholinesterase Activity. *Biochemical Pharmacology* 7 (2): 88–95.

Glasby J. S. (1991) *Directory of Plants Containing Secondary Metabolites*. CRC Press, London. pp488. doi:10.1201/b12561

Habig W.H., Michael J. P. and William B.J. (1974) Glutathione S-Transferases: The First Enzymatic Step in Mercapturic Acid Formation. *Journal of*

- Biological Chemistry 249 (22): 7130–7139.
- Hashim M.S. and Devi K.S. (2003) Insecticidal Action of the Polyphenolic Rich Fractions from the Stem Bark of *Streblus asper* on *Dysdercus cingulatus*. Fitoterapia 74 (7–8): 670–676.
- Lowry O.H. (1951) Protein Measurement with the Folin Phenol Reagent. Journal of Biological Chemistry 193: 265–275.
- Maazoun A., Ben Hlel T., Haouel S., Belhadj F., Mediouni B.J.J. and Marzouki M.N. (2017) Screening for insecticidal potential and acetylcholinesterase activity inhibition of *Urginea maritima* bulbs extract for the control of *Sitophilus oryzae* (L.). Journal of Asia-Pacific Entomology 20 (3): 752–760. doi: 10.1016/j.aspen.2017.04.004
- Mahmood I., Sameen R.I., Kanwal S., Alvina G. and Khalid R. H. (2016) Effects of Pesticides on Environment. In: Plant, Soil and Microbes: Volume 1: Implications in Crop Science, Springer International. pp253–269. doi: 10.1007/978-3-319-27455-3_13
- Mansour, S.A., Abdel-Tawab H.M. and Tarek M.H. (2009) Effects of Methomyl on Lipid Peroxidation and Antioxidant Enzymes in Rat Erythrocytes: In Vitro Studies. Toxicology and Industrial Health 25 (8): 557–63.
- Mishra J., Venkatesh D. and Naveen Kumar A. (2020) Biopesticides in India: Technology and Sustainability Linkages. 3 Biotech 10 (5): 210. doi: 10.1007/s13205-020-02192-7
- Niehaus Jr W.G. and Samuelsson B. (1968) Formation of Malonaldehyde from Phospholipid Arachidonate during Microsomal Lipid Peroxidation. European Journal of Biochemistry 6 (1): 126–130.
- Ozden S., Betul C., Selda G., Pelin A., Sehnaz B. and Buket A. (2009) Methiocarb-Induced Oxidative Damage Following Subacute Exposure and the Protective Effects of Vitamin E and Taurine in Rats. Food and Chemical Toxicology 47 (7): 1676–1684.
- Rastogi S., Dinesh K.K. and Ajay Kumar S.R. (2006) *Streblus asper* Lour.(Shakhotaka): A Review of Its Chemical, Pharmacological and Ethnomedicinal Properties. Evidence-Based Complementary and Alternative Medicine 3 (2): 217–222.
- Saini R., Himaani M., Tanisha G. and Neena K.D. (2023) Dietary Antioxidants and Their Potential Role in Human Disease Management. Current Nutrition & Food Science 19 (3): 262–281.
- Singh N.P. and Ram E.R. (1988) Filaria and Its Herbal Cure. New Botanist 15 (4): 201–205.
- Singh N.P. and Singh V.K. (1987) *Streblus asper* Lour.—an Ancient Indian Drug for the Cure of Filariasis. Acta Botanica Indica 15: 108–109.
- Sule R.O., Liam Condon and Aldrin V.G. (2022) A Common Feature of Pesticides: Oxidative Stress—the Role of Oxidative Stress in Pesticide-Induced Toxicity. Oxidative Medicine and Cellular Longevity 2022:5563759. doi: 10.1155/2022/5563759
- Vennila S., Ramasundram P., Sheo Raj and Kairon M.S. (2000) Cotton IPM and Its Current Status. ICAR - Central Institute for Cotton Research, Technical Bulletin 8: 1–13.

Bioefficacy of *Tagetes minuta* L. against *Aphis craccivora* Koch (Hemiptera, Aphididae)

V.V. Ashish*, Haseena Bhaskar*, U. Sreelatha[#] and K.B. Deepthy*

*Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Thrissur 680656, Kerala, India.

[#]Department of Floriculture and Landscaping, College of Agriculture, Kerala Agricultural University, Thrissur 680656, Kerala, India.

Email: ashishvv98@gmail.com, haseena.bhaskar@kau.in

ABSTRACT: Efficacy of solvent fractions of *Tagetes minuta* L. (hexane, chloroform and aqueous fractions of aerial parts) was evaluated against *Aphis craccivora* Koch at five different concentrations (0.025, 0.05, 0.1, 0.15 and 0.2%) in the laboratory. At higher concentrations, all the fractions recorded significant reduction in aphid count; maximum reduction after 120h of treatment was in hexane fraction at 0.2 per cent (90.00%), followed by chloroform fraction at 0.2 per cent (86.67%) and aqueous fraction at 0.2 per cent (85.00%). The best concentrations of the three solvent fractions of *T. minuta* identified in the laboratory were evaluated against the aphid on cowpea, in a pot culture experiment. By 10th day of treatment, all the fractions at 2 per cent concentration reduced population of aphids, with the highest reduction in hexane fraction. © 2023 Association for Advancement of Entomology

KEYWORDS: Mexican marigold, solvent fractions, hexane fraction, aphid management

INTRODUCTION

The cowpea aphid, *Aphis craccivora* Koch (Hemiptera, Aphididae) is an important sucking pest of leguminous crops throughout India (Jagdish *et al.*, 2011). It is a serious pest of cowpea, feeding on aerial parts of the plant and leading to significant yield loss (Veeranna and Adivappar *et al.*, 2019). It also causes indirect damage to the plant by excreting copious amount of honey dew, which leads to the development of sooty mould on the surface of leaves, thereby reducing the rate of photosynthesis; besides, it is also a vector of important viral diseases affecting the legumes (Ghosh *et al.*, 2017). Synthetic insecticides are

being extensively used for aphid management in vegetable crops in India. The popular chemical insecticides used against aphids include neonicotinoids like imidacloprid, acetamiprid and thiamethoxam. The excessive and indiscriminate use of synthetic pesticides for their management has, however, led to the development of resistance in the populations of aphids. *A. craccivora* has been reported to exhibit considerable degree of resistance to newer insecticides like imidacloprid (Dawood and Farghaly, 2016) and dinotefuran (Mokbel and Mohamed, 2009), among many others. Hence, there is a need for identifying alternative, environmentally benign, effective, and biodegradable pesticides with greater selectivity against the aphid pest. In this

* Author for correspondence

context, botanical pesticides are ideal alternatives for synthetic pesticides due to their rapid degradation, target specific nature and less phytotoxicity (Vasquez *et al.*, 2016). *Tagetes minuta* L., commonly known as Mexican marigold is an annual herb that belongs to the family, Asteraceae. The biological activities of the plant have been documented worldwide. Insecticidal effects of *T. minuta* have been studied against several pests of field crops, stored products and public health (Perich *et al.*, 1995; Shahzadi *et al.*, 2010; Phoofole *et al.*, 2013). Few studies have been conducted in India, particularly in Kerala, to evaluate the insecticidal effects of this plant; hence, a study was conducted to identify the insecticidal effects of the solvent fractions of *T. minuta* on *A. craccivora*.

MATERIALS AND METHODS

Adults of *A. craccivora* collected from the culture maintained, in the polyhouse of Department of Agricultural Entomology, College of Agriculture, Vellanikkara, Kerala, India were used to initiate the laboratory culture. The aphids were allowed to infest cowpea seedlings grown in pro trays maintained in rearing cages in the laboratory. New seedlings were allowed to be infested periodically to maintain the culture. In order to obtain uniform aged adult aphids for the study, a few seedlings from the culture were selected and all adults were removed, retaining the nymphs. On the next day, the freshly moulted adults were collected from those seedlings and used for the study.

Preparation of solvent fractions: *T. minuta* plants, cultivated in the pots, in the Department of Floriculture and Landscaping, College of Agriculture, Vellanikkara, Kerala, India were used for the study. The above-ground parts of the plant were harvested at the flowering stage, shade dried and pulverized. The botanical was then extracted sequentially, based on the polarity of the solvents into three separate fractions, using hexane (non-polar), chloroform (medium polar) and water (highly polar). The aerial parts of the plant were dried and pulverized material was weighed (100g each conical flask) and extracted using hexane (3 times of volume) by placing in a rotary shaker for 48h. The

solution was then filtered and the filtrate was concentrated by a rotary evaporator at 40°C, to obtain the hexane fraction. The residue obtained during filtration was re-extracted sequentially using chloroform, followed by water, following the same procedure as hexane extraction, to obtain chloroform fraction and aqueous fraction, respectively (Auamcharoen and Chandrapatya, 2015). The three solvent fractions were used for bioefficacy studies against *A. craccivora*.

Laboratory bioassay: The hexane, chloroform and aqueous fractions of *T. minuta* were evaluated at five different concentrations (0.025, 0.05, 0.1, 0.15 and 0.2 %) for efficacy against *A. craccivora*. The treatment concentrations were prepared by dissolving the required quantity of each fraction in distilled water along with the emulsifier, Triton X-100 (20µl 50ml⁻¹). Cowpea seeds were sown in individual paper cups (one seed per cup) filled with potting mixture (soil and vermicompost in 1:1 ratio) and the seeds were allowed to sprout and grow. At four leaf stage, adult aphids from the laboratory culture were released using a camel hair brush at the rate of 20 per plant and allowed to settle. The seedlings were then sprayed with appropriate treatments using a hand atomizer (3ml per seedling). Three replications were maintained for each concentration of the fractions. Seedlings sprayed with water plus emulsifier, served as control. An absolute control was also maintained without any treatment. The experiment was laid out in Completely Randomized Design (CRD) with 17 treatments and three replications. Nymphs produced viviparously if any, were removed from the experimental seedlings periodically, to avoid any experimental error due to population build up. Number of adult aphids on the plants was recorded at 24, 48, 72, 96 and 120h of spraying and reduction in aphid count was calculated. Data on reduction of aphid was subjected to analysis of variance using the software, GRAPES 1.0.0, developed by Kerala Agricultural University.

Evaluation of fraction in pot culture: A pot culture experiment was carried out to evaluate the efficacy of the most effective concentration of each fraction of *T. minuta* against *A. craccivora* in cowpea, during July - September, 2022. The best

concentration each of hexane (0.2%), chloroform (0.2%) and aqueous (0.2%) fractions were selected for evaluation based on their efficacy in the laboratory study against the adult aphids. The efficacy of the selected fractions was compared with those of neem oil emulsion (2%), azadirachtin (1% EC, 3ml L⁻¹) and horticultural mineral oil (2%). An untreated control was also maintained. The experiment was laid out in CRD with seven treatments, each replicated thrice with eight plants per replication.

The seeds of cowpea variety - Bhagyalakshmi (bush type) were sown in individual polybag (35cm x 20cm x 20cm) filled with potting mixture (soil, coir pith compost and cow dung in 2:1:1 ratio). All cultural practices were carried out as per Package of Practices Recommendations, KAU (2016). Aphids were allowed to infest the 20-day-old cowpea plants by keeping aphid infested cowpea seedlings in paper cups, at the base of each plant in polybag. Treatments were imposed 20 days after the release of aphids, using a hand sprayer. For recording the aphid population in treatments, three plants were selected randomly from each treatment replication. Number of aphids (both nymphs and adults) was recorded from three shoot bits of 5cm length per plant, excised from three tender shoots (Bindu, 1997). Pre-treatment count one day prior to treatment and post treatment counts of aphids at 1, 3, 7 and 10 days after treatment (DAT) application were counted by dislodging the aphids from each shoot bit, on to a white paper.

The data on mean number of aphids (per 5cm shoot length) before and after treatments were subjected to analysis of covariance (ANCOVA) using the software, GRAPES 1.0.0. In order to accommodate the variations in pre-count, the transformed data were analysed by taking population counts prior to the first application as covariate and ANCOVA was done for observations at 1, 3, 7 and 10 DAT. The result obtained was subjected to LSD (Least Significance Difference Test). The mean per cent reduction in population was also worked out 10 DAT.

Qualitative phytochemical analysis: Qualitative analysis of hexane (nonpolar) and chloroform

(medium polar) fractions of *T. minuta*, was performed to determine the predominant phytochemical constituents in the fractions. However, aqueous (polar) fraction was not analysed as ample literature on the constitution of polar fractions of *T. minuta* is available. The hexane and chloroform fractions were subjected to GC-MS/MS analysis at Sophisticated Analytical Instrumentation Facility (SAIF), IIT, Mumbai and the major compounds present in the fractions were recorded.

RESULTS AND DISCUSION

Laboratory evaluation of solvent fractions:

There was a sudden decline in aphid count on treated seedlings up to 48h of treatment, after which the population reduction was less pronounced. All three solvent fractions recorded the maximum reduction in aphid count at the highest concentration (0.2%) tested. Within 24h of treatment the hexane fraction (@0.2%) recorded 61.67 per cent reduction, followed by chloroform (60%) and aqueous fractions (58.33%), which were on par with each other. At lower concentration (0.15%), hexane fraction and aqueous fraction also reduced aphid count (60 and 50% respectively) and were on par with the above treatments. After 48h, hexane fraction (@0.2%) resulted in as high as 83.33 per cent reduction. Chloroform fraction (0.2%) also showed significant reduction (78.33%), on par the hexane fraction. Other concentrations of solvent fractions also showed reduction in aphid count at 48h. From 48 to 120h of treatment, the reduction in aphid count was less prominent in all the treatments. After 120 h of treatment, there were significantly higher reduction in aphid count in hexane fraction (@0.2%), hexane fraction (@0.15%), chloroform fraction (@0.2%), chloroform fraction (@0.2%) and aqueous fraction (@0.2%) (recording 90.00, 86.67, 86.67, 85.00 and 85.00% respectively) and were on par with one another (Table 1).

Evaluation of fractions in pot culture: All the treatments reduced the aphid population considerably 10 DAT. Among the fractions, hexane fraction recorded the lowest population (11.18 aphids per 5cm of shoot) resulting in 78.95 per cent reduction in aphid population. The hexane fraction

Table 1. Laboratory evaluation on the bioefficacy of *Tagetes minuta* solvent fractions against *Aphis craccivora*

Treatment	Reduction in aphid count (%) – hours after treatment				
	24h	48h	72h	96h	120h
Hexane fraction 0.025 %	23.33 ^{fg} (28.67)	40.00 ^{hi} (39.21)	53.33 ^{gh} (46.92)	56.67 ^{gh} (48.84)	56.67 ^f (48.84)
Hexane fraction 0.05 %	40.00 ^{cde} (39.21)	53.33 ^{fg} (46.94)	60.00 ^{fg} (50.85)	65.00 ^{fg} (53.76)	65.00 ^{de} (53.76)
Hexane fraction 0.1%	43.33 ^{cd} (41.13)	63.33 ^e (52.78)	68.33 ^{def} (55.82)	75.00 ^{de} (60.07)	76.67 ^{bc} (61.15)
Hexane fraction 0.15 %	60.00 ^{ab} (50.79)	75.00 ^b (60.07)	76.67 ^{bcd} (61.15)	80.00 ^{cd} (63.55)	86.67 ^a (68.66)
Hexane fraction 0.2 %	61.67 ^a (51.81)	83.33 ^a (65.95)	86.67 ^a (68.66)	88.33 ^a (70.11)	90.00 ^a (71.57)
Chloroform fraction 0.025 %	30.00 ^{ef} (33.16)	48.33 ^{gh} (44.04)	55.00 ^{gh} (47.88)	56.67 ^{gh} (48.84)	58.33 ^{ef} (49.80)
Chloroform fraction 0.05 %	36.67 ^{de} (37.20)	65.00 ^{de} (53.76)	70.00 ^{de} (56.84)	71.67 ^{ef} (57.86)	73.33 ^{bc} (59.00)
Chloroform fraction 0.1 %	38.33 ^{cde} (38.19)	66.67 ^{cde} (54.75)	71.67 ^{cde} (57.86)	75.00 ^{de} (60.00)	76.67 ^{bc} (61.15)
Chloroform fraction 0.15 %	48.33 ^{bcd} (44.01)	73.33 ^{bc} (58.93)	80.00 ^{abc} (63.55)	81.67 ^{bcd} (64.81)	85.00 ^a (67.40)
Chloroform fraction 0.2 %	60.00 ^{ab} (50.79)	78.33 ^{ab} (62.29)	85.00 ^a (67.40)	86.67 ^{ab} (68.66)	86.67 ^a (68.66)
Aqueous fraction 0.025%	15.00 ^{gh} (22.29)	36.67 ⁱ (37.26)	45.00 ^h (42.12)	46.67 ⁱ (43.09)	48.33 ^g (44.04)
Aqueous fraction 0.05%	28.33 ^{ef} (32.14)	45.00 ^{ghi} (42.12)	53.33 ^{gh} (46.92)	55.00 ^{hi} (47.91)	58.33 ^{ef} (49.83)
Aqueous fraction 0.1%	38.33 ^{cde} (38.22)	61.67 ^{ef} (51.76)	65.00 ^{ef} (53.76)	70.00 ^{ef} (56.79)	70.00 ^{cd} (56.79)
Aqueous fraction 0.15%	50.00 ^{abc} (45.00)	63.33 ^e (52.74)	71.67 ^{cde} (57.86)	75.00 ^{de} (60.07)	78.33 ^b (62.29)
Aqueous fraction 0.2 %	58.33 ^{ab} (49.80)	71.67 ^{bcd} (57.86)	81.67 ^{ab} (65.00)	83.33 ^{abc} (66.15)	85.00 ^a (67.40)
Control (Water + emulsifier)	13.33 ^h (21.34)	13.33 ^j (21.34)	16.67 ⁱ (24.05)	18.33 ^j (25.31)	18.33 ^h (25.31)
Absolute control	1.67 ⁱ (4.31)	10.00 ^j (18.43)	11.67 ⁱ (19.89)	13.33 ^j (21.34)	13.33 ^h (21.34)
LSD(0.05)	7.329	4.881	5.789	4.974	4.568

Figures in parentheses are arc sine transformed values; Means followed by common letter(s) do not significantly differ at P=0.05%

was on par with the neem oil emulsion (7.57 aphids/ 5cm shoot) that reduced aphid population by 87.59 per cent. The chloroform and aqueous fractions were on par with each other, reducing aphid population by 72.43 and 68.20 per cent, respectively. Aqueous fraction recorded aphid population (23.11) on par with azadirachtin (25.74), whereas, chloroform fraction was superior to azadirachtin recording a lower population of 18.41 aphids per 5 cm of shoot (Table 2).

Phytochemical constituents in solvent fractions: Qualitative phytochemical analysis of the bio-active fractions viz., hexane and chloroform fractions GC-MS/MS identified the presence of 4 α -phorbol 12,13-didecanoate, 4-o-methylphorbol 12, 13- didecanoate and milbemycin b in hexane fraction and 4-o-methylphorbol 12, 13- didecanoate and milbemycin b in the chloroform fraction.

In the laboratory, on treatment application, the aphids moved away from the treated seedlings, suggesting

a strong repellent and/or antifeedant action by the solvent fractions of *T. minuta*. In the pot culture study, the solvent fractions could significantly reduce the aphid population on cowpea. The efficacy of the hexane fraction of *T. minuta* was found comparable to neem oil while that of chloroform and aqueous fractions was comparable to azadirachtin 1 EC, which are widely recommended in the management of aphids in various crops (Sarvaiya *et al.*, 2018).

Volatile constituents present in several botanicals are known to trigger sudden repellent action in the aphids (Dardouri *et al.*, 2019). Volatile compounds such as D-limonene, dihydro-tagetone, (E)-tagetone, (Z)-tagetone, (Z)-beta-ocimene and allo-ocimene were reported to be present in *T. minuta* and are known to have repellent properties against insects (Kimutai *et al.*, 2015). Polar fractions of *T. minuta*, extracted using polar solvents such as methanol and water were reported to be composed

Table 2. Effect of *Tagetes minuta* solvent fractions on *Aphis craccivora* in cowpea pot culture

Treatment	Mean live aphids/ 5cm shoot length at - DAT					Reduction (%)
	PTC*	1	3	7	10	
Hexane fraction 0.2 %	53.11 (7.28)	20.37 ^e (4.51)	15.29 ^d (3.91)	12.59 ^c (3.54)	11.18 ^d (3.33)	78.95
Chloroform fraction 0.2%	66.78 (8.17)	27.33 ^b (5.22)	23.33 ^{bc} (4.81)	19.69 ^b (4.39)	18.41 ^c (4.27)	72.43
Aqueous fraction 0.2%	72.67 (8.52)	30.04 ^b (5.48)	20.74 ^c (4.55)	18.63 ^b (4.31)	23.11 ^{bc} (4.81)	68.20
Neem oil emulsion 2%	61.18 (7.82)	11.89 ^d (3.45)	9.96 ^c (3.16)	7.93 ^d (2.81)	7.59 ^{de} (2.75)	87.59
Azadirachtin 1% EC (3 ml L ⁻¹)	80.37 (8.96)	29.00 ^b (5.38)	25.85 ^b (5.08)	24.63 ^b (4.95)	25.74 ^b (5.07)	67.97
Horticultural mineral oil 2%	33.93 (5.82)	8.45 ^c (2.90)	6.48 ^f (2.55)	6.38 ^d (2.52)	5.59 ^e (2.36)	83.52
Untreated control	43.74 (6.61)	50.41 ^a (7.09)	44.36 ^a (6.66)	51.22 ^a (7.15)	61.78 ^a (7.85)	—
LSD (0.05)	NS	0.413	0.478	0.715	0.590	—

*PTC=Pre-treatment count; #Reduction at 10 DAT (%); Figures in parentheses are square root transformed values; Means followed by common letter (s) do not significantly differ at P=0.05 %

of various carbohydrates, proteins, phenols, flavonoids, terpenoids and alkaloids (Rikisahedew, 2018). The terpenoid, D-limonene present in the polar fraction of *T. minuta* is known to possess significant insecticidal properties (Karr and Coats, 1987).

Qualitative phytochemical analysis of hexane and chloroform fractions in the present study showed phorbol esters as predominant compounds. The antifeedant properties of the phorbol esters against some insect pests have already been documented. Phorbol esters extracted from *Jatropha* oil were evaluated for insecticidal activity against the third instar larvae of *Spodoptera frugiperda* (JE Smith). At highest concentration of the phorbol esters (0.25 mg ml⁻¹ w/v), food consumption and relative growth rate of *S. frugiperda* larvae reduced (by 33 and 42% respectively), apart from exhibiting contact toxicity (Devappa *et al.*, 2012). Phorbol esters are also known to affect signal transduction pathways by acting on the biological membranes (Goel *et al.*, 2007). Among the two phorbol esters identified, the role of 4 α -phorbol 12, 13-didecanoate as a prominent TRPV (Transient receptor potential cation channel) agonist has been widely documented (Alexander *et al.*, 2012). Similarly, milbemycin b, identified in both hexane and chloroform fractions, is a GABA gated chloride channel agonist (Ozoe, 2012), like the avermectin group of insecticides. The insecticidal activities of this compound have been widely documented against several insects (Bobade, 2019).

There are also reports on the detrimental effects of volatile compounds present in *T. minuta* viz., α -caryophyllene, limonene and (Z) ocimene on the reproduction of aphids. Tomova *et al.* (2005) reported up to 100 per cent reduction in reproduction in three species of aphids, viz., *Acyrtosiphon pisum*, *Myzus persicae* and *Aulacorthum solani*, on exposure to these compounds. The efficacy of the *T. minuta* fractions against *A. craccivora* in the present study could be the combined effect of repellency, antifeedancy, and neurotoxicity and reproduction inhibition. Though insecticidal effects of *T. minuta* against different species of aphids have been documented from different parts of the world,

most of the studies evaluated the crude aqueous extracts of the botanical in the field. Limited studies were conducted to evaluate the solvent fractions of *T. minuta*. When Ali *et al.* (2019) evaluated the bio-efficacy of aqueous leaf extracts of *T. minuta* along with three other botanicals viz., *Calotropis procera*, *Argemone mexicana* and *Azadirachta indica* against the mustard aphid, *Lipaphis erysimi* on Indian mustard, *Brassica juncea*, at higher concentration (1: 2.5 g ml⁻¹), *T. minuta* reduced *L. erysimi* population by 96.38 per cent. Kora and Teshome (2016) evaluated the aqueous extracts of chilli, garlic, ginger and *T. minuta* for their insecticidal property against green pea aphid, *Acyrthosiphon pisum* in the field. The aphid population was reduced to zero in the plants treated with *T. minuta*, chilli and ginger. The above studies show that *T. minuta* extracts can reduce aphid population build up considerably, in the field. Phoofolo *et al.* (2013) had evaluated the aphidicidal activities of the crude extracts *T. minuta* against the cabbage aphid, *Brevicoryne brassicae*. A comparison was made on lethal and sub-lethal effects of the crude extracts from acetone, methanol, water and a mixture of acetone/methanol/water (7:7:1 v:v). The mixture produced the most toxic extract, followed by methanol and water; whereas acetone extract was the least toxic. The study also demonstrated that the crude extract of *T. minuta* obtained using water as a solvent is as effective as crude extracts from organic solvent systems in terms of efficacy against cabbage aphids. This observation corresponds to the results of the present study where all the three solvent fractions of *T. minuta* (hexane, chloroform and water) were effective in reducing the aphid population.

ACKNOWLEDGEMENT

The authors are thankful to All India Network Project on Agricultural Acarology and Kerala Agricultural University for providing necessary facilities and financial support.

REFERENCES

- Alexander R., Kerby A., Aubdool A.A., Power A.R., Grover S., Gentry C. and Grant A.D. (2013) 4 α -

- phorbol 12, 13-didecanoate activates cultured mouse dorsal root ganglia neurons independently of TRPV4. *British Journal of Pharmacology* 168(3): 761–772. doi: 10.1111/j.1476-5381.2012.02186.x
- Ali A., Rizvi P.Q. and Khan F.R. (2019) Bio-efficacy of some plant leaf extracts against mustard aphid, *Lipaphis erysimi* Kalt on Indian mustard, *Brassica juncea*. *Journal of Plant Protection Research* 50(2): 130–132.
- Auamcharoen W. and Chandrapatya A. (2015) Acaricidal and ovicidal efficacies of *Leucaena glauca* Benth. seed crude extracts on *Tetranychus urticae* Koch (Acari: Tetranychidae). *Journal of Biopesticides* 8(2): 68–81.
- Bindu (1997) Biocontrol of pests of vegetable cowpea. M.Sc. (Ag) thesis, Kerala Agricultural University, Thrissur 680656, Kerala, India.
- Bobade A.F. (2019) GC-MS Analysis of bioactive compound in ethanolic extract of *Pithecellobium dulce* leaves. *American Society of Plastic Surgeons* 3: 8–13.
- Dardouri T., Gautier H., Issa R., Costagliola G. and Gomez L. (2018) Repellence of *Myzus persicae* (Sulzer): evidence of two modes of action of volatiles from selected living aromatic plants: Repellent volatiles against *Myzus persicae*. *Pest Management Science* 75(6): 1571–1584. doi: 10.1002/ps.5271.
- Dawood A.I. and Farghaly S.F. (2016) Resistance to Confidor (imidacloprid) in Two Sap-Sucking Insects and Cross-Resistance to Several Insecticides. *Egyptian Academic Journal of Biological Sciences F, Toxicology and Pest Control* 8(1): 1–5.
- Devappa R.K., Angulo-Escalante M.A., Makkar H.P. and Becker K. (2012) Potential of using phorbol esters as an insecticide against *Spodoptera frugiperda*. *Industrial Crops and Products* 38: 50–53.
- Ghosh A., Chakrabarti S., Mandal B. and KrishnaKumar N.K. (2017) Aphids as vectors of the plant viruses in India. In: *A century of plant virology in India*, Springer, Singapore. pp515–536.
- Goel G., Makkar H.P., Francis G. and Becker K. (2007) Phorbol esters: structure, biological activity, and toxicity in animals. *International Journal of Toxicology* 26(4): 279–288.
- Jagdish J., Bhaskar H., Priya P. and Ranjith A.M. (2011) Dispersal pattern of *Aphis craccivora* Koch in cowpea in Kerala. *Research Journal of Agricultural Sciences* 2(3): 746–747.
- Karr L.L. and Coats J.R. (1988) Insecticidal properties of d-limonene. *Journal of Pesticide Science* 13(2): 287–290.
- KAU (2016) Package of Practices Recommendations: Crops (15th Ed.). Kerala Agricultural University, Thrissur: pp183–207.
- Kimutai A., Ngeiywa M., Njagi P.G.N., Mulaa M., Ingonga J., Nyamwamu L.B., Ombati C. and Ngumbi P. (2015) Effects of essential oils of lemon grass, *Cymbopogon citratus* and the Mexican marigold, *Tagetes minuta* on mortality and oviposition in adult sandflies, *Phlebotomus duboscqi*. *International Journal of Biomedical Research* 6(9): 679. doi:10.7439/ijbr.v6i9.2351.
- Kora D. and Teshome E. (2016) Field evaluation of some botanical extracts against the pea aphid *Acyrthosiphon pisum* (Homoptera: Aphididae) on field pea *Pisum sativum* L. *Journal of Entomology and Zoology Studies* 4(4): 336–339.
- Mokbel E.S.S. and Mohamed A.I. (2009) Development of resistance in field strain of *Aphis craccivora* to the dinotefuran insecticides from the new class neonicotinoids and its effect on some enzymes content. *Egyptian Academic Journal of Biological Sciences C, Physiology and Molecular Biology* 1(1): 65–69.
- Ozoe Y. (2012) Target receptors in the control of insect pests, Part I. by E. Cohen. *Advances in Insect Physiology* 44: 211–286.
- Perich M.J., Wells C., Bertsch W. and Tredway K.E. (1995) Isolation of the insecticidal components of *Tagetes minuta* (Compositae) against mosquito larvae and adults. *Journal of the American Mosquito Control Association* 11(3): 307–310.
- Phoofolo M.W., Mabaleha S. and Mekbib S.B. (2013) Laboratory assessment of insecticidal properties of *Tagetes minuta* crude extracts against *Brevicoryne brassicae* on cabbage. *Journal of Entomology and Nematology* 5(6): 70–76.
- Rikisahedew J.J. (2018) Histo-phytochemical evaluation and characterisation of the foliar structures of *Tagetes minuta* L. (Asteraceae) (Doctoral dissertation). Master's thesis. Engineering and Science University of KwaZulu-Natal Westville South Africa.
- Sarvaiya R.M., Rathod N.P. and Patel R.M. (2018) Bio-Efficacy of Bio-Pesticides against Aphid, *Aphis craccivora* Koch Infesting Fenugreek. *International Journal of Current Microbiology and Applied Sciences* 7(10): 2634–2640.

- Shahzadi I., Hassan A., Khan U.W. and Shah M.M. (2010) Evaluating biological activities of the seed extracts from *Tagetes minuta* L. found in Northern Pakistan. *Journal of Medicinal Plants Research* 4(20): 2108–2112.
- Tomova B.S., Waterhouse J.S. and Doberski J. (2005) The effect of fractionated *Tagetes* oil volatiles on aphid reproduction. *Entomologia Experimentalis et Applicata* 115(1): 153–159.
- Vasquez C., Balza D., Jimenez M.A., Colmenarez Y. and Rios Y. (2016) Use of plant extracts as an alternative control method against phytophagous mites in South America. *Current Topics in Phytochemistry* 13: 36–40.
- Veeranna H.K. and Adivappar N. (2019) Seasonal Incidence of Aphid, *Aphis craccivora* (Koch) Infecting Yard Long Bean, *Vigna unguiculata* sub spp. *Sesquipedalis*. *International Journal of Current Microbiology and Applied Sciences* 8(12): 2197–2203.

(Received July 26, 2023; revised ms accepted September 16, 2023; published December 31, 2023)

Exploration of natural enemy fauna of aphids and associated ant species from eastern dry zone of Karnataka, India

V.S. Chinnu¹, Vidya Mulimani^{*2}, S. Shanas³ and N. Sumithramma¹

¹Department of Entomology, College of Agriculture, GKVK, University of Agricultural Sciences, Bengaluru 560065, Karnataka, India.

²AINP on Acarology, Department of Entomology, College of Agriculture, GKVK, University of Agricultural Sciences, Bengaluru 560065, Karnataka, India.

³Integrated Farming System Research Station, Kerala Agricultural University, Karamana, Thiruvananthapuram 695002, Kerala, India.

Email: vidyamulimani34@gmail.com

ABSTRACT: Purposive surveys were conducted at regular intervals in different districts of eastern dry zone of Karnataka (Zone 5) during the year 2021-22, to document the natural enemy fauna and the ant species associated with aphids. Out of 34 aphid species recorded, 16 species of aphids were preyed on by 14 species of predators belonging to Coccinellidae, Syrphidae, Hemerobiidae and Chamaemyiidae and seven species of aphids were found parasitized by eight species of parasitoids belonging to Braconidae. Predators belonging to Coccinellidae and Syrphidae were recorded more abundantly with six species in each family, preying on 12 and 10 aphid species, respectively. Among the aphid parasitoids, *Aphidius* spp. was the more abundant taxa. Eleven species of ants belonging to Formicinae, Myrmicinae and Dolichoderinae were found associated with 17 aphid species. Ants belonging to the genus *Camponotus* were found to be more abundant and associated with 10 species of aphids. A comprehensive list of predators, parasitoids and ants associated with different aphid species was put together during this study. © 2023 Association for Advancement of Entomology

KEY WORDS: Purposive surveys, abundance, predators, parasitoids, aphidicolous ants

INTRODUCTION

Aphids (Hemiptera, Aphididae) are small soft bodied sap sucking insects. More than 450 species of aphids are found to be associated with different crop plants (Blackman and Eastop, 2000) of which about 100 species are of economic importance. Chemical insecticides have been used regularly for the management of aphid pests but not without risk of resurgence, destruction of natural enemies, development of resistance, phytotoxic effects,

environmental pollution and residual toxicity. Changing scenario of modern sustainable agriculture emphasizes the need for biological control for effective management of aphids. Aphids are good candidates for biological control (Joshi *et al.*, 2010) as they serve as a consistent and abundant food source for many natural enemies (Singh and Singh, 2016). Aphids have a mutualistic relationship with the ants which is by way of the aphids providing rich supply of food for ants in the form of honey dew and in return receiving

* Author for correspondence

protection from their natural enemies. The risk to aphids of death caused by predators can be decreased through recruiting more ants by producing more attractive honeydew (Stadler and Dixon, 2005). Presence of ants also accelerates growth of aphids and establishment of aphid colony (Saha *et al.*, 2018). Ants that attend aphids belong largely to the most evolved subfamilies Myrmicinae, Formicinae and Dolichoderinae (Tizado *et al.*, 1993). Noteworthy contributions on the natural enemies of aphids in India were made by Rao (1969), Raychaudhuri (1978), Ghorpade (1981), Ghosh and Raychaudhuri (1982), Sary and Ghosh (1983), Agarwala and Ghosh (1988), Singh *et al.* (1999), Joshi (2005), Dey and Akthar (2007), Akthar *et al.* (2011), Chaudhary and Singh (2012), Joshi and Sangma (2015), Bhat and Bhagat (2017), Khan *et al.* (2017), Bhat *et al.* (2020), Kale *et al.* (2020) and Maji *et al.* (2023). In India, aphid-ant association has been mainly dealt with Gadiyappanavar (1970), Roy and Behura (1980), Kurl and Misra (1980), Datta *et al.* (1982, 1983), Devi and Singh (1986), Verghese and Tandon (1987), Devi *et al.* (1987), Devi *et al.* (2001), Bisht *et al.* (2002), Joshi (2005), Kataria and Kumar (2013) and Rakshan and Ahmad (2015). The present investigation was carried out with an aim to record the natural enemies and ant species associated with aphids from eastern dry zone of Karnataka (Zone 5).

MATERIALS AND METHODS

In order to record the various natural enemies associated with different species of aphids, purposive surveys were conducted at regular intervals (15 days) in six districts of eastern dry zone of Karnataka *viz.*, Bengaluru Urban (13.0801° N; 77.5406° E), Bengaluru Rural (13.3535° N; 77.5406° E), Kolar (13.1320° N; 78.1783° E), Chikkaballapura (13.3354° N; 78.0824° E), Tumkur (13.2818° N; 77.1860° E) and Ramanagara (12.9576° N; 77.2261° E) during 2021-22. Field collection of aphid infested plant parts and associated aphid predators was carried out. The aphids collected were preserved in small plastic vials containing ethyl alcohol (70%), properly furnished with labels and the slides were prepared in accordance with the method suggested by Eastop

and van Emden (1972). Identification of aphids up to species level was carried out using the keys available from Aphids on the world's plants: an online identification and information guide and on confirmation of the identity with the specialist. Immature stages of predators associated with aphids were collected in small plastic containers and brought to the laboratory. These were reared to adult stages in rearing containers, providing respective host aphids as food (Joshi *et al.*, 1997). To record the parasitoid species associated with aphids, a part of the collection of aphid infested plant material and also colony with mummified aphids were brought to the laboratory (Joshi, 2005). Emerging adult parasitoids and predators were collected and processed for study. Identification of the predators and parasitoids were carried out with the help of specialists in the respective fields. Ants associated with different species of aphids were collected directly from the field and preserved in small vials containing alcohol (70%) for identification.

RESULTS AND DISCUSSION

During the course of investigation, out of 34 aphid species recorded, 16 species of aphids were preyed on by 14 species of predators belonging to four families and seven species of aphids were found parasitized by eight species of parasitoids belonging to Braconidae. The aphidophagous predators belonged to Coccinellidae, Syrphidae, Hemerobiidae and Chamaemyiidae. Six species each of aphidophagous coccinellid beetles and syrphids were recorded preying on 12 and 10 aphid species, respectively. Members of Hemerobiidae were recorded feeding on six aphid species and *Leucopis* sp. belonging to Chamaemyiidae was recorded preying on three aphid species (Table 1). Out of the eight aphid parasitoids, *Aphidius* spp. was recorded parasitizing five aphid species. Other parasitoids recorded include *Binodoxys* sp., *Lysiphlebus* sp. and *Trioxys* sp. which were recorded feeding on single aphid species each (Table 2). Out of 34 species of aphids reported, 17 species of aphids were found associated with 11 species of ants (Table 3). Three subfamilies of Formicidae were reported. Subfamily Formicinae was found more abundant with five species,

Table 1. Predatory species associated with different species of aphids in eastern dry zone of Karnataka during 2021-22

Predator	Aphid	Host plants
<i>Cheilomenes sexmaculata</i> (F.) Coccinellidae	<i>Aphis craccivora</i> Koch	<i>Cajanus cajan</i> (L.); <i>Cyamopsis tetragonoloba</i> (L.); <i>Dolichos lablab</i> L.; <i>Vigna unguiculata</i> (L.)
	<i>A. odinae</i> (van der Goot)	<i>Anacardium occidentale</i> L.
	<i>Brevicoryne brassicae</i> (Linnaeus)	<i>Brassica oleraceae</i> L. var. <i>capitata</i>
	<i>Hysteroneura setariae</i> (Thomas)	<i>Melinis repens</i> (Willd.) Zizka
	<i>Hyadaphis coriandri</i> Das	<i>Anethum graveolens</i> L.
	<i>Macrosiphum rosae</i> (Linnaeus)	<i>Rosa</i> sp.
	<i>Myzus persicae</i> (Sulzer)	<i>Br. juncea</i> L.
	<i>Rhopalosiphum maidis</i> (Fitch)	<i>Zea mays</i> L.
	<i>Schoutedenia emblica</i> (Patel and Kulkarni)	<i>Phyllanthus emblica</i> L.
<i>Coccinella transversalis</i> Fabricius Coccinellidae	<i>A. craccivora</i>	<i>Arachis hypogea</i> Willd
	<i>Br. brassicae</i>	<i>Br. oleraceae</i> L. var. <i>capitata</i>
	<i>M. persicae</i>	<i>Br. juncea</i>
	<i>R. maidis</i>	<i>Z. mays</i>
<i>Propylea dissecta</i> (Mulsant) Coccinellidae	<i>A. craccivora</i>	<i>Arachis hypogea</i> C. <i>cajan</i>
<i>Pseudaspidimerus</i> sp. Coccinellidae	<i>A. citricida</i> (Kirkaldy)	<i>Artocarpus heterophyllus</i> Lamk.
	<i>A. odinae</i>	<i>Garcinia indica</i> Choisy
<i>Scymnus nubilus</i> Mulsant Coccinellidae	<i>R. maidis</i>	<i>Z. mays</i>
<i>S. latemaculatus</i> Motschulsky Coccinellidae	<i>A. craccivora</i>	<i>Gliricidia maculate</i> (Jacq.)
	<i>A. gossypii</i> Glover	<i>Hibiscus rosa sinensis</i> L.
	<i>A. nerii</i> Boyer de Fonscolombe	<i>Calotropis gigantea</i> (L.)
<i>Asarkina belli</i> Ghorpade Syrphidae	<i>A. gossypii</i>	<i>H. rosa sinensis</i> .
<i>Betasyrphus</i> sp. Syrphidae	<i>A. craccivora</i>	<i>C. cajan</i>
	<i>A. nerii</i>	<i>C. gigantea</i>
	<i>Lipaphis pseudobrassicae</i> (Davis)	<i>Br. juncea</i>
	<i>M. persicae</i>	<i>Br. juncea</i>
<i>Dideopsis aegrota</i> (F.) Syrphidae	<i>A. odinae</i>	<i>Anacardium occidentale</i> L.
<i>Episyrphus viridaureus</i> (Wiedemann); Syrphidae	<i>Macrosiphum rosae</i> (Linnaeus)	<i>Rosa</i> sp.
<i>Ischiodon scutellaris</i> (Fabricius) Syrphidae	<i>A. craccivora</i>	<i>C. cajan</i>
	<i>A. gossypii</i>	<i>Cucumis sativa</i> L.; <i>Chromolaena odorata</i> (L.)
	<i>A. citricida</i>	<i>Citrus</i> sp.
	<i>Hyperomyzus carduellinus</i> (Kirkaldy)	<i>Sonchus</i> sp.

<i>Serratoparagus serratus</i> (Fabricius) Syrphidae	<i>A. craccivora</i>	<i>C. cajan</i> ; <i>Cyamopsis tetragonoloba</i> (L.); <i>G. maculata</i>
	<i>A. citricida</i>	<i>Citrus</i> sp.
	<i>R. maidis</i>	<i>Z. mays</i>
Unidentified Hemerobiidae	<i>A. craccivora</i>	<i>Vigna unguiculata</i> (L.)
	<i>A. odinae</i>	<i>Aralia</i> sp.
	<i>A. gossypii</i>	<i>Lantana camara</i> L.
	<i>Macrosiphum euphorbiae</i> (Thomas)	<i>Rosa</i> sp.
	<i>R. maidis</i>	<i>Zea mays</i> L.
	<i>Uroleucon compositae</i> (Theobald)	<i>Phyllocephalum scabridum</i> (DC.)
<i>Leucopis</i> sp. Chamaemyiidae	<i>A. gossypii</i>	<i>H. rosachinensis</i>
	<i>Hysteroneura setariae</i> (Thomas)	<i>Melinis repens</i> (Willd.) Zizka
	<i>R. maidis</i>	<i>Z. mays</i>

Table 2. Parasitoid species [Braconidae, Aphidiinae] associated with different species of aphids in the eastern dry zone of Karnataka during 2021-22

Parasitoid	Host aphid	Host plants
<i>Aphidius matricariae</i> Haliday	<i>Aulacorthum solani</i> (Kaltenbach)	<i>Brugmansia suaveolens</i> (Humb. & Bonpl. Ex Willd.)
<i>Aphidius</i> sp.	<i>Macrosiphum rosae</i> (Linnaeus)	<i>Rosa</i> sp.
<i>Aphidius</i> sp.	<i>Myzus persicae</i> (Sulzer)	<i>Brassica juncea</i> L.
<i>Aphidius</i> sp.	<i>Macrosiphoniella sanborni</i> (Gillette)	<i>Chrysanthemum indicum</i> L.
<i>Aphidius</i> sp.	<i>Lipaphis pseudobrassicae</i> (Davis)	<i>Brassica oleraceae</i> L. var <i>capitate</i>
<i>Binodoxys</i> sp.	<i>A. gossypii</i> Glover	<i>Duranta</i> sp.
<i>Lysiphlebus</i> sp.	<i>A. craccivora</i> Koch	<i>Arachis hypogea</i> Willd.
<i>Trioxys</i> sp.	<i>A. craccivora</i>	<i>Cyamopsis tetragonoloba</i> (L.)

followed by Myrmicinae and Dolichoderinae with three species each.

Among the coccinellid predators, more common was *Cheilomenes sexmaculata* (Fabricius), which was found preying on nine species of aphids followed by *Coccinella transversalis* Fabricius which preyed on four aphid species. Pervez (2004) provided a catalogue of predaceous coccinellids of India and its prey, which also gives an account of *Ch. sexmaculata* and *Co. transversalis* as predators. In the study conducted by Megha *et al.* (2015) on coccinellids in different crops at Dharwad region of Karnataka, *Ch. sexmaculata* was the

dominant species, consistent with the present findings. The catalogue of predaceous coccinellids of India gives the account of *Propylea dissecta* (Mulsant) feeding on *Aphis craccivora* Koch (Pervez, 2004). Agarwala and Ghosh (1988) provided prey records of aphidophagous coccinellids in India, which gives an early record of *Pseudaspidimerus* sp. preying on the *Aphis citricida* (Kirkaldy) and *A. odinae* (van der Goot). Megha *et al.* (2015) recorded *Scymnus nubilus* Mulsant feeding on *Rhopalosiphum maidis* (Fitch) from Dharwad region of Karnataka and the record of *S. latemaculatus* Motschulsky preying on *A. craccivora*, *A. gossypii* Glover and *A. nerii* Boyer

Table 3. List of aphidocolous ant species recorded from eastern dry zone of Karnataka during 2021-22

Ant species	Associated aphid species	Host plants
Subfamily - Formicinae		
<i>Anoplolepis gracilipes</i> (Smith)	<i>Aphis gossypii</i> Glover	<i>Gardenia resinifera</i> Roth
	<i>A. nerii</i>	<i>Calotropis gigantea</i> (L.)
	<i>A. odinae</i>	<i>Mussaenda erythrophylla</i> Lam.
	<i>Pseudoregma bambucicola</i> (Takahashp)	<i>Bambusa vulgaris</i> Schrad. ex J. C. Wendl.
	<i>Rhopalosiphum maidis</i>	<i>Zea mays</i> L.
<i>Camponotus</i> sp.	<i>A. craccivora</i>	<i>Cajanus cajan</i> (L.) <i>Cyamopsis tetragonoloba</i> (L.) <i>Gliricidia maculate</i> (Jacq.)
	<i>A. fabae</i>	<i>Solanum nigrum</i> L.
	<i>A. gossypii</i>	<i>Ruellia brittoniana</i> Leonard <i>Catharanthus rosea</i> (L.)
	<i>A. spiraecola</i> Patch	<i>Bidens Pilosa</i> L.
	<i>A. citricida</i> (Kirkaldy)	<i>Artocarpus heterophyllus</i> Lamk. <i>Citrus aurantifolia</i> Christm., <i>Citrus</i> sp.
	<i>A. odinae</i> (van der Goot)	<i>Hamelia patens</i> Jacq. <i>Mussaenda erythrophylla</i> Lam.
	<i>Cinara tujaefilina</i> (del Guercio)	<i>Thuja chinensis</i> Borders and Gausen
	<i>Hysteroneura setariae</i> (Thomas)	<i>Eleusine corocana</i> (L.) <i>Melinis repens</i> (Willd.) Zizka
	<i>Rhopalosiphum maidis</i> (Fitch)	<i>Zea mays</i> L.
	<i>Schoutdenia emblica</i> (Patel "and Kulkarni)	<i>Phyllanthus emblica</i> L.
<i>Crematogaster</i> sp.	<i>A. craccivora</i>	<i>Solanum torvum</i> Sw.
	<i>A. gossypii</i>	<i>Hibiscus rosa sinensis</i> L. <i>Lantana camara</i> L.
<i>Oecophylla smaragdina</i> Smith	<i>A. gossypii</i>	<i>Chromolaena odorata</i> (L.)
	<i>A. citricida</i> (Kirkaldy)	<i>Citrus</i> sp.
	<i>A. odinae</i> (van der Goot)	<i>Anacardium occidentale</i> L.
	<i>Schoutdenia emblica</i> (Patel and Kulkarni)	<i>Phyllanthus emblica</i> L.
<i>Paratrechina</i> sp.	<i>A. gossypii</i>	<i>Ocimum sanctum</i> L.
Subfamily - Myrmicinae		
<i>Lophomyrmex</i> sp.	<i>A. gossypii</i>	<i>Chromolaena odorata</i> (L.)
<i>Myrmecaria brunnea</i> (Saunders)	<i>A. craccivora</i>	<i>Cyamopsis tetragonoloba</i> (L.) <i>Dolichos lablab</i> L. <i>Moringa oleifera</i> Lam.
	<i>A. gossypii</i>	<i>Parthenium hysterophorus</i> L. <i>Tecoma stans</i> (L.)
	<i>A. odinae</i>	<i>Mussaenda erythrophylla</i> Lam. <i>Tagetes erecta</i> L.
	<i>H. setariae</i>	<i>Melinis repens</i> (Willd.) Zizka

<i>Solenopsis</i> sp.	<i>A. craccivora</i> <i>A. gossypii</i> <i>A. odinae</i> <i>H. setariae</i> <i>Uroleucon compositae</i> (Theobald)	<i>Arachis hypogea</i> <i>Vigna radiata</i> (L.) <i>Abelmoschus esculentus</i> Moench <i>Anacardium occidentale</i> <i>Eleusine indica</i> (L.) <i>Guizotia abyssinica</i> (L.f.) Cass.
Subfamily - Dolichoderinae		
<i>Tapinoma melanocephalum</i> (F)	<i>Aphis spiraeicola</i> Patch	<i>Chromolaena odorata</i> (L.)
<i>Technomyrmex albipes</i> (Smith)	<i>A. odinae</i>	<i>Pentas</i> sp.
<i>Technomyrmex</i> sp.	<i>A. craccivora</i>	<i>Arachis hypogea</i> Cordyline sp.
	<i>A. odinae</i>	<i>Aralia</i> sp.
	<i>Cerataphis brasiliensis</i> (Hempel)	<i>Areca catechu</i> L.
	<i>Pentalonia caladii</i> Boyer de Fonscolombe	<i>Alpinia zerumbet</i> (Pers.)
	<i>Pentalonia nigronervosa</i> Coquerel	<i>Musa</i> sp.
	<i>Melanaphis sacchari</i> (Zehntner)	<i>Saccharum officinarum</i> L.
	<i>Schoutdenia emblica</i> (Patel and Kulkarni)	<i>Phyllanthus emblica</i> L.

de Fonscolombe have also been made earlier (Chaudhary and Singh, 2012), similar to the findings of this study.

Among the syrphids recorded during the present study, *Ischiodon scutellaris* (Fabricius) and *Betasyrphus* sp. were found in abundance preying on four aphid species each. The syrphid, *Serratoparagus serratus* (Fabricius) was recorded feeding on three species of aphids. *Asarkina belli* Ghorpade, *Dideopsis aegrota* (Fabricius) and *Episyrphus viridaureus* (Wiedemann) were also recorded, each feeding on a single species. Similar reports of these aphid-syrphid associations were also given by Ghorpade (1981). Hemeroibiidae are major biocontrol agents that are used against aphids in several parts of the world. Members of Chamaemyiidae such as *Leucopis glyphinivora* Tanas. are potential biocontrol agents against aphids (Singh and Singh, 2016).

During the investigation, eight aphid parasitoid species belonging to Braconidae were recorded. Joshi (2005) reported *Aphidius* spp. and *Binodoxys* sp. parasitizing the aphids analogous to the records made during the present study. Similarly, the catalogue of aphid, parasitoids (Braconidae, Aphidiinae) from India also provides an account of

Aphidius sp., *Lysiphlebus* sp. and *Trioxys* sp. parasitizing the aphids (Akthar *et al.*, 2011).

In the present study the *Camponotus* ants were more abundant, associated with ten species of aphids. Seven species of aphids were attended by *Technomyrmex* sp., *Anoplolepis gracilipes* (Smith) and *Solenopsis* sp. were found associated with five species of aphids. *Oecophylla smaragdina* Smith and *Myrmecaria brunnea* (Saunders) attended four species of aphids. *Crematogaster* sp. was found associated with two species of aphids. *Paratrechina* sp., *Lophomyrmex* sp., *Tapinoma melanocephalum* (Fabricius) and *Technomyrmex albipes* (Smith) attended one species of aphids each. Rohini (2017) also recorded 26 aphid species from Chikkamagaluru district and reported 11 species of ants associated with 12 species of aphids, where the genus *Camponotus* was encountered more commonly tending six species of aphids, *Solenopsis* sp. attended three species of aphids and *Anoplolepis gracilipes*, *Crematogaster* sp., *Myrmecaria* sp., *Oecophylla smaragdina*, *Technomyrmex albipes* and *Tapinoma* sp. attended one aphid species each. Similarly, Joshi (2005) recorded 66 species of aphids of which 23 species of aphids were attended by eleven species

of ants. *Camponotus compressus* (F.) was found most abundant, associated with 15 aphid species. A total of 22 species of aphid natural enemies which is constituted by 14 species of predators and eight species of parasitoids were found associated with 16 and seven aphid species, respectively in the eastern dry zone of Karnataka.

ACKNOWLEDGEMENTS

Authors thank Dr. Sunil Joshi, Principal Scientist (Entomology) & Head of Division of Germplasm Collection and Characterisation, ICAR - NBAIR, Bengaluru; Dr. Yeshwanth H.M., Taxonomist, National Centre for Biological Sciences (NCBS), Bengaluru; Dr. Prabhu C. Ganiger, Entomologist, PC Unit (ICAR), AICRP on Small Millets, UAS, GKVK, Bengaluru; Dr. J. Poorani, Principal Scientist, ICAR - National Research centre for Banana, Tiruchirapalli, Tamil Nadu; Dr. K.J. David, Sr. Scientist (Agril. Entomology), Division of Germplasm Collection and Characterisation, ICAR - NBAIR, Bengaluru, Dr. A.N. Sringeswara, Curator, Mahatma Gandhi Botanical Garden, GKVK, University of Agricultural Sciences, Bengaluru and Dr. Shahid Bin Zeya, Professor, Department of Zoology, Aligarh Muslim University, Uttar Pradesh for the identification services provided.

REFERENCES

- Agarwala B.K. and Ghosh A.K. (1988) Prey records of aphidophagous Coccinellidae in India. A review and bibliography. *Tropical Pest Management* 34(1): 1–14.
- Akthar M.S., Dey D. and Usmani M.K. (2011) A catalogue of aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) from India. *Insecta Mundi* 151: 1–31.
- Bhat D.M. and Bhagat R.C. (2017) Host range and diversity of syrphid predators (Insecta: Diptera) of aphids on vegetable crops of Kashmir, with New Host Aphid/Plant Records. *Trends in Biosciences* 10(6): 1446–1448.
- Bhat D.M., Khan S.A., Ahanger F.A. and Sheikh M.A. (2020) Diversity of aphid pests (Homoptera: Aphididae) and their natural bio-control agents in vegetable crop ecosystems of Jammu & Kashmir, India. *International Journal of Current Microbiology and Applied Sciences* 9(5): 2529–2546.
- Bisht R.S., Rana D.S. and Katoch A.R. (2002) Records of aphidocolous ants (Hymenoptera: Formicidae) from Garhwal hills, Uttaranchal. *Journal of Aphidology* 16: 15–19.
- Blackman R.L. and Eastop V.F. (2000) *Aphids on the world's crops: an identification and information guide* (No. Ed. 2), John Wiley & Sons Ltd, Chichester, UK. 476pp.
- Chaudhary H.C. and Singh R. (2012) Records of the predators of aphids (Homoptera: Aphididae) in eastern Uttar Pradesh. *The Aphidological Society of India* 25: 13–30.
- Datta S.K., Raychaudhuri D. and Agarwala B.K. (1982) Study on aphid tending ants in India. I. New records of aphid and ant species in their association. *ENTOMON* 7(3): 327–328.
- Datta S.K., Raychaudhuri D. and Agarwala B.K. (1983) Study on aphid tending ants in India. II. New records of aphid and ant species in their association. *ENTOMON* 8: 23–25.
- Devi C.M. and Singh T.K. (1986) New aphidocolous ants from India. *Newsletter Aphidological Society India* 5: 5–7.
- Devi C.M., Paonam, M.S. and Singh T.K. (1987) Aphids (Homoptera: Aphididae) and their associated ants (Hymenoptera: Formicidae) of fruit trees in Manipur, north-east India. *Journal of Aphidology* 1: 102–104.
- Devi C.M., Singh T.K. and Nilamani L. (2001) New aphidocolous ants (Hymenoptera: Formicidae) from India. Subfamily: Myrmicinae. *Journal of Aphidology* 1: 102–104.
- Dey D. and Akthar M.S. (2007) Diversity of natural enemies of aphids belonging to Aphidiinae (Hymenoptera: Braconidae) in India. *Journal of Asia-Pacific Entomology* 10(4): 281–296.
- Eastop V. F. and van Emden H. F. (1972) The insect material. In: *Aphid Technology* (ed. H. F. van Emden). Academic Press, London and New York. pp1–31.
- Gadiyappanavar R.D. (1970) Study of Aphid Pests of Crops of Mysore State and the Biology of Ragi Root Aphid, *Tetraneura nigriabdominalis* (Homoptera: Aphididae). M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Bangalore, Karnataka, India.
- Ghorpade, K.D. (1981) Insect prey of Syrphidae (Diptera) from India and neighbouring countries: a review

- and bibliography. International Journal of Pest Management 27(1): 62–82.
- Ghosh A.K. and Raichaudhuri D.N. (1982) Ecology of natural enemy complex of Aphidoidea (Homoptera) in some areas of eastern India. In: Proceeding of the symposium on ecology of animal population. Zoological Survey of India 3: 55–69.
- Joshi S. (2005) Faunistic studies on aphididae (Hemiptera) of Karnataka and bioecology of the aphid parasitoid, *Diaeretiella rapae* (Mintosh) (Hymenoptera: Braconidae). Ph.D. Thesis (Unpub.), University of Agricultural Sciences, Bangalore, Karnataka, India.
- Joshi S. and Sangma R.H.C. (2015) Natural enemies associated with aphids and coccids from Sikkim, India. Journal of Biological Control 29 (1): 3–7.
- Joshi S., Rabindra R. J. and Rajendran T.P. (2010) Biological control of aphids. Journal of Biological Control 24 (3): 185–202.
- Joshi S., Venkatesan T. and Rao N.S. (1997) Host range and predatory fauna of *Aphis craccivora* Koch (Homoptera: Aphididae) in Bangalore, Karnataka. Journal of Biological Control 11(1/2): 59–63.
- Kale P., Bisen, A., Naikwadi B., Bhure K. and Undirwade D.B. (2020) Diversity study of aphids and associated predatory fauna occurred in major Kharif and Rabi crop ecosystems of Akola, Maharashtra, India. International Journal of Chemical Studies 8(4): 3868–3876.
- Kataria R. and Kumar D. (2013) On the Aphid–ant association and its relationship with various host plants in the Agroecosystems of Vadodara, Gujarat, India. Halteres 4: 25–32.
- Khan A.A., Shah M.A. and Riyaz S. (2017) Records of aphid and their natural enemies in agro-ecosystem with special reference to horticultural ecosystem of Kashmir. Journal of Entomology and Zoology Studies 5(4): 189–203.
- Kurl S. P. and Misra S.D. (1980) Interaction between the aphid and their attended ants. Journal of Environmental Research 1: 1–6.
- Maji A., Pal S., Gurung B. and Sahoo S.K. (2023) Diversity of aphids and their predatory coccinellids from West Bengal. Indian Journal of Entomology 85(2): 332–336.
- Megha R.R., Vastrad A.S., Kamanna B.C. and Kulkarni N.S. (2015) Species complex of coccinellids in different crops at Dharwad region. Journal of Experimental Zoology (India) 18(2): 931–935.
- Pervez A. (2004) Predaceous coccinellids in India: predator-prey catalogue (Coleoptera: Coccinellidae). Oriental Insects 38(1): 27–61.
- Rao V.P. (1969) Survey for natural enemies of aphids in India. Final Technical Report., U.S.P.L. 480 Project. C.I.B. C. Indian Station, Bangalore. pp1–93.
- Rakshan R. and Ahmad M.E. (2015) Study of mutualistic ants associated with *Aphis craccivora* (Hemiptera: Aphididae) on various host plants of family Fabaceae in Northeast Bihar (India). European Scientific Journal 11(18): 317–327.
- Raychaudhuri D.N. (1978) Some parasites and predators of aphids from northeast India and Bhutan. ENTOMON 3: 91–94.
- Rohini B.S. (2017) Species complex of aphids on horticultural crops in selected taluks of Chikkamagaluru district. M.Sc. (Horti.) Thesis, University of Agricultural & Horticultural Sciences, Shivamogga (COH, Mudigere), Karnataka, India.
- Roy D.K. and Behura B.K. (1980) Three more species of ants attending *Aphis gossypii* Glover. Newsletter Aphidological Society of India 1: 4–5.
- Saha S., Das T. and Raychaudhuri D. (2018) Myrmecophilous association between ants and aphids-an overview. World News of Natural Sciences 20: 62–67.
- Singh R. and Singh G. (2016) Aphids and their biocontrol. In: Ecofriendly pest management for food security. Academic Press. pp63–108
- Singh R., Upadhyay B.S., Singh D. and Chaudhary H.C. (1999) Aphids (Homoptera: Aphididae) and their parasitoids in north-eastern Uttar Pradesh. Journal of Aphidology 13: 49–62.
- Stadler B. and Dixon A.F. (2005) Ecology and evolution of aphid-ant interactions. Annual Review of Ecology, Evolution, and Systematics 36: 345–372.
- Stary P. and Ghosh A.K. (1983) Aphid parasitoids of India and adjacent countries (Hymenoptera: Aphidiidae). Technical Monograph, Zoological Survey of India 7: 1–96.
- Tizado E.J., Tinaut A. and Nafriaw J.N. (1993) Relationships between ants and aphids in the province of León (Spain) (Hym: Formicidae; Hom: Aphididae). Vie et Milieu/Life & Environment 43 (1): 63–68.
- Verghese A. and Tandon P.L. (1987) Interspecific associations among *Aphis gossypii*, *Menochilus sexmaculatus* and *Camponotus compressus* in a guava ecosystem. Phytoparasitica 15: 289–297

A new species of *Cigaritis* Donzel, 1847 (Lycaenidae, Aphnaeinae) from the southern Western Ghats of Peninsular India

S. Ramasamy Kamaya Naicker^{1,6}, Sujitha Prabhakaran Chandrika², Baiju Kochunarayanan³, Jebine Jose⁴, Manoj Kripakaran⁵, Rajkumar Chidambaram Palaniappan⁶, Vinayan Padmanabhan Nair⁷ and Kalesh Sadasivan^{8,9*}

¹Palace, Saptur 625705, Peraiyur Taluk, Madurai, Tamil Nadu, India.

²Department of Zoology, University of Kerala, Kariyavattom 695581, Thiruvananthapuram, Kerala, India.

³Sreerangam, Puravoorkonam, Karakulam 695564, Kerala, India.

⁴Kaduvathookil House, Elikulam 686577, Kottayam, Kerala, India.

⁵Shyladhri, TC 10/386(1), Njandoorkonam, Powdikonam 695587, Thiruvananthapuram, Kerala, India.

⁶Vanam Trust, 63/2, Lake Road, Sri Ram Nagar 625531, Theni, Tamil Nadu, India.

⁷XV/446 A1, Nethaji Housing Colony, Trichambaram, Taliparamba 670571, Kannur, Kerala, India.

⁸Greeshmam, BN439, Bapuji Nagar, Medical College 695011, Trivandrum, Kerala, India.

⁹Travancore Nature History Society (TNHS), MBRR 65, Jyothis, Mathrubumi Road, Vanchiyoor 695035, Thiruvananthapuram, Kerala, India.

Email: srknaicker@gmail.com; sujeespc@gmail.com; baijupaluvally@gmail.com; jebinkj2002in@gmail.com; manojvbm1@yahoo.com; drcpraj@gmail.com; vinayanpnair@gmail.com; kaleshs2002in@gmail.com

ABSTRACT: A new species *Cigaritis meghamalaiensis* **sp. nov.** (Lycaenidae, Aphnaeinae) is described from the Meghamalai Hills of the Periyar landscape of the southern Western Ghats. Images of adults and illustrations of male genitalia are presented. Information on myrmecophilous immature stages is provided and its ecology is discussed. The new species is very distinct from all the known *Cigaritis* species in WG, and is diagnosed based on the following combination of characters—upper side of both wings marked extensively blue in males; discal and post-discal bands on forewing underside conjoined and lying parallel from their origin at the costa; post-basal band in hindwing underside continuous and not broken into three smaller bands and this post-basal band ends at vein 1b, is not continued along it to reach discal band. The discal and post-discal bands on the underside of the forewing is conjoined and lying parallel from their origin at the costa which is a unique feature that distinguishes the new species from all other *Cigaritis* species occurring in Peninsular India and Sri Lanka. A key to all known species of *Cigaritis* from the Western Ghats is provided. © 2023 Association for Advancement of Entomology

KEYWORDS: Meghamalai Tiger Reserve, myrmecophily, new taxon, silverline, butterfly, crematogastor

INTRODUCTION

The Silverlines are strong-winged lycaenids in the subfamily Aphnaeinae Distant, 1884. Evans (1932)

treated Indian taxa of Aphnaeinae Distant, 1884 under *Aphnaeus* Hübner, [1819] in synonymy with *Spindasis* Wallengren, 1857 and considered

* Author for correspondence

Cigaritis Donzel 1847 as a senior synonym of *Apharitis* Riley 1925. Evans (1932) treated, species with prominent hindwing lobe and the tails at veins 1b and 2 nearly equal under *Spindasis* Wallengren, 1857 (= *Aphnaeus* Hübner, [1819]); and those species with ill-developed hindwing lobe and tail at v2 half as long as in vein 1 were treated under *Apharitis* Riley 1925 (= *Cigaritis* Donzel, 1847). Thus, the Western Ghats taxa were kept under the genera *Spindasis* and *Apharitis* by him. However, Heath (1997) synonymised *Apharitis* with *Spindasis*. The genus *Spindasis* was later synonymized with the genus *Cigaritis* Donzel, 1847, and *Aphnaeus* Hübner, [1819] was restricted to African taxa by Heath *et al.* (2002). Heath (1997), Heath *et al.* (2002), Heath and Pringle (2011), and later Boyle *et al.* (2015), using molecular data synonymized *Spindasis* with *Cigaritis*—the senior synonym. There are seven species of *Cigaritis* in the Western Ghats namely *C. vulcanus* (Fabricius, 1775), *C. schistacea* (Moore, [1881]), *C. ictis ictis* (Hewitson, 1865), *C. elima elima* (Moore, 1877), *C. lohita lazularia* (Moore, 1881), *C. lilacinus* (Moore, 1884), and *C. abnormis* (Moore, [1884]). Of these, except *C. lilacinus* all others have been reported from the southern Western Ghats (Unpublished data – Sadasivan *et al.*). A distinct *Cigaritis* species in the high elevations of Periyar Tiger Reserve, Idukki district, Kerala in 2018, and its myrmecophilous immature stages was observed. On further exploration in 2021, this species was found to be common in Meghamalai (Megamalai) of Tamil Nadu and adjoining Periyar Tiger Reserve of Kerala (Fig.1). This *Cigaritis* species was found to be new to science and is described here.

MATERIALS AND METHODS

The taxonomy of *Cigaritis* follows Evans (1932), Heath (1997), Heath *et al.* (2002), Heath and Pringle (2011) and Boyle *et al.* (2015). Identification of species follows Evans (1932), Wynter-Blyth (1957), and van der Poorten and van der Poorten (2018). Photographs of the specimens were taken with a Canon EOS 70D DSLR fitted with a 180mm macro lens and MPE 65 f 2.8 1–5x lens. The genitalia were studied by soaking overnight in KOH, then dissected under a stereo-zoom microscope

(HEADZ Model HD81) and preserved in glycerol. Illustrations were drawn by the senior author using the stereo-zoom microscope. The length of the forewing (FW) is measured as the longest straight-line distance from the wing base to the wing tip following Van Hook *et al.* (2012). Terminology for the wing pattern follows Evans (1932) and genitalia descriptions follow Corbet & Pendlebury (1992). The holotype and paratypes will be deposited in the insect collection of the Zoological Survey of India (ZSI), Western Ghat Regional Centre (WGRC), Kozhikode, Kerala, and Bombay Natural History Society (BNHS), Mumbai.

Abbreviations

BNHS	Bombay Natural History Society
PTR	Periyar Tiger Reserve, Kerala
SMTR	Srivilliputhur-Meghamalai Tiger Reserve
TNHS	Travancore Nature History Society
UpF	Upperside of forewing
UnF	Underside of forewing
UpH	Upperside of hindwing
UnH	Underside of hindwing
WG	Western Ghats
WLS	Wildlife Sanctuary
ZSI	Zoological Survey of India, Kozhikode

RESULTS AND DISCUSSION

Systematics

Family Lycaenidae Leach, 1815

Subfamily Aphnaeinae Distant, 1884

Genus *Cigaritis* Donzel, 1847

Cigaritis meghamalaiensis Sadasivan & Naicker
sp. nov. LSID urn:lsid:zoobank.org:act:A8D4F48B-46E6-4692-81D4-0D8F2EE4DA03

Holotype (Figs. 2A–B): TLRG 1001; Kardana Estate, Meghamalai, Theni District, Tamil Nadu

State, India; Col. SRK; 15.vi.2021, 1400m ASL, from a private estate; dry pinned specimen; will be deposited in the insect collections of ZSI, WGRC, Kozhikode Kerala, India.

Paratypes (1 male and 2 females) (Figs. 2C–G): ♂: TLRG 1002; bearing the same data as the holotype; dry pinned specimen; will be deposited in the insect collections of BNHS, Mumbai, Maharashtra, India. ♀ TLRG 1003 and ♀ TLRG 1004 bearing the same data as the holotype; dry pinned specimens, will be deposited in the insect collections of ZSI, WGRC, Kozhikode, Kerala, India and BNHS, Mumbai, Maharashtra, India, respectively.

Other materials (observed, not collected): 2 ♀♀, 26. xii. 2016, Eravangalar, PTR, Kerala, 1400m ASL (KS & JJ); 21 ♂♂, and 14 ♀♀, 16.iv.2021, Kardana Estate, Chinnamanur Range, Theni District, Tamil Nadu, 1420m ASL (SRK); 6 ♂♂ and 4 ♀♀, 21.iv.2023, KSR Estate, Meghamalai Range, Theni District, Tamil Nadu, 1320m ASL (SRK).

Description of the Holotype (♂ TLRG 1001)

Head. Antennae dark brownish-black, inferolateral striations grey, and tip distinctly marked in pale

orange-white; palpi dorsally blackish and covered with small thick whitish hairs ventrally; eyes grey with black speckling in life.

Thorax. dark grey bearing long pale greyish white hairs; legs very pale pinkish white, distally speckled in brown.

Forewing. Measures 16mm, costal margin and termen straight, apex minimally obtuse. Upperside with ground color velvety black, marked extensively with blue. Most of the spaces 1b and 2, three-fourths of space 3 from base, and just over half of space 4 are marked in metallic blue; a little over basal third of space 4 blue, rest of it black with a small blue spot at its middle; origin of space 5 blue; inferior half of cell blue. Entire costa marked broadly in black, this black border thickest at apex, then tapers on termen towards tornus; dorsum wholly marked in blue. Underside with ground color pale pinkish-brown, marked with a band at base of wing, and other long bands which are darker than ground color as follows—post-basal, discal, post-discal, sub-apical, submarginal bands pinkish-orange, centrally marked with silver scales, bordered with black; all long bands start at costa; post-basal band ends at origin of v2; discal band short, ends at origin of v3;




Fig. 1 Distribution of *Cigaritis meghamalaiensis* sp. nov. in Meghamalais of Periyar Landscape




Fig. 2 *Cigaritis meghamalaiensis* **sp. nov.** Images of the types and paratypes. A and B – TLRG 1001 Holotype male, A – dorsal and B ventral views; C and D –TLRG 1002, Paratype male, C – dorsal and D – ventral view; D and E – paratype female TLRG 1003, D – dorsal view and E – ventral view; F and G – paratype female TLRG 1004, F – dorsal view and G –ventral view. All images © Kalesh Sadasivan




Fig. 3 *Cigaritis meghamalaiensis* sp. nov. Field images of males, females and seasonal forms. A–male. Typical color; B–female, typical color; C–male upperside; D–female upperside; E–dry season male underside; F–dry season female underside; G–mating; H–oviposition. All images © Ramasamy Naicker




Fig. 4 *Cigaritis meghamalaiensis* sp. nov. Early stages, larval pens, and attending ants. A—freshly laid egg © Kalesh Sadasivan; B—larvae being attended by *Crematogaster* ants inside the larval pen © Kalesh Sadasivan; C—intermediate instar larvae and its attending ant © Kalesh Sadasivan; D—larval pen under the bark of a shola tree opened to reveal the walls (white arrows), and two larvae inside it marked 1 and 2 (blue arrows) © Jebin Jose

post-discal band consists of two conjoined bands, meeting at base of space 4; lower band turns basally and crosses into middle of space 1b; subapical band curved inwards ending at middle of space 3; submarginal band with its outer black border ill-defined, ends at v1b; a submarginal series of blackish-brown streaks on each space along termen; cilia brownish-black.

Hindwing. Upperside with ground color grey, whole of spaces 1c, 2, 3, 4, and cell marked in blue; origin of space 5 bears a small blue patch along vein 4; space 1b pale grey; costa broadly dark grey, termen narrowly marked in dark greyish black, dorsum pale greyish white with long pale grey hairs; whole of cell bears long bluish hairs; tornus pale orange white; tornal spot black; tails black, basal quarter pale orange-white, extreme tip white. Underside ground color as in forewing but slightly more orangish; short basal, longer-post-basal, discal, sub-apical and submarginal bands; longer bands originate at costa, run towards tornus; post-basal band continuous, not broken in three spots; discal band just crosses v1b, but does not reach tornus; post-basal and post-discal bands do not meet each other; sub-apical band crosses v3 at its middle; submarginal band just reaches 1b; all bands except the submarginal band bears central silver scales; silver scaling is sparse generally but well defined as a curvilinear streak running from end of submarginal band to middle of dorsal margin; tornal spots black, medial most spot almost twice as large as lateral.

Abdomen. Dorsally dark greyish-violet to violet-black, anterolaterally all segments bearing a cinnabar-red transverse streak that tapers towards ventrum, rest of segment yellowish-white; ventrally clothed in pale dirt white hairs.

Male genitalia (Fig. 6). Tegumen broad; uncus in lateral view broad, truncated, and flattened distally, appearing as a blunt tip tooth; dorsally tegumen and uncus appearing horseshoe shaped with a u-shaped gap separating the halves; uncus tip flattened and appearing spatulated in dorsal view; subuncal process shorter than uncus, thin, and directed towards opposite side and tips pointing posterolaterally; vinculum moderately thick, with a shallow concavity cephalad; saccus thicker than

vinculum; caudal plate of saccus absent; valva with a middorsal auricular process; dorsal process of valva long triangular, curved inferomedially and its tip directed posteroinferiorly, with respect to rest of valva; aedeagus as in fig. 6D, with its tip bearing a sectorized triangular plate, edges of which is toothed.

Description of female (Figs. 2D–G; 3B, D, F, H; 5D): A rounded greyish form of male without azure-blue upper sides. Wing span 35–38mm.

Head and Thorax. As in males.

Forewing. Measures 17mm, colour dark brownish-black with pale greyish-blue scaling in basal two-thirds of space 1 and base of space 2; termen rounded, apex rounded in comparison to males; cilia greyish-brown. All bands and marking as in male but sub-apical band more angulated towards termen in females in comparison to males.

Hindwing. A paler shade of forewing, greyish-brown, discal area clothed heavily in long bluish-grey hairs; tornal region yellowish-orange bearing a large medial black tornal spot and a smaller one laterally between tails; tails colored as in males. Cilia grey.

Abdomen. As in males with reddish lateral markings less prominent, speckled in black scales.

Variation: Not much variation was observed in the adults, except for the extent of silver scaling inside the bands and the lesser extent of reddish-orange hue on the underside in DSF individuals. Male genitalia is consistent. Forewing length was slightly variable to some extent, in males 15–17mm and females 16–18mm. On dry preservation, the blue shade on wings developed a greenish tinge.

Etymology: The new species is named after the Meghamalai region where it was discovered. Meghamalai means ‘cloud mountain’, reflecting the montane habitat of this very local species, which is restricted to the sub-tropical evergreen ‘sholas’ or cloud forests of the Periyar landscape. We suggest the common name ‘Cloud-forest Silverline’.

Ecology: Flight period observed was from

December to June. The butterfly, unlike its congeners, is very uninclined to fly and often falls easy prey to predators like *Monilesaurus acanthocephalus* Pal, Vijayakumar, Shanker, Jayarajan & Deepak, 2018 (Squamata, Agamidae). The butterfly is restricted to the sub-tropical evergreen forests and keeps to the forest edges where they perform mating and basking. The females were observed flying around trees occupied by *Crematogaster* ants. Another species that flies in the same elevation is *Cigaritis lohita*. The adults of *C. lohita* are not uncommon on the southern WG and their larvae have been observed on various plant families like Cannabaceae, Euphorbiaceae, Loranthaceae, Mimosaceae, and Myrsinaceae. However, the locally preferred species is *Maesa indica* (Roxb.) A. DC. (Myrsinaceae), and the larvae were seen attended by *Crematogaster rothneyi civa* Forel, 1902.

Immature stages and myrmecophily: Mating was noted in April (Fig. 3G), and oviposition was noted in mid-April and late December. Females lay eggs (Fig. 4A) on dry bark of trees such as *Neolitsea* (Benth. & Hook. f.) Merr. (Lauraceae) inside the shola and shrubs like *Clerodendrum infortunatum* L. on shola edges (Figs. 3H; 5D), invariably in the presence of the ant *Crematogaster wroughtonii* Forel, 1902 and their nests (Figs. 4 B–D). Oviposition was observed between 11.00 a.m. and 1.30 p.m. Eggs are laid on stems inside ant nests, dark crevices on tree trunks and fallen branches with moss and lichen. The females preferred trees and shrubs whose stem diameter was less than 20 cm, and eggs were laid at heights of 1 meter or less from the ground well away from any foliage. Immature stages were observed inside *Crematogaster* ant larval pens (Fig. 4D), under the bark of *Neolitsea cassia* (L.), Kosterm., a shola tree at Eravangalar in Periyar (1400m ASL). Each larval pen contained 3–4 larvae in various stages of development (Fig. 4D). We observed that the larvae scrape and eat the soft bark of the tree and are sheltered under the hard bark in small pens created by *Crematogaster* ants, possibly with the droppings of the caterpillars and vegetable matter (Fig. 4D). These larval pens were in the main trunk of the tree at a height of 1.5–2m from the ground,

very far away from any leaves of the tree. The ant nests were very far away from these larval pens, and thus the possibility of feeding by trophallaxis is suggested rather than the larvae being parasitic or predatory on these ant and their broods.

Of the seven species of *Cigaritis* known from the Western Ghats, six are reported on the southern Western Ghats namely *C. vulcanus*, *C. schistacea*, *C. ictis ictis*, *C. elima elima*, *C. lohita lazularia*, and *C. abnormis*. Of these, as per our field observation, *C. vulcanus* is a ubiquitous species seen in all elevations from sea coasts to 1200m, *C. schistacea* is an uncommon midland species (200–1200m), while *C. ictis* and *C. elima* are distributed below 800m, especially on the drier eastern slopes and *C. lohita* is a species which is found from the seacoast to about 1800m on the WG. *Crematogaster abnormis* is reported to occur on the lower eastern slopes of Coorg, Wayanad, Nilgiri, and Anamalai landscapes below 800m. Only *C. lohita* shares the elevational habitat of the new species. The ‘unidentified’ *Cigaritis* sp. mentioned in Sujitha *et al.* (2023) is described here as new to science and the myrmecophilous association with *Crematogaster wroughtonii* is confirmed. The new species is very distinct from all the known *Cigaritis* species in WG, and is diagnosed based on the following combination of characters—upper side of both wings marked extensively in blue; discal and post-discal bands on forewing underside conjoined and lying parallel from their origin at the costa; post-basal band in hindwing underside continuous and not broken into three smaller bands and this post-basal band ends at vein1b, is not continued along it to reach discal band. The presence of extensive blue coloration on Fw readily separates this new species from *C. elima elima*, *C. ictis ictis*, *C. schistacea*, and *C. vulcanus*, all of which have some form of orange stripes on the forewing. In addition, the unbroken post-basal band UnH distinguishes the new species from *C. elima elima*, *C. ictis ictis* as well as *C. abnormis*, and *C. lilacinus*. The hindwing underside post-basal band ends at vein1b, not continuing along it to reach the discal band separates the new taxon from *C. lohita*.

The discal and post-discal bands UnF conjoined and



Fig. 5 *Cigaritis meghamalaiensis* **sp. nov.**, habitat and host plants. A–Sub-tropical Evergreen forests of Meghamalais (1400m ASL) © Ramasamy Naicker; B–*Neolitsea cassia* (L.), Kosterm., a host tree © Kalesh Sadasivan; C–typical climate inside the misty cloud forests © Jebin Jose; D–female ovipositing inside *Crematogaster* nest on *Clerodendrum infortunatum* L. © Ramasamy Naicker

lying parallel from their origin at the costa is a unique feature that distinguishes *C. meghamalaiensis* **sp. nov.** from all known species of *Cigaritis* in Peninsular India and Sri Lanka. The discovery of a new species of *Cigaritis* from the southern Western Ghats reiterates the possibility of discovering new species which may have sought refuge in the montane sholas and cloud forests, which are under severe anthropogenic stress. The nature of myrmecophilous interaction needs to be studied in

detail, and the possibility of finding this new species must be kept in consideration wherever the attending host ant *C. wroughtonii* occurs in the Western Ghats.

**Key to *Cigaritis* Donzel, 1847 of Western Ghats
Modified from Wynter-Blyth (1957), based on
the males**

1. UnH the second band from the post-basal band continued along v1b to meet the discal band




Fig. 6 *Cigaritis meghamalaiensis* **sp. nov.**, Illustration of male genitalia (scale bar 1.5 mm). A—right lateral view of the genitalia with the aedeagus removed; B—ventral view of the valva; C—dorsal view of the uncus; D—right lateral view of the aedeagus

- | | |
|--|--|
| <p>near the tornus; below creamy yellow to cinnamon red, bands black to red; male above azure blue (Figs. 7A, B)<i>Cigaritis lohita lazularia</i></p> <p>– UnH post-basal band ends at vein 1b, not continue along it to reach the discal band.....2</p> <p>2. UnH the post-basal band continuous as judged by continuity of outer black margin of that band.....3</p> <p>– UnH post-basal band from base broken into 3 spots and not continuous.....5</p> | <p>3. UpF and UpH with orange stripes, in grey background; discal and post discal bands always separate at their origin at the costa.....4</p> <p>– UpF and UpH are marked extensively in blue on a black background; UnF discal and post discal bands conjoined and lying parallel from their origin at the costa. (Figs. 7G, H)<i>Cigaritis meghamalaiensis</i> sp. nov.</p> <p>4. A conspicuous small patch of blue scales near the orange tornal patch on UpH; orange patches on UpF apex restricted. (Figs. 8C, D).....<i>Cigaritis schistacea</i></p> |
|--|--|




Fig. 7 *Cigaritis* of Western Ghats. A–*Cigaritis lohita* male ventral view © Kalesh Sadasivan; B–*Cigaritis lohita* male dorsal view © Kalesh Sadasivan; C–*C. abnormis* male ventral view © Milind Bhakare; D–*C. abnormis* female dorsal view © Prateik More; E–*C. lilacinus* male ventral view © Kalesh Sadasivan; F–*C. lilacinus* male dorsal view © Haneesh KM; G–*C. meghamalaiensis* **sp. nov.** male ventral view © Ramasamy Naicker; H–*C. meghamalaiensis* **sp. nov.** male dorsal view © Ramasamy Naicker




Fig. 8 *Cigaritis* of Western Ghats. A—*Cigaritis vulcanus* male ventral view © Milind Bhakare, B—*C. vulcanus* male dorsal view © Milind Bhakare; C—*C. schistacea* male ventral view © Milind Bhakare; D—*C. schistacea* male dorsal view © Jebin Jose; E—*C. ictis* male ventral view © Milind Bhakare; F—*C. ictis* male dorsal view © Milind Bhakare; G—*C. elima* male ventral view © Haneesh KM; H—*C. elima* male dorsal view © Haneesh KM

- UpH without blue patch; orange patches on the UpF extensive. (Figs. 8A, B)*Cigaritis vulcanus*
- 5. UnH the lower two basal bands conspicuous, UpF apical orange patch present in males, above bright blue; three seasonal forms, pale yellow WSF with dark edged dark yellow bands, cinnamon or khaki autumn form with black-edged bands or khaki spring form with bands defined by silver lines and black margins absent.....6
- UnH the lower two basal bands inconspicuous and markings abnormal and narrow and only central bands are seen; male above only lightly blue, no orange apical patches; dull reddish brown below; forewing apex violet-brown and generally grey otherwise (Figs. 7C, D)*Cigaritis abnormis*
- 6. UpF with orange patches in brown background.....7
- UpF lilac blue in grey background (Figs. 7E, F).....*Cigaritis lilacinus*
- 7. UpF apical orange patch is developed and conspicuous; the basal blue patch does not reach vein 2; last band (postdiscal) on UnH meets or leans towards the submarginal band. (Figs. 8E, F)*Cigaritis ictis ictis*
- UpF apical orange patch restricted, basal blue patch reaches vein 2; the last band (postdiscal) on UnH never meets and leans away from the submarginal band. (Figs. 8G, H)*Cigaritis elima elima*

ACKNOWLEDGEMENTS

The authors are grateful to Srinivas R. Reddy IFS, PCCF & Chief Wildlife Warden, Tamil Nadu Forest Department; Padmavathe R., IFS, Field Director SMTR; Anand S., IFS, Deputy Director SMTR Theni; Sivaji T., Forest Range Officer, Chinnamanur; Suresh Kumar D., Forest Range officer Berigem Range; Ajay N., Forest Range Officer, Meghamalai; Pethanasamy V., Mutheesh J. and other field staff of SMTR. The authors are

thankful to Chidambara Kannan, Planter Kardana Estate-Metla Division, Meghamalai; Md. Jafer Palot and George Mathew for their encouragement. The authors thank Milind Bhakare, Haneesh K.M., Prateik More and Sunil Bhoite for providing images for comparison. The authors thank the members of TNHS, Thiruvananthapuram for providing the field support. The second author acknowledges the financial assistance received from the University of Kerala as a Post-Doctoral Fellowship (Ac EVII /2019-2020 dt.11.11.2021).

REFERENCES

- Boyle J., Kaliszewska Z., Espeland M., Suderman T., Fleming J., Heath A. and Pierce N. (2015) Phylogeny of the Aphnaeinae: Myrmecophilous African butterflies with carnivorous and herbivorous life histories. *Systematic Entomology* 40 (1): 169–182 doi:10.1111/syen.12098.
- Corbet A.S. and Pendlebury H.M. (1992) *The Butterflies of the Malay Peninsula*. 4th edition. Revised by J.N. Eliot. Malayan Nature Society, Kuala Lumpur. 595pp, 69 plates.
- Evans W.H. (1932) *Identifications of Indian Butterflies*. 2nd edition, Bombay Natural History Society, Bombay. x + 454pp, 32 pl.
- Heath A. (1997) A review of African genera of the tribe Aphnaeini (Lepidoptera: Lycaenidae). *Metamorphosis* 8 (Supplement 2): 1–60.
- Heath A., Newport M.A. and Hancock D. (2002) *The butterflies of Zambia*. African Butterfly Research Institute and The Lepidopterists' Society of Africa. i-xvii, 1–137.
- Heath A. and Pringle E.L. (2011) Taxonomic notes on the tribe Aphnaeini and genus *Thestor* Hubner (Lepidoptera: Lycaenidae) arising from D'Abrera (2009). *Metamorphosis* 22(1): 2–11.
- Sujitha P.C., Kalesh S., Nair V.P., Manoj K. and Prasad G. (2023) Myrmecophily in Lycaenidae Butterflies (Rhopalocera, Lepidoptera) of Kerala, Western Ghats, Southern India. *Oriental Insects* doi:10.1080/00305316.2023.2228299
- Van der Poorten G.M. and van der Poorten N.E. (2018) *Field guide to the Butterflies of Sri Lanka*. Lepodon Books, Toronto. 250pp.
- Van Hook T., Williams E., Brower L., Borkin S and Hein J. (2012) A standardized protocol for ruler-based measurement of wing length in monarch

- butterflies, *Danaus plexippus* L. (Nymphalidae, Danainae). International Journal of Tropical Insect Science 22: 42–52.
- Wynter-Blyth M.A. (1957) Butterflies of the Indian Region. Bombay Natural History Society, Bombay. 523pp.

(Received October 02, 2023; revised ms accepted November 11, 2023; published December 31, 2023)

Redescription of *Cyphochilus niveosquamosus* (Blanchard, 1851) (Coleoptera, Scarabaeidae, Melolonthinae) with notes on two species of the genus from India

K.V. Prakash^{1*} and H.M. Yeshwanth²

¹All India Network Project on Soil Arthropod Pests, Department of Entomology, University of Agricultural Sciences, G.K.V.K., Bengaluru 560065, Karnataka, India.

²Collection Facility, National Centre for Biological Sciences, TIFR, G.K.V.K., Bengaluru 560065, Karnataka, India.

Email: kvpento@gmail.com

ABSTRACT: *Cyphochilus niveosquamosus* (Blanchard, 1851) from India is redescribed. The habitus and parameres of *C. candidus* (Olivier, 1789) and *C. septentrionalis* Waterhouse, 1867 redescribed by Sabatinelli (2020) are also illustrated. The species are also redescribed based on the newly examined specimens in this study. A checklist with distribution of eight species of *Cyphochilus* from India is provided.

© 2023 Association for Advancement of Entomology

KEY WORDS: *Cyphochilus candidus*, *C. septentrionalis*, habitus, parameres, checklist

INTRODUCTION

The genus *Cyphochilus* Waterhouse, 1867 is characterized by the strongly asymmetric labrum among Scarabaeidae. The beetles possess an exceptional ability to scatter visible light from their scales to produce brilliant whiteness (Vukusic *et al.*, 2007). Sabatinelli, (2020) revised the “Section II” of the genus defined by Waterhouse (1867) as “the species having the mesosternum armed by a spur” and redescribed *C. candidus* (Olivier, 1789) and *C. septentrionalis* Waterhouse, 1867 for the first time and recognised two new species viz., *C. gandhii* Sabatinelli, 2020 and *C. satyarthii* Sabatinelli, 2020 from north India. So far, 64 species of the genus *Cyphochilus* have been described worldwide (Sabatinelli, 2020a, b, c; Sabatinelli and Pham, 2021; Zhao, 2021) and no information on the south Indian species except for the original

description of species is available. Despite, being a melolonthine beetle, hitherto none of the members of this genus have been reported as pest of any crop. In this paper, *C. niveosquamosus* (Blanchard, 1851) is redescribed from south India along with new locality record and notes of the two north Indian species, *C. candidus* (Olivier, 1789) and *C. septentrionalis* Waterhouse, 1867.

MATERIALS AND METHODS

The specimens were examined under a Nikon SMZ-1000 stereo binocular microscope for the study of the external morphology. For the study of male genitalia, the abdomen was separated from the rest of the body, cleaned in hot water and later macerated in hot KOH (10%), washed in distilled water and placed in glycerine for separating genitalia and further observation and imaging. The dissected

* Author for correspondence

genitalia and the rest of the specimen were suitably marked to ensure their association.

Images were made using Canon 7D camera with 100 mm and MPE65 mm lenses, series of images were taken at different depths and were subsequently stacked using Zerene Stacker software 1.04 to obtain high resolution images and grouped in plates using Adobe Photoshop CS2. All the specimens studied are deposited in the Department of Entomology, University of Agricultural Sciences, Bengaluru, abbreviated in the material examined section as UASB.

Abbreviations: Characters, measurements, mensural procedures, and ratios are those used by Sabatinelli (2020).

BL: Body length (from anterior margin of clypeus to apex of elytra, in dorsal view)

BW: Body width (across the elytral humeri, in dorsal view)

BWX: Greatest body width (across the elytral maximal width, in dorsal view)

CW/L: Clypeal ratio (width measured along clypeo-frontal suture divided by greatest length of clypeus)

F/O: Interocular ratio (minimum frons width across eyes divided by transverse compound eye diameter in dorsal view)

A2-7L/CL: Antennal ratio (derived from length of basal segments 2-7 divided by antennal club length)

PnW/L: Pronotal ratio (pronotal greatest width divided pronotal length along midline in dorsal view)

MstL: Mesosternal process length, in lateral view

RESULTS AND DISCUSSION

Genus *Cyphochilus* Waterhouse, 1867

Type species: *Melolontha candida* Olivier, 1789 by subsequent designation by Medvedev, 1951: 231.

Diagnosis: Principal diagnostic characters of *Cyphochilus* are: Labrum strongly asymmetric and

body surface covered with scales.

Cyphochilus niveosquamosus (Blanchard, 1851)
Figs. 1A-D, 2 A-D

Leucopholis niveosquamosa Blanchard, 1851: 158.

Waterhouse, 1867: 142 (*Cyphochilus*)

Material examined: INDIA: Tamil Nadu: 4♂, 1♀, Valparai: Near UPASI, Tea Research Station, 1062m, 10°16'N; 76°58'E, 02.vii.2014, leg. H.M. Yeshwanth (UASB).

Description: *Size* – BL: 17.74mm, BW: 7.81mm, BWX: 8.8mm at mid-point of elytra. *Colour* – Integument dark reddish brown including antennal club and legs; dorsal surface with uniformly pale white lanceolate scales each originating from a small puncture, scales denser along lateral margins of pronotum and elytra. *Head* – CW/L: 2.72; clypeus subrectangular, anterior margin straight, slightly reflexed; clypeo-frontal suture distinct; eyes prominent (F/O: 3); frons with an indentation behind the clypeo-frontal suture; antennal club longer than antennomeres 2-7 combined (A2-7/CL: 0.54). *Pronotum* – Transverse (PnW/L: 2), strongly convex, anterior margin concave, posterior margin arched at middle, anterior angles slightly obtuse, posterior angles angularly rounded, lateral margin angularly protruded almost at midlength, anterior half of the lateral margin devoid of scales. *Scutellum* – Semi-circular, covered with scales of varied size usually smaller than scales on elytra. *Elytra* – Elongate, parallel sided, irregularly punctate, stria not recognizable, suture raised, each elytron with two distinct and one feeble costae running parallel to each other. *Pygidium* – Broad, apex arched, dorsal surface with uniform small scales, apex and sides with a series of short hairs. *Thoracic sterna* – With long yellowish brown pubescence and sparse spiniform scales; prosternal process tubercular; mesosternal process peg-like, narrowed and rounded anteriorly, ventral surface slightly convex with a median longitudinal carina extending from a short distance before apex to midlength and dorsal apical surface flat (MstL: 1.18mm), ventrally glabrous. *Abdomen* – Ventrites with sparse scales,




Fig.1 *Cyphochilus niveosquamosus* (Blanchard, 1851), A-B: Male with white scales; A: Dorsal view, B: Ventral view, C-D: Male with yellow scales; C: Dorsal view, D: Ventral view

intersegmental suture prominent, border of 5th sternite with a row of short hairs, 1-4 ventrites glabrous. *Legs*—Protibia tridentate, apical tooth acute; subapical tooth triangular and positioned at same level as inner apex of protibia. *Aedeagus*—Basal piece 1.4× longer than paramere, medially longitudinally keeled, basi-dorsal angle smoothly curved, in lateral view gradually widened distally, lateral ventral margin in distal 1/3rd curved dorsally. Parameres fused till very close to apex, symmetrical, in lateral view gradually narrowed distally in basal 0.8 length then widened, distal apex bifid in lateral view, ventro-apical angles of lower branch projected and pointed, dorso-apical angle of upper branch rounded.

Variability: BL; 17.17–19.7mm (x=18.5mm, n=5); vestiture of scales pale yellow.

Remarks: Blanchard (1851) reported *Leucopholis niveosquamosa* from Mysore (erstwhile princely state). Here the species is recorded from Valparai (Tamil Nadu, south India). The specimens collected in this study are much shorter (around 18mm) than the specimens studied by Blanchard (1851) and Waterhouse (1866-67) which ranged from 24–25mm.

***Cyphochilus candidus* (Olivier, 1789)**

Figs. 3 A-E, 4 A-E

Melolontha candida Olivier, 1789: 15.




Fig. 2 *Cyphochilus niveosquamosus* (Blanchard, 1851)
Aedeagus - A: Dorsal view, B: Ventral view, C - D: Lateral views of paramere

Material Examined: INDIA: Meghalaya: 4♂, Ri-Bhoi, ICAR-Research Complex for North Eastern Hill Region; Umiam, 1031m, 25°41'N; 91°55'E, 3.vi.2013, leg. H. M. Yeshwanth; 1♂, Shillong, 1996; leg. S.K. Gangwar; 1♂, East Khasi hills: Mawkdok, 6.vi.2013, leg. H.M. Yeshwanth (UASB).

Description: *Size* – BL: 23.6mm, BW: 10.66mm, BWX: 12.08mm at mid-point of elytra. *Colour* – Integument dark reddish brown including antennal club and legs; dorsal surface with pale yellowish scale, each scale arising from a small puncture, denser along pronotal margins and sides of elytra. *Head* – CW/L: 2.67; anterior margin of clypeus straight, slightly reflexed; eyes prominent (F/O: 4.8); frons with a subtriangular depression behind clypeo-frontal suture; antennal club longer than antennomeres 2-7 combined (A2-7/CL: 0.65), one

of the setae on antennomere 7 approximately equal to 1/3 as long as antennal club. *Pronotum* – Transverse (PnW/L: 2), strongly convex, anterior margin concave, posterior margin arched at middle, anterior angles right angled, posterior angles angularly rounded and obtuse. *Scutellum* – Semi-circular densely covered with scales. *Elytra* – No visible stria or costa on elytra. *Pygidium* – Convex, apex deeply notched and apical margin reflexed, dorsal surface covered with scales. *Thoracic sterna* – With long yellowish brown pubescence and sparse scales; prosternal process tubercular; mesosternal process well developed (MstL: 2.4mm), acute, ventrally glabrous. *Abdomen* – Sternites with sparse scales, medioapical part of 8th sternite glabrous. *Legs* – Protibia tridentate, apical tooth acute; subapical tooth triangular and positioned at same level as inner apex of protibia.



Fig. 3 *Cyphochilus candidus* (Olivier, 1789), A: Dorsal view (Male with yellow scales); Aedeagus - B: Ventral view, C: Dorsal view, D: Left paramere with a spur, E: Right paramere

Aedeagus – Basal piece along with parameres ‘S’ shaped, about 1.3× as long as paramere, without median keel. Parameres asymmetrical, strongly curved dorsally, inner margin of parameres dorsally projected at midline, projection of left paramere more prominent than right one, left paramere with a spur, broad with rounded margin in basal half, strongly constricted and spine like in distal half.

Variability: BL: 22.32–24.71mm (x=23.5mm, n=3); vestiture of scales white (Fig. 4A-E). Shape of the spur on left paramere varied from subovate to subrectangular.

***Cyphochilus septentrionalis* Waterhouse, 1867, Figs. 5 A-E**

Cyphochilus septentrionalis Waterhouse, 1867: 141.

Cyphochilus pygidialis Nonfried, 1893: 332. Synonymised by Sabatinelli, 2020

Cyphochilus pygidialis v. *angeri* Nonfried, 1893: 333. Synonymised by Sabatinelli, 2020

Material studied: INDIA: Manipur: 1♂, Ukhrul; 1647 m, 25°06’N; 94°21’E, 23.viii.2014, leg. H.M. Yeshwanth (UASB).




Fig. 4 *Cyphochilus candidus* (Olivier, 1789), A: Dorsal view (Male with white scales); Aedeagus - B: Ventral view, C: Dorsal view, D: Right paramere, E: Left paramere with a spur




Fig. 5 *Cyphochilus septentrionalis* Waterhouse, 1867, A: Dorsal view (Male); Aedeagus - B: Ventral view, C: Dorsal view, D: Right paramere, E: Left paramere with a spine

Description: *Size* – BL: 23.44mm, BW: 10mm, BWX: 11.87mm at mid-point of elytra. *Colour* – Integument dark reddish brown including antennal club and legs; dorsal surface with uniformly yellow lanceolate scales, each scale originating from a small puncture, scales on clypeus, frons, pronotal and elytral disc comparatively larger; sides of pronotum, lateral and apical margin of elytra with smaller scales. *Head* – CW/L: 2.85; anterior margin of clypeus straight, slightly reflexed; clypeo-frontal suture feeble; eyes not prominent (F/O: 7.9); a subtriangular depression on frons just behind clypeo-frontal suture distinct; antennal club slightly shorter than antennomeres 2-7 combined (A2-7/CL: 1.12). *Pronotum* – Transverse (PnW/L: 2), strongly convex, anterior margin concave, posterior margin arched at middle, anterior angles almost right angled, posterior angles conically rounded and obtuse angled. *Scutellum* – Semi-circular, sparsely covered with scales. *Elytra* – Without visible stria or costa. *Pygidium* – Convex, apex rounded and apical margin reflexed, dorsal surface with scales. *Thoracic sterna* – With long yellowish brown pubescence and spiniform scales; prosternal process tubercular; mesosternal process well developed (MstL: 2.6 mm), acute, ventrally glabrous. *Abdomen* – Sternites with sparse scales, medioapical part of 8th sternite glabrous. *Legs* – Protibia tridentate, apical tooth acute; subapical tooth triangular and positioned at same level as inner apex of protibia. *Aedeagus* – Basal piece 1.33× longer than parameres and without median keel. Parameres asymmetrical, fused at base and dorsally curved, a posteriorly directed pointed spine on ventral inner margin at midlength of left paramere and a short tooth-like projection on ventral inner margin of right paramere subapically, right paramere concavely excavated along the inner margin subapically, left paramere slightly sinuate along inner margin subapically.

Remarks: This species resembles *C. candidus* (Olivier, 1789) externally but can be differentiated by the aedeagus. *C. candidus* has a subovate to subrectangular spur with distal attenuated process on the outer surface of left paramere, whereas *C. septentrionalis* has a posteriorly directed pointed spur on the ventral inner margin.

Updated checklist of species of *Cyphochilus* Waterhouse, 1867 known from India

C. candidus (Olivier, 1789) –India: Sikkim, West Bengal, Assam, Meghalaya, Manipur

C. gandhii Sabatinelli, 2020– India: West Bengal, Sikkim, Assam, Arunachal Pradesh

C. manipurensis Nonfried, 1893 – India: Manipur

C. niveosquamosus (Blanchard, 1851) – India: Karnataka, Tamil Nadu

C. oberthueri Brenske, 1903 – India: Tamil Nadu

C. satyarthii Sabatinelli, 2020– India: Sikkim, West Bengal

C. septentrionalis Waterhouse, 1867 – India: Uttar Pradesh, Sikkim, Assam; Meghalaya, West Bengal and Nagaland

C. waterhousei Brenske, 1903 – south India

ACKNOWLEDGEMENTS

The authors are grateful to Dr. Guido Sabatinelli, Natural History Museum, Geneva, Switzerland for confirming the identification of *C. niveosquamosus*. Special thanks to Prof. C.A. Viraktamath for helpful suggestions. Sincere thanks to Ming-Zhi Zhao, College of Plant Protection, South China Agricultural University, Guangzhou, China for improvising the manuscript.

REFERENCES

- Blanchard C.E. (1851) [New taxa]. In: Milne-Edwards H.: Catalogue de la Collection Entomologique du Muséum d'Histoire Naturelle de Paris. Classe des Insectes. Ordre des Coléoptères. Tome 1. Gide et Baudry, Paris (1850)1–128 (1851)129–240.
- Brenske E. (1903) Neue Arten der Melolonthiden (Coleopt.) Gattung *Cyphochilus*. Insecten Börse 20: 380–381.
- Olivier G.A. (1789) Entomologie, ou Histoire naturelle des insectes, avec leurs caractères générique et spécifiques, leur description, leurs synonymie et leur figure senluminées. Coléoptères. Tome premier. Baudoin, Paris. xx + 497pp., 65 pls.
- Medvedev S.I. (1951) Plastinchatousye (Scarabaeidae),

- posdem. Melolonthinae, ch. 1 (chrushchi). Fauna SSSR, zhestkokrylye. Tom. 10, vyp. 1. Moskva, Leningrad: Izd. Akad.Nauk. SSSR. 512pp.
- Nonfried A.F. (1893) Beiträge zur Käferfauna von Manipur (Vorderindien). Berliner entomologische Zeitschrift 38: 327–340.
- Sabatinelli G. (2020a) Taxonomic notes on the genus *Cyphochilus* Waterhouse 1867 (Coleoptera, Scarabaeoidea, Melolonthinae) with description of 10 new species. Revue suisse de Zoologie 127: 157–181.
- Sabatinelli G. (2020b) Taxonomic notes on the genus *Cyphochilus* Waterhouse 1867 (Coleoptera, Scarabaeoidea, Melolonthinae) (part 2) with description of nine new species and a new subspecies. Munis Entomology & Zoology 15: 301–318.
- Sabatinelli G. (2020c) Taxonomic notes on the genus *Cyphochilus* Waterhouse 1867 (Coleoptera, Scarabaeoidea, Melolonthinae) (part 3) with description of three new species and a new subspecies. Acta Societatis Zoologicae Bohemicae 84: 51–65.
- Sabatinelli G. and Phjm P. (2021) Taxonomic notes on the genus *Cyphochilus* (Coleoptera: Scarabaeoidea: Melolonthinae) (part 4) with description of eight new species from Indochina and China. Revue suisse de Zoologie 128 (1): 157–172.
- Waterhouse C.O. (1867) On some new lamellicorn beetles belonging to the family Melolonthidae. The Entomologist's Monthly Magazine 4: 141–146.
- Vukusic P., Hallam B. and Noyes J. (2007) Brilliant whiteness in ultrathin beetle scales, Science 315: 348.
- Zhao M.Z. (2021) On the genus *Cyphochilus* Waterhouse, 1867 from Hainan Island, China (Coleoptera: Scarabaeidae: Melolonthinae). Faunitaxys 9 (18): 1–8.

(Received September 09, 2023; revised ms accepted October 02, 2023; published December 31, 2023)

Evaluation of pongamia oil soap against red spider mites, *Tetranychus urticae* Koch and its effect on mite predators in brinjal ecosystem

P. Jeevitha^{1*}, K.M. Sreekumar² and Mukul Kumar Deka¹

¹Department of Entomology, Assam Agricultural University, Jorhat 78501 3, Assam, India.

²Department of Entomology, College of Agriculture, Padannakkad 671314, Kerala Agricultural University, Kerala, India.

Email: jeevithadpi9@gmail.com

ABSTRACT: The efficacy of pongamia oil soap was tested on red spider mites and on its predators (rove beetle, *Oligota* sp. and predatory gall midge, *Feltiella acarisuga* Vallot). Pongamia oil soap at 0.6, 1, 2 and 3 per cent along with neem oil 0.6 per cent, soap solution 0.5 per cent and thiamethoxam 25 WG were investigated and observations were recorded on, a day prior to and, 1, 3, 5, 7 and 14 days after spray application. The results revealed that pongamia oil soap @ 3, 2 and 1 per cent shown mite population reduction up to 86.50, 84.57 and 79.27 per cent respectively. Richness of mite predators didn't vary significantly at one day after spray and the maximum population was observed in soap solution (4.33/5 plants). Population of mite predators was relatively high at 14 days after spray with soap solution (35.67 per five plants) which was at par with pongamia oil soap @ 0.6, 1 and 2 per cent.

© 2023 Association for Advancement of Entomology

KEY WORDS: Rove beetle, predatory gall midge, thiamethoxam, neem oil, soap solution

INTRODUCTION

Brinjal is one of the major vegetables crops in India which positions second in brinjal (*Solanum melongena* L.) production with 12.80 MT of production (National Horticulture Board, 2018). There are 140 different types of insect and non-insect pests have a history of damaging the crop. The destructive pest of brinjal is fruit and shoot borer (BFSB), *Leucinodes orbonalis* Guenee (Lepidoptera, Crambidae) which causes enormous yield loss of as high as 70-92 per cent (Rosaiah, 2001). When infestation levels are high, sucking pests like red spider mites; *Tetranychus urticae*

Koch (Tetranychidae, Acari) dramatically lower crop output and inflict damage.

Millettia pinnata (L.) Pierre, often known as “pongam,” “Indian beech,” or “karanj,” is a multifunctional tree that is especially prized for the oil that is extracted from its seeds (27–40% oil). The primary furanoflavones responsible for karanj oil's insecticidal effects on pests, are karanjin, pongapin, kanjone, and diketone pongamol (Bringi, 1987). Different pongamia extracts shown antifeedant and oviposition-deterrent efficacy against various agricultural pests (Kumar *et al.*, 2006). A field experiment was conducted to

* Author for correspondence

evaluate the efficacy of four different concentrations of soap formulation of pungam (*Milletia/Pongamia*) oil against the red spider mite, *Tetranychus urticae* Koch and its effects against its natural enemies, (rove beetle, *Oligota* sp. and a predatory gall midge, *Feltiella acarisuga* Vallot population), in an eggplant ecosystem.

MATERIALS AND METHODS

Pungam oil soap made based on the technology used in “Ready to use Neem Oil Garlic Soap”, the biocide approved by Kerala Agricultural University (Varma, 2018). Pongamia oil was obtained from Tamil Nadu Agricultural University, Coimbatore to make pongamia oil soap required for the experiment. To prepare pongamia oil soap, 55g of caustic soda was mixed with 100ml of distilled water and left undisturbed for 4 hours. The solution blended scrupulously into the solution made up of 55g of soap stone power and one litre of pongamia oil. The final pongamia oil soap solution kept for solidification. The pongamia oil soap solution pH was determined (10.5) in Soil Science and Agricultural Chemistry lab, College of Agriculture, Padannakkad and saponification value was 194mg KOH/g of oil. Neem oil soap 0.6 per cent and standard check (Thiamethoxam) were also included as treatments to compare the results of pongamia oil soap.

Experiments were performed with the KAU released brinjal variety “Surya” under Randomized Block Design with 8 treatments (Table 1) and three replicates at Instructional Farm II, Karuvachery, College of Agriculture, Padannakkad. Eggplant seeds were sown in pro trays and 30 days old seedlings transplanted in a plot of size 3.4 X 2.8m² each with the spacing of 60 X 60cm. Pongamia oil soap 3, 2, 1 and 0.6 per cent spray solution were prepared by using pongamia oil soap 30, 20, 10 and 6g pongamia oil soap respectively and knapsack sprayer was used for spraying. Five representative plants among 12 plants were selected and tagged for recording observations. The observations were documented one day before, 1, 3, 5, 7 and 14 days after treatment application. The nymphs and adult red spider mites (RSM) and its natural enemies

(predatory gall midge and rove beetle) were counted from the 2 cm² leaf area of three leaves (one each from top, middle and lower) from represented plants. To calculate the per cent reduction in red spider mite population, Henderson and Tilton formula was used (Herderson and Tilton, 1955).

$$\text{Per cent reduction} = \left\{ 1 - \frac{n \text{ in C before treatment} \times n \text{ in T after treatment}}{n \text{ in C after treatment} \times n \text{ in T before treatment}} \right\} \times 100$$

The data on mites and its natural enemy's population count were analyzed after square root transformation by analysis of variance (ANOVA). Web Agri Stat Package (WASP) software was used to analyze the data on population count of mites and its enemies.

RESULTS AND DISCUSSION

Red spider mite population:

RSM population, ranging from 43.67 to 54.53 per 3 leaves recorded one day before the treatments, showed no significant difference among treatments. Pongamia oil soap 3per cent had significantly lower mites (5.53 RSM/3 leaves), comparable to 2 per cent pongamia oil soap on a day after spray application. The soap solution recorded the maximum mite population (91.40 RSM/3 leaves), comparable to the control plot (48.53 RSM/3 leaves). Pongamia oil soap at 1 and 0.6 per cent showed statistically equivalent results in reducing -mite populations (11.53 and 12.67 RSM/3 leaves respectively). Neem oil soap at 0.6 per cent (20.20 RSM/3 sheets) were found to be equivalent to thiamethoxam 25 WG (33.73 RSM/3 sheets) and both were comparable to pongamia soap at 0.6 and 1 percent.

Three days after spray, pongamia oil soap (3%) had the lowest mite population – (16.33 RSM/3 leaves), followed by pongamia oil soap at 2, 1, 0.6 per cent and neem oil soap (20.47, 21.07, 21.93 and 27.27 RSM/3 leaves respectively) and they were on par with each other. The 0.5 per cent soap solution recorded the highest mite count (55.07 RSM/3 leaves), comparable to the control (50.73

Table 1. Average population density of red spider mites, *Tetranychus urticae*

Treatments	Number of mites per 3 cm ² area of three leaves *					
	Second application					
	1 DBS	1 DAS	3 DAS	5 DAS	7 DAS	14 DAS
Thiamethoxam 25 WG 2g 10L ⁻¹	54.53 (6.93)	33.73 (5.74) ^{bc}	35.13 (5.69) ^{ab}	61.53 (7.83) ^a	74.47 (8.61) ^{ab}	85.33 (9.22) ^a
Pongamia oil soap 3%	43.67 (6.59)	5.53 (2.23) ^d	16.33 (4.02) ^b	24.40 (4.93) ^c	35.13 (5.92) ^f	47.67 (6.92) ^d
Pongamia oil soap 2%	50.47 (7.09)	7.87 (2.78) ^d	20.47 (4.50) ^b	34.87 (5.89) ^{bc}	36.60 (6.03) ^{ef}	55.73 (7.43) ^{cd}
Pongamia oil soap 1%	46.20 (6.70)	11.53 (3.34) ^{cd}	21.07 (4.55) ^b	35.67 (5.96) ^b	41.67 (6.43) ^{def}	62.47 (7.93) ^{bcd}
Pongamia oil soap 0.6%	51.00 (7.11)	12.67 (3.45) ^{cd}	21.93 (4.63) ^b	38.93 (6.22) ^b	50.60 (7.10) ^{cde}	64.93 (8.07) ^{abcd}
Neem oil soap 0.6%	47.13 (6.85)	20.20 (4.37) ^{bcd}	27.27 (5.16) ^b	37.93 (6.13) ^b	53.93 (7.33) ^{cd}	65.47 (8.09) ^{abc}
Soap solution 0.5%	67.00 (7.70)	91.40 (9.10) ^a	55.07 (7.30) ^a	76.67 (8.74) ^a	85.13 (9.18) ^a	73.20 (8.58) ^{ab}
Control	47.47 (6.89)	48.53 (6.96) ^{ab}	50.73 (7.09) ^a	64.73 (8.02) ^a	59.40 (7.70) ^{bc}	74.73 (8.67) ^{ab}
C.D. (P=0.05)	NS	2.60	1.86	1.03	1.17	1.16

* Mean of five observations; Means followed by similar alphabets don't differ significantly by DMRT at 5%; Figures in parentheses denotes square root transformed values

RSM/3 leaves). Plants treated with thiamethoxam 25 WG (35.13 RSM/3 leaves) were comparable to all other treatments tested. Observations made 5 days after treatment pongamia soap (3%) showed least 24.40 RSM/3 leaves, followed by pongamia soap (2 and 1%), neem oil soap and pongamia soap (0.6%) with 34.87, 35.67, 37.93 and 38.93 RSM/3 leaves respectively. The soap solution (0.5%) recorded the highest mite count of 76.67 RSM/3 leaves. This was comparable to controls (64.73 RSM/3 leaves) and thiamethoxam 25 WG (61.53 RSM/3 leaves).

Mite populations were lowest on plants treated with pongamia soap @ 3 per cent (35.13 RSM/3 leaves)

on day 7 after treatment, compared to pongamia soap @ 2 and 1 percent with 36.60 and 41.67 RSM respectively. It was 50.60 RSM/3 leaves in pongamia soap (0.6%) which was on par with neem oil soap 0.6 per cent (53.93 RSM/3 leaves), and pongamia soap 1 and 2 per cent. The maximum population was observed in 0.5 per cent soap solution (85.13 RSM/3 leaves), followed by thiamethoxam 25 WG (74.47 RSM/3 leaves), and the control (59.40 RSM/3 leaves).

Increased mite populations were observed for all treatments. Pongamia soap (3%) recorded 47.67 RSM/3 leaves after 14 days of spraying - followed by pongamia oil soap 2 per cent (55.73 RSM/3

leaves), 1 per cent (62.47 RSM/3 leaves) and 0.6 per cent (64.93 RSM/3 cards) pongamia soap. Neem oil soap 0.6 per cent (65.47 RSM/3 leaves) was comparable to soap solution 0.5 per cent (73.20 RSM/3 leaves), pongamia oil soap 0.6 and 1 per cent, control (74.73 RSM/3 leaves) and thiamethoxam treated plot showed maximum mite population (Table 1).

Percentage reduction in spider mite population:

Pongamia oil soap 3, 2, and 1 percent were statistically equivalent, reducing mite populations by 86.50, 84.57, and 79.27 per cent, respectively whereas soap solution 0.5 per cent (0.17%) and the insecticide thiamethoxam 25 WG (19.60%) were comparable with control (0.00%) in population decline. Pongamia oil soap 0.6 per cent with a population reduction of 76.40 per cent was equivalent to 1, 2, and 3 per cent pongamia oil soap at 1 day post-treatment. Three days after spray application, a maximum population reduction was recorded in pongamia soap 3 per cent which was on par with pongamia oil soap 2, 1 and 0.6 per cent with 61.17, 57.90, 55.10 and 53.93 per cent decrease in population of mites respectively. At 0.5 per cent soap solution, it was comparable to the control (0.00%) and thiamethoxam 25 WG (32.43%) with only a 16.00 per cent reduction. Neem oil soap 0.6 per cent reduced the population by 45.73 per cent among botanicals, comparable with pongamia soap at 3, 2, 1 and 0.6 per cent.

The pongamia oil soap 3 per cent was the highest among all treatments in population reduction (55.67%), while the soap solution increased mite populations with a negative sign (-23.3%). This was consistent with thiamethoxam 25 WG (5.33%) and comparable to the control (0.00%) at 5 days of application. Pongamia soap @ 2 and 1 per cent, and neem oil soap 0.6 per cent were statistically similar in population decline, recording 44.00, 41.40, and 38.00 3 per cent respectively. After 7 days of application, pongamia soap @ 3 and 2 per cent showed reduction in mite population (42.00 and 40.33%, respectively). Pongamia oil soap @ 1 and

0.6 per cent showed statistically similar results in reducing mite population (18.00 and 17.02% respectively). However, the soap solution, thiamethoxam, and untreated plots showed no reduction in population and recorded negative results (of -47.00, -35.00, and 0.00%, respectively).

On the 14th day of application, population reduction was statistically non-significant among pongamia soap 3, 2, 1, 0.6 per cent and neem oil soap 0.6 per cent treated plots (31.33, 19.00, 18.67, 17.67 and 16.33% respectively). However, thiamethoxam 25 WG and soap solution 0.5% failed to reduce mite numbers, indicated by negative signs of -32 and -4.3 per cent, both comparable to the control plot (Table 2).

Relative abundance of mite predators:

Data on a day before spray application revealed that the mite predator's numbers didn't differ significantly among different treatments and it was at a range of 1.33 to 4.00/5 plants. Mite predator's abundance did not change significantly 1 day after spray application it was at a range of 1.33 to 4.33/5 plants. Three days after spray, predator populations were statistically non-significant and evenly distributed (1.67 to 5.67). After 5 days of spray also, there was no significant difference in – predator (1.67 to 7.67/5 plants). Observations on 7 DAS revealed that the 0.5 per cent soap solution treatment recorded more predator population of 11.00/5 plants, and significantly different from the other treatments. Pongamia oilseed soap 0.6 per cent recorded 9.33/5 plants and was statistically equivalent to 0.5 per cent soap solution. Pongamia soap 2 per cent and 0.6 per cent neem soap treatments were statistically similar with comparable results to control - (4.00, 4.00 and 4.33/5 plants, respectively), but the standard test and 3 per cent pongamia soap showed the lowest population- (0.33 and 2.33/5 plants respectively).

Abundance of mite predators was relatively high in all treatments on 14 DAS. Soap solution (0.5%) recorded highest at 35.67/5 plants which was comparable to pongamia oil soap at 0.6, 1 and 2 per cent with mite predator populations of 20.00,

Table 2. Percentage reduction in red spider mite's population during field evaluation

Treatments	Percentage reduction in mites *				
	Second application				
	1 DAS	3 DAS	5 DAS	7 DAS	14 DAS
Thiamethoxam 25 WG 2g 10L ⁻¹	19.60 ^c	32.43 ^{bc}	5.33 ^c	-35 ^c	-32 ^b
Pongamia oil soap 3%	86.50 ^a	61.17 ^a	55.67 ^a	42.00 ^a	31.33 ^a
Pongamia oil soap 2%	84.57 ^a	57.90 ^a	44.00 ^{ab}	40.33 ^a	19.00 ^a
Pongamia oil soap 1%	79.27 ^a	55.10 ^a	41.40 ^{ab}	18.00 ^b	18.67 ^a
Pongamia oil soap 0.6%	76.40 ^{ab}	53.93 ^a	37.07 ^b	17.02 ^b	17.67 ^a
Neem oil soap 0.6%	57.13 ^b	45.73 ^{ab}	38.00 ^{ab}	6.33 ^{bc}	16.33 ^a
Soap solution 0.5%	0.17 ^c	16.00 ^{cd}	-23.3 ^c	-47.00 ^c	-4.3 ^b
Control	0.00 ^c	0.00 ^d	0.00 ^c	0.00 ^c	0.00 ^b
C.D. (P=0.05)	21.03	21.04	17.78	13.14	15.96

* Mean of observations of five plants; Means followed by similar letters are not significantly different by DMRT at 5%; DAS- Days After Spray; NS – Non-Significant

19.67 and 18.00/5 plants respectively. All remaining treatments were comparable to control plots (9.33/5 plants) containing neem oil soap @ 0.6 per cent, pongamia oil soap @ 3 per cent, and standard checks, each with a mite predator population of 12.33, 11.33, and 8.00/5 plants (Table 3).

Overall, pongamia soap solutions (@ 0.6, 1, 2, and 3%) were superior to control in reducing the RSM population. The greatest reduction was observed immediate day after spraying, and all treatments except soap solution remained effective up to 5 days after spraying, after which they declined. This observation is consistent with the acaricidal effect of pongamia oil against the horseshoe mite (*Tetranychus ludeni*). Among the various sprayed chemical treatments, 0.6ml l⁻¹ Abamectin 1.8 EC showed the greatest reduction in mite population (70.32%) on 14th day of spraying, followed by spiromesifen and fenpyroximate while pongamia oil at 2 ml/l decreased by 49.58% on the fourteenth day (Raina, 2016). Among botanicals, tulsi leaf extract (5 and 10%), neem oil (3 and 5%) and notchi leaf extract (5 and 10%) recorded superior results

followed by 63.53 and 62.98 per cent egg reduction were recorded by pongamia oil soap at 3 and 5 per cent respectively (Raghavendra *et al.*, 2017).

The results obtained in the current study show that the miticidal effect of pongamia soap 0.6 per cent is better than that of 0.6 per cent neem oil soap. This result is in consistent with the study reporting, among the plants evaluated, pongamia oil was clearly the superior treatment, showing high mortality with a low LC₅₀ value of 0.008 per cent whereas neem oil recorded higher LC₅₀ value of 0.230 per cent at 24 h after treatment (Islam *et al.*, 2017). In a another study same results was found with abamectin reducing the population of red spider mite (*Tetranychus urticae*) on okra by 74.64 per cent, recording the highest efficacy (Singh *et al.*, 2018).

The nirgundi and pongamia oils were given better remedies than neem oil as evidenced by the minimum LC values of 78.40 and 194.20ppm respectively, and neem oil was 1469.88ppm against tea mite *Oligonychus coffeae* (Handique *et al.*,

2018). A reduction of up to 32.43 per cent was recorded on the third day after spraying, and from the fifth day onwards the effect began to decline, indicating an increase in the number of mites on the seventh and fourteenth day. In a study, thiamethoxam, clothianidin, and imidacloprid on cotton, corn, and tomatoes each showed increases in spider mite populations, and noted a 30 per cent increase in spider mite populations on cotton plants treated with thiamethoxam at the end of the study while in the current study are 32 per cent more mite numbers at 14DAS (Szczeplaniec *et al.*, 2013).

Data on rove beetle, *Oligota* and predatory gall midge, *Feltiella acarisuga* show that pongamia oil soap treated plants recorded higher population compared to even control plot. Soap solution 0.5 per cent recorded highest of 35.67/5 plants, followed by pongamia oil soap 0.6, 1 and 2 per cent. All remaining treatments were comparable to control plots (9.33/5 plants) containing neem oil soap @ 0.6 per cent, pongamia oil soap @ 3 per cent, and standard checks, each with a mite predator population of 12.33, 11.33, and 8.00/5 plants on 14th day after spray. As no previous studies on the

Table 3. Relative abundance of mite predators during field evaluation

Treatments	Number of mite predators per 3 cm ² area of three leaves*					
	Second application					
	1 DBS	1 DAS	3 DAS	5 DAS	7 DAS	14 DAS
Thiamethoxam 25 WG 2g 10L ⁻¹	1.67 (1.46)	1.67 (1.24)	1.67 (1.35)	1.67 (1.39)	0.33 (1.68) ^d	8.00 (2.67) ^b
Pongamia oil soap 3%	1.67 (1.46)	2.00 (1.41)	1.67 (1.46)	2.67 (1.76)	2.33 (1.68) ^{cd}	11.33 (3.09) ^b
Pongamia oil soap 2%	2.33 (1.68)	2.33 (1.52)	2.00 (1.56)	2.67 (1.77)	4.00 (2.12) ^c	18.00 (4.15) ^{ab}
Pongamia oil soap 1%	2.00 (1.56)	2.67 (1.63)	3.00 (1.86)	4.67 (2.13)	8.33 (2.96) ^b	19.67 (4.32) ^{ab}
Pongamia oil soap 0.6%	2.00 (1.56)	3.00 (1.71)	3.33 (1.95)	6.67 (2.64)	9.33 (3.13) ^{ab}	20.00 (4.42) ^{ab}
Neem oil soap 0.6%	1.33 (1.27)	1.33 (1.14)	2.33 (1.68)	1.67 (1.25)	4.00 (2.09) ^c	12.33 (3.50) ^b
Soap solution 0.5%	4.00 (2.03)	4.33 (1.94)	5.67 (2.43)	7.67 (2.81)	11.00 (3.38) ^a	35.67 (5.92) ^a
Control	4.00 (2.09)	2.67 (1.61)	2.67 (1.74)	6.00 (2.41)	4.33 (2.18) ^c	9.33 (3.05) ^b
C.D. (P=0.05)	NS	NS	NS	NS	0.50	1.89

• Means followed by similar alphabets do not differ significantly @ 0.05 DMRT; Figures in parentheses denotes square root transformed values; DAS- Days After Spray; NS – Non-Significant

effects of pongamia treatment on predatory mites were available in the literature, the results on predatory mite populations are compared with neem products. Adverse effects of applying phytopesticides to the predatory mite *Amblyseius andersoni* were reported (Castagnoli *et al.*, 2002). Eggs were unaffected by Biopyrene Plus (Pyrethrin 8 EC) (92.31% hatching) and Neemazal 10 EC (98.73% hatching) after treatment. Only 14.01 per cent toxicity was recorded on females by neemazal treatment whereas others showed 100 per cent toxicity. Survival percentage of protonymphs by neemazal treatment was 51.33 per cent and death was Zero.

It's been observed that the overall toxic effects of thiamethoxam were more than 90 per cent to *Phytoseiurus persimilis* by all different routes of exposure. However, the local exposure could result in low mortality and residual exposure (Pozzebon *et al.*, 2011). The present study is in accordance with the findings in which the neonicotinoids are mildly toxic to *Neoseiulus fallacis* (Jamil *et al.*, 2016), and neonicotinoids are moderately to highly toxic to *Galendromus occidentalis* and *G. fallacis* (Bostanian *et al.*, 2009). It may be concluded pongamia oil soap showed fair control effect on red spider mites, in brinjal better than neonicotinoid thiamethoxam. At the same time, pongamia soap at different concentration showed no negative effects on mite predators hence it is absolutely safe to use against brinjal pests.

ACKNOWLEDGEMENT

The authors highly acknowledge Kerala Agricultural University for the access on the online research papers and other papers and valuable support for this article.

REFERENCES

- Bostanian N.J., Thistlewood H.A., Hardman J.M., Laurin M.C. and Racette G. (2009) Effect of seven new orchard pesticides on *Galendromus occidentalis* in laboratory studies. *Pest Management Science: formerly Pesticide Science* 65: 635–639.
- Bringi N.Y. (1987) *Non-Traditional Oil Seeds and Oils in India*. Oxford and IBH Publication Pvt. Ltd, New Delhi, India.
- Castagnoli M., Angeli G., Liguori M., Forti D. and Simoni S. (2002) Side effects of botanical insecticides on predatory mite *Amblyseius andersoni* (Chant). *Journal of Pesticide science* 75: 122–127.
- Handique G., Rahman A. and Roy S. (2018) Toxicity of three plant-based oils against *Oligonychus coffeae* (Acari: Tetranychidae). *Journal of Entomological Science* 53: 93–95.
- Henderson C.F. and Tilton E.W. (1955) Tests with acaricides against the brown wheat mite. *Journal of Economic Entomology* 48: 157–161.
- Islam T., Biswas M.J.H., Howlader M.T.H. and Ullah M.S. (2017) Laboratory evaluation of *Beauveria bassiana*, some plant oils and insect growth regulators against two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae). *Persian Journal of Acarology* 6: 203–211.
- Jamil R.Z.R., Vandervoort C., Gut L.J., Whalon M.E. and Wise J.C. (2016) Lethal time of insecticides on the predator mite *Neoseiulus fallacis* (Acari: Phytoseiidae) following topical exposure. *The Canadian Entomologist* 148: 353–360.
- Kumar V., Chandrashekar K. and Sidhu O.P. (2006) Efficacy of karanjin and different extracts of *Pongamia pinnata* against selected insect pests. *Journal of Entomological Research* 30: 103–108.
- National Horticulture Board (2018) Horticultural Statistics At A Glance 2018. <http://agricoop.nic.in/sites/default/files/Horticulture>.
- Pozzebon A., Duso C., Tirello P. and Ortiz P.B. (2011) Toxicity of thiamethoxam to *Tetranychus urticae* Koch and *Phytoseiulus persimilis* Athias Henriot (Acari Tetranychidae, Phytoseiidae) through different routes of exposure. *Pest Management Science* 67: 352–359.
- Raghavendra K.V., Chinniah C., Ramaraju K., Muthiah C. and Balakrishnan K. (2017) Acaricidal action of plant derivatives and natural oils against two spotted spider mite, *Tetranychus urticae* Koch in rose (*Rosa* sp.) ecosystem. *Bull Environ Pharmacol Life Sciences* 6: 9–15.
- Raina S. (2016) Seasonal incidence of major pests of cowpea, *Vigna unguiculata* (L.) Walp. with special reference to management of mite pest. M.Sc. (Ag) thesis, Sher-e-Kashmir University of Agricultural Sciences and Technology, Jammu. 86pp.
- Rosaiah B. (2001) Evaluation of different botanicals

- against the pest complex of brinjal. *Pestology* 25: 14–16.
- Singh A.K., Koul K., Shankar U., Singh S.K., Mondal A. and Singh M. (2018) Seasonal incidence and management of red spider mite, *Tetranychus urticae* Koch on okra, *Abelmoschus esculentus* (L.) Moench. *Journal of Entomological and Zoological Studies* 6: 650–656.
- Szczepaniec A., Raupp M.J., Parker R.D., Kerns D. and Eubanks M.D. (2013) Neonicotinoid insecticides alter induced defenses and increase susceptibility to spider mites in distantly related crop plants. *PloS one* 8: e62620.
- Varma Y.C.K (2018) Ready To Use Veppenna Veluthulli Soap. *Kalpadhenu*. Jan-March 2018: 45–46.

(Received May 29, 2023; revised ms accepted August 10, 2023; published December 31, 2023)

Eri silkworm pupae as fish meal replacement in common carp (*Cyprinus carpio* L.) feed in indoor aquaria

R.K. Gokulakrishnaa and Selvamuthukumaran Thirunavukkarasu

Department of Entomology, Faculty of Agriculture, Annamalai University, Chidambaram 608002, Tamil Nadu, India.

Email: entogokul230696@gmail.com; entoselva@gmail.com

ABSTRACT: Eri silkworm (*Samia ricini* D.) pupae based fish feed, for common carp (*Cyprinus carpio* L.) advanced fry, in indoor aquaria was evaluated with eight different diets. Eri silkworm pupae meal (ESWPM) based fish feeds influenced the growth of common carp significantly. Fortnightly growth measurements of the carp in eight treatments showed maximum in ESWPM alone and in ESWPM 100 per cent treatments. The proximate nutrient and amino acid composition varied in the fish feed diets. ESWPM based fish feed recorded significant impact on the carp with reference to its weight gain, specific growth rate, feed conversion ratio and relative growth rate. The technology got patent.

© 2023 Association for Advancement of Entomology

KEYWORDS: By-product utilization, *Samia ricini*, pupae, *Cyprinus carpio*, feed

Fish meal has long been a significant source of protein in the fish feed industry. Furthermore, wild fish catches have been progressively dropping due to aquaculture's heavy reliance on them as a feed source for farmed fish (FAO, 2014). Aquaculture feed costs make up between 40 and 70 percent of the price of the fish produced (Wilson, 2002; Rana *et al.*, 2009). Due to the shortage of fish meal, other protein sources with comparable nutritional characteristics have been thoroughly researched (Daniel, 2018). Studies on insect protein as a partial or complete substitute for the fish meal are attempted (Van Huis, 2013). Antifungal and antibacterial peptides in insect meal may also help the meal's shelf life.

Some conflicting opinions also arise about chitin-

containing insect-based fish feed, which reduces nutrient absorption and digestibility and interferes with the fish's normal physiological functions. However, the three enzymes required for chitin digestion: chitinase, chitobiose, and lysozyme (Lindsay *et al.*, 1984; Fines and Holt, 2010), are present in both carnivorous and omnivorous fish species. Further, Aquatic organisms' immune systems are modulated by chitin, its derivatives, and active substances found in insect exoskeletons, such as antimicrobial peptides (AMPs). Chitin intake in moderate proportion can enhance fish gut health, immunity, and resistance to infectious diseases (Hoffman *et al.*, 1997; Kim and Rajapakse, 2005; Lin *et al.*, 2012; Harikrishnan *et al.*, 2012). Chitin is responsible for some of the immunostimulatory

* Author for correspondence

effects of insect meal (Su *et al.*, 2017; Li *et al.*, 2017; Gasco *et al.*, 2018; Henry *et al.*, 2018).

According to Chakrabarthy *et al.* (1973), silkworm pupae are a better meal for Indian major carps than mustard oil cake and rice bran. Among silkworms, the two popular types are mulberry silk worm (MSW), feeding on mulberry, and eri silk worm (ESW), feeding on castor. Using the under-utilized by-products, MSW and ESW pupae, containing nearly 54 and 62 per cent protein, respectively, as an alternative protein source provides ample scope (Zegeye, 2020; Altomere *et al.*, 2020). Moreover, the enormous potential for using ESW pupae as a feed resource is made possible by their composition, which includes 26.21 per cent lipid, 8 per cent moisture content, 56.83 per cent Poly Unsaturated Fatty Acids (PUFA), high methionine and lysine content, low omega-6/omega-3 fatty acid, and sodium: potassium ratio (Mrinal Ray and Gangopadhyay, 2021).

One of the most popular and commercially significant freshwater fish is the common carp, *Cyprinus carpio*, which accounts for 11 per cent of global freshwater aquaculture production (FAO, 2014). Global production of common carp accounts 3.4 per cent of the world's fish production. During its transitions from the advanced fry to the fingerlings stage, insects become one of its main sources of nutrition (Aquaculture feed and fertilizer resource information system, FAO, 2014). The current study aimed to determine ESW pupae's suitability as a protein supplement replacing fish meal in common carp feed.

MATERIALS AND METHODS

The ESW culture was maintained in the Department of Entomology, Faculty of Agriculture, Annamalai University, Tamil Nadu, India. On the fifth day of spinning, cocoons were cut open, pupae taken out, dried in a hot air oven (Technico, TLPPL 131) at 60°C for 12 hours, and made into powder. Various ratios of feed ingredients, *viz.*, ESW pupae powder, fish meal, groundnut cake, rice bran, fish oil, tapioca flour, and vitamin and mineral premix (Table 1), were ground and sieved through a 250-micron

mesh. The ingredients were combined with enough water, mixed into a soft dough, cooked under pressure for 15 minutes, cooled, and then pushed through a manual noodle's maker (with a 1mm diameter sieve plate) to obtain noodle-like threads. They were then baked in a hot air oven at 60°C for six hours to lower the moisture content to 8 per cent, broken and sieved to obtain the required pellet size (0.5 - 1.5mm), depending on the age of the fish. The prepared isocaloric and isonitrogenous ESW pupae-based fish feeds were packed in airtight plastic containers separately and stored at room temperature.

Cyprinus carpio advanced fries with a mean weight range from 0.300 – 0.450g were purchased from a commercial fish farm (M/s Shakthi fish farm). They were acclimatized for about two weeks with a continuous oxygen supply using an aquarium air pump (Sebo, AP 500) and commercial fish feed (Vrudhi Plus, Godrej Agrovet Limited, Mumbai). Then they were randomly stocked into an experimental glass tank (60cm x 45cm x 45cm). Feed was given daily once with a ratio of five per cent fish body weight. The faecal matter was siphoned out and a 75 per cent water change was done at weekly intervals. The experimental design was planned as a Completely Randomised Design (CRD) consisting of eight treatments, three replications, and ten fish per replication. Growth indices like Weight Gain (g), Specific Growth Rate (SGR), Feed Conversion Ratio (FCR), Relative Growth Rate (RGR), Survival Rate (%) and Fish In: Fish out (FIFO) (%) were calculated for ten weeks. Experimental feeds' proximate nutrient composition and essential amino acid composition analysis using three samples from each treatment were carried out following Sadasivam and Manickam (2005) and Bandlamori *et al.* (2012) protocol, respectively. The data were subjected to one-way ANOVA and analysed using SPSS Statistical package version 16.

Formulas for growth indices:

Weight Gain (g) = Final wet weight – Initial wet weight

Table 1. Composition of experimental diets (per 100 g) in the treatments

Feed Composition	ESWPM 20 %	ESWPM 40 %	ESWPM 60 %	ESWPM 80 %	ESWPM 100 %	ESWPM alone	Control	Positive* Control
Fish meal (g)	24	18	12	6	0	-	30	-
Eri Silkworm pupae powder (g)	6	12	18	24	30	62	0	-
Groundnut cake powder (g)	20	20	20	20	20	-	20	-
Rice husk (g)	37	37	37	37	37	-	37	-
Fish oil (ml)	2	2	2	2	2	-	2	-
Tapioca flour (g)	10	10	10	10	10	-	10	-
Vitamin and Mineral premix (g)	1	1	1	1	1	-	1	-
Bentonite	-	-	-	-	-	3	-	-
Cellulose	-	-	-	-	-	35	-	-

ESWPM - Erisilk worm pupae meal; *Commercial fish feed used as such

$$\text{Specific Growth Rate (SGR)} = \frac{(\ln(W_t) - \ln(W_0))}{t(d)} \times 100$$

W_0 [g]= the weight in grams at the beginning of the period

W_t [g]= the weight in grams at the end of the period

t [d]= period, expressed in number of days

\ln = natural logarithm

$$\text{Feed Conversion Ratio (FCR)} = \frac{\text{Feed Fed}}{\text{Wet weight gain}}$$

$$\text{Relative Growth Rate (RGR) (\%)} = \frac{\text{Final Weight} - \text{Initial Weight}}{\text{Initial Weight}}$$

$$\text{Survival (\%)} = \frac{\text{Number of live fishes at end of the experiment}}{\text{Total numbers of fishes Stocked}} \times 100$$

$$\text{Fish In: Fish Out (FIFO) (\%)} = \text{FCR} \times (\% \text{ Fish meal} + \% \text{ Fish oil in feed})$$

RESULTS AND DISCUSSION

Impact of eri silkworm pupae-based fish feed on common carp's growth performance

It was found that the ESW pupae-based fish feed significantly influenced common carp's growth indices. This performance of ESW pupae meal (ESWPM) at 20 and 40 per cent, recorded poor fortnight weight gains (Table 3). Such poor performance resulted due to the poor protein content of the feed fed (as both treatments contain more proportion of fish meal over ESWPM than other treatments). Nandeesh *et al.* (1990) reported such a reduced growth in common carp when fed with a diet containing 10 and 20 per cent non-defatted pupae meal. In contradiction to poor performance of ESWPM treatment, Jayaram and Shetty (1980a) reported better growth in 30 per cent defatted pupae meal fed catla and common carp.

During the first fortnight measurement, the maximum growth was recorded in ESWPM 100 per cent (5.1g) and ESWPM alone (5g) both treatments recorded on par with each other. Positive

Table 2. Proximate nutrient and essential amino acid composition of experimental feeds in the Treatments

Composition	ESWPM (20 %)	ESWPM (40 %)	ESWPM (60 %)	ESWPM (80 %)	ESWPM (100 %)	ESWPM alone	Control	Positive Control (Commercial fish feed)
Crude protein (%)	38.1	38.3	38.0	38.4	38.5	38.5	38.4	38
Crude lipid (%)	8.3	8.2	8.1	8.0	8.6	9.4	8.2	8.5
Crude fibre (%)	1.1	1.3	1.1	1.2	1.1	1.2	1.2	0.8
Metabolizable energy (MJKg ⁻¹)	17.5	17.2	17.4	17.3	17.6	18	17.3	17.0
Iron (mg 100 g ⁻¹)	13.1	13.0	13.3	13.5	13.8	13.5	13.7	12.9
Zinc (mg 100 g ⁻¹)	20.5	21.0	21.8	20.9	21.3	21.9	21.2	20.6
Calcium (mg 100 g ⁻¹)	1.4	1.5	1.5	1.9	2.0	2.1	1.4	1.3
Phosphorus (mg 100g ⁻¹)	1.1	1.2	1.3	1.2	1.4	1.5	1.4	1.5
Methionine (g 100 g ⁻¹)	1.1	1.1	1.4	1.3	1.5	1.6	1.5	1.4
Cystine (g 100 g ⁻¹)	1.3	1.3	1.2	1.4	1.5	1.6	1.9	1.6
Lysine (g 100 g ⁻¹)	2.7	2.5	3.3	3.0	3.3	3.7	4.0	3.2

control and ESWPM 80 per cent recorded next best weight gains. However, the trend was little bit changed in the second fortnight, ESWPM alone recorded higher value (12.6 g) followed by ESWPM 100 per cent (12.1 g). In third fortnight measurement, ESWPM 100 per cent and ESWPM alone both treatments were on par with each other. It was followed by positive control and ESWPM 80 per cent recorded on par with each other. Control treatment and ESWPM 60 per cent also were on par with each other. In fourth fortnight interval, ESWPM 80 per cent recorded weight a head than positive control. In last fortnight measurement (fifth fortnight interval), trend recorded in third fortnight interval evidenced. Irrespective of all the fortnight interval measurements ESWPM 40 and 20 per cent records the poor growth performance (Table 3). Weight gain (%) in experimental feed-fed fish over positive control represent the significant performance of ESWPM alone (23.57 % increase over positive control) and ESWPM 100 per cent (20.29 % increase over positive control) among other treatments.

Rangacharyulu *et al.* (2003) reported a 13 per cent higher weight gain when fish meal was entirely replaced with silkworm pupae silage in the Indian major carp's diet. The reason behind the enhanced performance is due to the presence of hydrolysed protein in ensiled silkworm pupae as compared to the complex proteins in fish meal (Manikandavelu *et al.*, 1992; Anon, 1999). Wan *et al.* (2017) reported that silkworm pupae meal is an attractive and sustainable functional feed component in carp diet with enhanced growth performance. Better protein and fat digestibility of carp diet containing silkworm pupae meal than the fish meal was reported by Nandeeshu *et al.* (1990) were in line with the present study findings. However, Nandeeshu *et al.* (1989 a, b) and Ji *et al.* (2015) reported least growth performance, impaired antioxidant enzyme status, decreased digestive function, and unfavourable changes in hepatic and intestinal morphology in fish fed with defatted silkworm pupae entirely or in combination with fish meal at different proportions. Further, Nandeeshu *et al.* (2000) found 50 per cent silkworm pupae

Table 3. Weekly growth performance (weight gain) of fish fed on experimental feeds (g) at fortnightly intervals

Treatments	Initial	1 st	2 nd	3 rd	4 th	5 th
ESWPM 20 %	0.358 (0.598) ^b	2.4 (1.549) ^d	4.8 (2.191) ^f	9 (3.000) ^e	14.4 (3.860) ^e	19.8 (4.505) ^e
ESWPM 40 %	0.333 (0.576) ^b	2.4 (1.549) ^d	5.7 (2.387) ^e	10.2 (3.193) ^d	15.2 (3.964) ^e	21.3 (4.669) ^d
ESWPM 60 %	0.337 (0.580) ^b	3.5 (1.871) ^c	8.8 (2.966) ^d	14.3 (3.781) ^c	18.7 (4.382) ^d	28.2 (5.357) ^c
ESWPM 80 %	0.333 (0.576) ^b	3.9 (1.975) ^b	9.8 (3.130) ^c	16.2 (4.025) ^b	22.5 (4.796) ^b	31.1 (5.621) ^b
ESWPM 100 %	0.365 (0.604) ^a	5.1 (2.258) ^a	12.1 (3.480) ^b	19.9 (4.462) ^a	27.2 (5.263) ^a	36.2 (6.060) ^a
ESWPM alone	0.424 (0.652) ^a	5 (2.236) ^a	12.6 (3.559) ^a	20.5 (4.528) ^a	28 (5.338) ^a	37.1 (6.132) ^a
Control	0.404 (0.635) ^a	3.5 (1.871) ^c	8.6 (2.932) ^d	14.2 (3.768) ^c	18.6 (4.370) ^d	27.9 (5.329) ^c
Positive Control (Commercial feed)	0.414 (0.643) ^a	3.8 (1.949) ^b	9.5 (3.082) ^c	15.6 (3.949) ^b	21.3 (4.669) ^c	30.2 (5.543) ^b
SEd	0.008	0.023	0.036	0.047	0.054	0.065
C.D(0.05)	0.016	0.046	0.072	0.094	0.109	0.130

Mean of three replications; Values within parentheses are square root transformed

inclusion diet were optimal to maintain growth performance and meat quality in common carp, contradicting the present finding of optimal growth performance in inclusion diets containing ESWPM above 60 per cent. As the fish meal incorporated might be contaminated, the resulting protein quality might also be inferior leading to conflicting results. Moreover, eri silkworm pupae's nutritive profile is completely different from mulberry silkworm pupae and fishmeal.

Poor weight gain recorded in fish fed with low proportion of ESW pupae powder might be due to amino acid profile imbalance. This was corroborated by the Kaushik and Seiliez (2010) and Wan *et al.* (2017). This was evidenced by the differences in amino acid profile of different feeds. The proximate nutrient and amino acid composition different

treatment fish feeds indicated variation (Table 2).

Similarly, Hora and Pillay (1962) found higher silkworm pupae incorporation led to offensive odour. Such odour related issues were observed in the present study only in the treatment ESWPM alone. Remaining ESW pupae-based fish feeds was in tune with the findings of Jayaram and Shetty (1980b) who reported no adverse flavour.

There was significant impact of eri silkworm pupae-based fish feed on common carp's growth indices (Table 4). Among the experimental feeds, fish fed with ESWPM alone recorded the highest weight gain of 36.67g followed by fish fed with ESWPM 100 per cent (35.86 g) and positive control (29.81g). Treatment ESWPM 60, 40 and 20 percent recorded least weight gains. Wan *et al.* (2017) studied the

Table 4. Growth indices of different experimental feeds fed common carp

Treatments	Weight gain	Specific Growth Rate (SGR)	Feed Conversion Ratio (FCR)	Relative Growth Rate (RGR)	Survival Rate (%)	Fish In: Fish Out (FI:FO) (%)
ESWPM 20 %	19.44	2.79	2.5	54.3	100	20
ESWPM 40 %	20.96	2.79	2.3	62.94	100	32.2
ESWPM 60 %	27.86	3.21	1.7	82.67	100	34
ESWPM 80 %	30.76	3.34	1.6	86.48	100	41.6
ESWPM 100 %	35.86	3.68	1.3	92.37	100	-
ESWPM alone	36.67	3.93	1.3	98.24	100	-
Control	27.49	3.46	1.8	68.04	100	57.6
Positive Control (Commercial feed)	29.81	3.61	1.6	72	100	-

effect of marine (Ragworm) and terrestrial invertebrate meals (Silkworm Pupae) in the diet of *C. carpio* and reported an increased weight gain of 12 per cent in diets containing fish meal + silkworm pupae when compared with the reference control diet.

Maximum percent increase in fish weight per day indicated by the parameter, Specific Growth Rate (SGR) was recorded in ESWPM alone. On an average, it recorded 3.93 per cent daily weight gain. The ESWPM 100 per cent and positive control recorded 3.68 and 3.61 per cent daily weight gains, respectively (Table 4).

Least Feed Conversion Ratio (FCR) of 1.3 was obtained from ESWPM alone and ESWPM 100 per cent followed by ESWPM 80 per cent and positive control records 1.6. Poor FCR noticed in ESWPM 40 and 20 per cent, hence both this feed composition is not recommended for commercial pisciculture (Table 4).

Relative Growth Rate (RGR) was higher in ESWPM alone (98.24) followed by ESWPM 100 per cent (92.37). This denotes the significant better performance than other treatments. Least RGR

values 62.94 and 54.3 were in ESWPM 40 and 20 per cent respectively. No negative impact on survival rate (%) was recorded in all the experimental diets fed fishes. Trend in FIFO was exact *i.e.*, the feed contains higher proportion of fish meal records higher values correctly.

From the present research, it was found that treatments, ESWPM alone and ESWPM replacing 100 per cent fish meal performed better than other experimental feeds. The technology got the patent from the Controller General of Patents, Designs & Trade marks, Chennai [No.136821 dt 18-12-2023]. However, the turbidity and the bad odour caused by ESWPM within few days after application need to be addressed.

REFERENCES

- Altomere A.A., Baron G., Aldini G., Carini M and Amato A. (2020) Silkworm pupae as source of high-value edible proteins and of bioactive peptides. Food Science and Nutrition 8: 2652–2661.
- Anon (1999) Final report (CFTRI, Mysore) of Research Project entitles Development of technology for the effective utilization of silkworm pupae with special reference to aquaculture feeds, grant-in-

- aid project by the Department of Biotechnology, Ministry of Science and Technology, Government of India, Project no. GAP-0084, April 1996-Sept 1999.
- Bandlamori S.V., Mondal M., Singh C.R and Karkada A.M. (2012) Evaluation of nutritional composition of hybrids of waste silkworm pupa *Bombyx mori* L. as a potential raw material for poultry feed-A sustainable technology for future. International Journal of Engineering Research and Technology 1: 1–5.
- Chakrabarthy R.D., Sen P.R. and Kowtal G.V. (1973) Observations on the relative usefulness of different feeds for carp spawn and fry. Journal of Inland Fishery Society of India 5: 182–188.
- Daniel N. (2018) A review on replacing fish meal in aqua feeds using plant protein sources. International Journal of Fisheries and Aquatic Studies 6(2): 164–179.
- FAO (2014) In: Graziano da Silva, J. (Ed.), The state of World Fisheries and Aquaculture, Opportunities and Challenges. FAO, Rome. p3.
- Fines B.C. and Holt G.J. (2010). Chitinase and apparent digestibility of chitin in the digestive tract of juvenile cobia, *Rachycentron canadum*. Aquaculture 303: 34–39.
- Gasco L., Finke M. and Van Huis A. (2018) Can diets containing insects promote animal health. Journal of Insects as Food and Feed 4(1): 1–4.
- Harikrishnan R., Kim J.S., Balasundaram C. and Heo M.S. (2012) Dietary supplementation with chitin and chitosan on haematology and innate immune response in *Epinephelus bruneus* against *Philasterides dicentrarchi*. Experimental Parasitology 131: 116–124.
- Henry M.A., Gasco L., Chatzifotis S. and Piccolo G. (2018) Does dietary insect meal affect the fish immune system? The case of mealworm, *Tenebrio molitor* on European sea bass, *Dicentrarchus labrax*. Developmental and Comparative Immunology 81: 204–209.
- Hoffman J., Johansen A., Steiro K., Gildberg A., Stenberg E and Bogwald J. (1997) Chito oligosaccharides stimulate Atlantic salmon, *Salmo salar* L., head kidney leukocytes to enhanced superoxide anion production in vitro. Comparative Biochemistry Physiology. Part B, Biochemistry and Molecular Biology 118(1): 105–115.
- Hora S.L. and Pillay T.V.R. (1962) Handbook on fish culture in the Indo-Pacific fisheries Region. FAO, Biology Technical Paper 14: 203 pp.
- Jayaram M.G. and Shetty H.P.C. (1980a) Digestibility of two pelleted diets by *Cyprinus carpio* and *Labeo rohita*, Mysore Journal of Agricultural Sciences 14: 578–584.
- Jayaram M.G. and Shetty H.P.C. (1980b) Studies on the growth rates of catla, rohu and common carp fed on different formulated feed, Mysore Journal of Agricultural Sciences 14: 598–606.
- Ji H., Zhang J.L., Huang J.Q., Cheng X.F and Liu C. (2015) Effect of replacement of dietary fish meal with silkworm pupae meal on growth performance, body composition, intestinal protease activity and health status in juvenile Jian carp (*Cyprinus carpio* var. Jian). Aquaculture Research 46: 1209–1221.
- Kaushik S.J. and Seiliez I. (2010) Protein and amino acid nutrition and metabolism in fish: current knowledge and future needs. Aquaculture Nutrition 41: 322–332.
- Kim S.K. and Rajapakse N. (2005) Enzymatic production and biological activities of chitosan oligosaccharides (COS): A review. Carbohydrate Polymers 62(4): 357–368.
- Li S., Ji H., Zhang B., Zhou J and Yu H. (2017) Defatted black soldier fly (*Hermetia illucens*) larvae meal in diets for juvenile Jian carp (*Cyprinus carpio* var. Jian): Growth performance, antioxidant enzyme activities, digestive enzyme activities, intestine and hepatopancreas histological structure. Aquaculture 477: 62–70.
- Lin S., Mao S., Guan Y., Lin X and Luo L. (2012) Dietary administration of chito oligosaccharides to enhance growth, innate immune response and disease resistance of *Trachinotus ovatus* Fish Shellfish Immunology 32(5): 909–913.
- Lindsay G.J.H., Walton M.J., Adron J.W., Fletcher T.C., Cho C.Y and Cowey C.B. (1984) The growth of rainbow trout (*Salmo gairdneri*) given diets containing chitin and its relationship to chitinolytic enzymes and chitin digestibility. Aquaculture 38: 315–334.
- Manikandavelu D., Neethiselvan, Jagatheeshan G. and Sundarsanam K. (1992) Use of fermented ensilage-based diet in enhancing the growth of common carp (*Cyprinus carpio* var. *communis*). Fish Technology 29(2): 111–113.
- Mrinal Ray M. and Gangopadhyay D. (2021). Effect of maturation stage and sex on proximate, fatty acid and mineral composition of eri silkworm from

- India. Journal of Food Composition and Analysis 100: 103898.
- Nandeesha M.C., Gangadhara B., Varghese T.J and Keshavanath P. (2000) Growth response and flesh quality of common carp, *Cyprinus carpio* fed with high levels of non-defatted silkworm pupae. Asian Fish Society 13: 235–242.
- Nandeesha M.C., Srikanth G.K., Keshavanath P., Varghese T.J., Basavaraja N and Das S.K. (1990) Effects of non-defatted silkworm-pupae in diets on the growth of common carp, *Cyprinus carpio*. Biological Waste 33: 17–23.
- Nandeesha M.C., Shikanth G.K., Varghese, T.J., Keshavanth P and Shetty H.P.C. (1989a) Growth performance of Indian major carp *Catla catla* (Ham) on fish meal free diets. In: S.S. De Silva (ed). Finfish nutrition Research in Asia. Proceedings of the third Asian Fish Nutrition Network Meeting, Asian Fisheries Society Special Publication No.4. Asian fisheries society, Manila, Philippines. pp 137–142.
- Nandeesha M.C., Shikanth G.K., Varghese T.J., Keshavanth P and Shetty H.P.C. (1989b) Influence of silkworm pupae-based diets on growth, organoleptic quality and biochemical composition of catla-rohu hybrid. In: E.A. Huisman, N. Zonneveld and Bouwmans, A.H. M: (eds), Aquaculture Research in Asia: Management Techniques and Nutrition. Pudoc. Press, Wageningen. pp211–220.
- Rana K.J., Siriwardena S. and Hasan M.R. (2009) Impact of rising feed ingredient prices on aquafeeds and aquaculture production. In: FAO (Ed.), Fisheries and Aquaculture Technical Paper. Food, Agriculture Organization of the United Nations, Rome, Italy. pp63.
- Rangacharyulu P.V., Giri S.S., Paul B.N., Yashoda K.P., Rao R.J., Mahendrakar N.S., Mohanty S.N and Mukhopadhyay P.K. (2003) Utilization of fermented silkworm pupae silage in feed for carps. Bioresources Technology 86: 29–32.
- Sadasivam S. and Manickam A. (2005) Biochemical Methods. New age International Pvt. Limited, New Delhi.
- Su J., Gong Y., Cao S., Lu F., Han D., Liu H and Xie S. (2017) Effects of dietary *Tenebrio molitor* meal on the growth performance, immune response and disease resistance of yellow catfish (*Pelteobagrus fulvidraco*). Fish Shellfish Immunology 69: 59–66.
- Van Huis A. (2013) Potential of insects as food and feed in assuring food security. Annual Review of Entomology 58: 563–583.
- Wan A.H., Snellgrove D.L. and Davies S.J. (2017) A comparison between marine and terrestrial invertebrate meals for mirror carp (*Cyprinus carpio*) diets: Impact on growth, haematology and health. Aquaculture Research 48(9): 5004–5016.
- Wilson R.P. (2002) Protein and amino acids. In: Halver, J.E., Hardy, R.W. (Eds.), Fish Nutrition. Academic press, London. eBook 9780080494920.
- Zegeye M.D. (2020) Nutritional Evaluation of Insect's Pupae-Larvae and its Utilization in Poultry Compound Feed. The Open Agriculture Journal 14: 1–8.

(Received July 07, 2023; revised ms accepted October 21, 2023; published December 31, 2023)

Record of *Anoplocnemis phasianus* (Fabricius, 1781) (Hemiptera, Heteroptera, Coreidae) from Goa, India

Vijaykumar S. Gadekar[#], Ayesha E. Shetkar and Aishwarya S. Naik^{*}

Department of Zoology, ADK Science College, Dodamarg, Sindhudurg 416602, Maharashtra, India

[#]Department of Zoology, Sangola College, Sangola, Solapur 413307, Maharashtra, India

Email: aishwaryanaik2002@gmail.com

ABSTRACT: In the survey on diversity of Coreidae, *Anoplocnemis phasianus* (Fabricius, 1781) is recorded for the first time in the state of Goa. External morphology of *A. phasianus* is described with its present geographical distribution, taxonomic photo plate, host plants, and natural photographs of the nymphs and adults are provided. © 2023 Association for Advancement of Entomology

KEY WORDS: Survey, diversity, host plant, morphology

Coreidae, commonly known as squash bug or leaf-footed bug, are medium to large sized bugs with four jointed antennae; some species in this family are bright colored with a head that is narrower than the pronotum, a four-segmented beak, front wings with veins, and three-segmented tarsi; extended hind tibiae in some species form a leaf like appearance (Gupta *et al.*, 2012). Their repugnatory glands release an unpleasant odour (Moody, 1930) that are supporting the defensive mechanisms against the predator species. They feed on cucurbits like squash and pumpkin, but some of them are pests of different agricultural crops (Bonjour and Fargo, 1989; Bonjour *et al.*, 1990). While conducting a bug survey in Goa, nymphs and adults of *Anoplocnemis phasianus* (Fabricius) were observed on *Senna obtusifolia* (Linn.) and it was identified by using literature of British Fauna of India (Distant, 1902), this is the first record of this taxon in Goa. Young shoots, flowers, leaves, and stems of *S. obtusifolia* were found to be infested frequently.

Anoplocnemis phasianus (Fabricius, 1781) (Plate 1: figs. 1-7; Plate 2: figs. 1-12)

Lygaeus phasiana, Fabricius 1781, *Spec. Ins*, 2: p 361

Lygaeus grossipes, Fabricius 1803. *Syst. Rhyng.*, 2: p 205.

Cerbus tumidipes, Herrich- Shaeffer 1842, *Wanz. Ins.*, 6: p.54.

Mictis punctum, affinis, bicolor. Westwood 1842, *in Hope Cat*, 2: p. 10.

Anoplocnemis phasiana: Distant 1902, *Fauna. Brit. Ind.*, 1:p. 346.

Specimens examined:

Male, 13. viii. 2021, Verna (North Goa), elevation (15m), coordinates (15°21'36" N; 73°55'44" E), Coll. Ayesha Shetkar, deposited in ADKS College, Dodamarg; Male, 18. viii. 2021, Verna (North Goa), elevation (15m), coordinates (15°21' 36" N; 73°55'44" E), Coll. Aishwarya Naik, deposited in

* Author for correspondence



Plate 1: Figures 1-7. *Anoplocnemis phasiana* (Fabricius). 1. Dorsal view of female (brown colour); 2. Dorsal view of male (brown colour); 3. Dorsal view of male (black colour); 4. Fourth instar nymph, female; 5. Fourth instar nymph, female; 6 to 7. Fifth instar nymphs on host plant.

ADKS College, Dodamarg; Female, 21. viii. 2022, Sal (North Goa), elevation (2m), coordinates (15°41'53" N; 73°55'47" E), Coll. Ayesha Shetkar; Female, 21. x. 2022, Kasarpal (North Goa), elevation (20 m), coordinates (15°38'44" N; 73°56'26" E), Coll. Parshuram Naik, deposited in ADKS College, Dodamarg. Host plant *Senna obtusifolia* (Linn.). Measurements (in mm). Male: total body length about 26.9 to 27.1 and Female: total body length about 24.8 to 25.

Diagnostic characters: (Male)

Head: Head dorsally black, longer than broad, pubescence, ventrally brownish to black; clypeus somewhat longer than paraclypei; antenniferous tubercle prominent and slightly overhanging the median lobe, widely separated; antennal segments black to brownish and setose; first three segments black; apical segment luteous at base and apex; antennal segments I stout and longer than segment

II; segments I and IV slightly subequal; segment III smallest; segments II and III stout and slender; segment IV spindle shaped; eyes large and globose with dark brown colour, ocelli very close to eyes with light brown colour; buccula black and short, extending beyond antenniferous tubercle; rostrum short and black, extending beyond procoxae.

Thorax: Pronotum black, densely granular and setose, and sloping towards the head, length shorter than width; anterior lobe narrow and acute, separated by transverse sulcus from broad posterior lobe; humeral angles rounded, ridge like appearance at the base; lateral margins obliquely straight; posterior margins slightly sinuate; base of pronotum moderately sloping downwards towards scutellum; scutellum triangular, somewhat acute at the apex covered with setae; corium and clavus brownish, covered with fine setae; corium and clavus brownish, covered with fine setae and covered




Plate 2: Figures 1-12. *Anoplocnemis phasianus* (Fabricius), 1-2 Dorsal view female (colour variation); 3-4. Dorsal view of male (colour variation); 5. Thorax, male; 6. Antennae; 7. Scutellum; 8. Thorax, female; 9. Rostrum; 10. Hind leg, dorsal view; 11-12. Ventral view of abdomen, male (colour variation)

with some apparent punctures; legs black; posterior femora strongly incrassated and curved in the middle with prominent spine and many minute spines; whole dorsal surface granular and setose; the entire surface of tibia slightly granular and setose; posterior tibiae much flattened; membrane brownish, with network like venation, which is not extending beyond the apical segment of abdomen.

Abdomen: Abdomen long, elongate and brownish to black; connexivum black and visible; sternum black covered with setae; a semicircular elevated ridge at segments II and III; spiracles large and prominent; the whole area setose, and covered with numerous fine black granules.

In the past, researchers on Coreidae species included Distant (1902), Basu and Mitra (1977 a, b and c, 1978, 1996, 2003, 2004) in India. Coreidae is distributed throughout the world, including countries and Islands such as Archipelago, China, India, Myanmar, Malay Peninsula, Nepal, Sri Lanka, and Thailand (Distant, 1902, 1908 and 1918; Schuh and Slater, 1995; Cassis and Gross, 2002; Dolling, 1991; Aukema *et al.*, 2013). There are over 1,802 species under 252 genera in the world (Schuh and Slater, 1995), out of which 143 species under 45 genera are in India (Distant, 1902, 1908, 1910). *Anoplocnemis phasianus* was noted in Assam, Kerala (Thiruvananthapuram), Madhya Pradesh,

Meghalaya (Khasi hills), Nagaland (Naga Hills), Sikkim, Tamil Nadu (Sholinganallur), Uttarakhand (Gwaldam-Garud Tehsil), and West Bengal (Darjeeling and Buxaduar, Jalpaiguri district). Although *A. phasianus* was recorded from adjacent states such as Maharashtra (Mumbai and Bor Ghat) and Karnataka (Distant, 1902). There is no report of *A. phasianus* from Goa till date.

ACKNOWLEDGEMENTS

The authors are extremely grateful to Mr. Rajesh Sutar, Dodamarg, who helped in the photo plate preparation for this article. The authors are thankful to the friends who helped in the collection of the bugs. The authors appreciate the facilities and encouragement provided by the ADK Science College, Dodamarg administration.

REFERENCES

- Aukema B., Rieger C. and Rabitsch W. (2013) Catalogue of the Heteroptera of the Palaearctic Region. Volume 6. Supplement. The Netherlands Entomological Society, Amsterdam. xxiv + 629pp.
- Basu R.C. and Mitra S.C. (1977a) New records of Coreidae (Insecta: Hemiptera) from West Bengal (India). Newsletter, Zoological Survey of India 3(4): 206–209.
- Basu R.C. and Mitra S.C. (1977b) New records of Coreidae (Insecta: Hemiptera) from Kameng division (Arunachal Pradesh), India. Newsletter, Zoological Survey of India 3(6): 415–418.
- Basu R.C. and Mitra S.C. (1977c) Some coreoidea (Insecta: Hemiptera) from Subansiri and Lohit districts of Arunachal Pradesh (NEFA), India. Zoological Survey of India 3: 431–433.
- Basu R.C. and Mitra S.C. (1978) New records of Coreidae (Insecta: Hemiptera) from Sikkim, India. Zoological Survey of India 44(9): 413–414.
- Basu R.C. and Mitra S.C. (1996) Insecta: Hemiptera: Coreoidea. Fauna of West Bengal, State Fauna Series 3(5): 449–467.
- Basu R.C. and Mitra S.C. (2003) Insecta: Hemiptera: Heteroptera: Coreoidea. Fauna of Sikkim, State Fauna Series 9(2): 489–508.
- Basu R.C. and Mitra S.C. (2004) Insecta: Hemiptera: Heteroptera: Coreoidea, Fauna of Manipur, State Fauna Series 10: 223–237.
- Bonjour E.L. and Fargo W.S. (1989) Host effects on the survival and development of *Anasa tristis* (Heteroptera: Coreidae). Environmental Entomology 18: 1083–1085.
- Bonjour E.L., Fargo W.S. and Rensner P.E. (1990) Ovipositional preference of squash bugs (Heteroptera: Coreidae) among cucurbits in Oklahoma. Journal of Economic Entomology 83: 943–947.
- Cassis G. and Gross G.F. (2002) Hemiptera: Heteroptera (Pentatomomorpha). In Zoological Catalogue of Australia, (eds) W.W.K. Houston and G.V. Maynard. Volume 27.3B., CSIRO, Melbourne, Australia. 737pp.
- Distant W.L. (1902) The Fauna of British India, including Ceylon and Burma. Rhynchota, 1 (Heteroptera). Taylor & Francis, London. 438pp.
- Distant W.L. (1908) The Fauna of British India, including Ceylon and Burma. Rhynchota, 4 (Heteroptera). Taylor & Francis, London. 446pp.
- Distant W.L. (1918) The Fauna of British India, including Ceylon and Burma. Rhynchota, 7 (Heteroptera). Taylor & Francis, London. 151pp.
- Dolling W.R. (1991) The Hemiptera. Oxford University Press, London. 274pp.
- Gupta R., Singh D., Kaur H and Kaur R. (2012) A survey report on family Coreidae (Hemiptera: Heteroptera) from North India. Annals of Entomology 30 (2): 11–16.
- Moody D.L. (1930) The morphology of the repugnatory glands of *Anasa tristis* De Geer. Annals of the Entomological Society of India 23: 81–104.pp
- Schuh R.T. and Salter J.A. (1995) True Bugs of the World (Hemiptera: Heteroptera). Classification and Natural History. Cornell University Press. 336pp.

Efficacy of insecticides against cashew leaf miner, *Acrocercops syngamma* Meyrick (Lepidoptera, Gracillariidae)

O. Chethana*, Jayalaxmi Narayan Hegde, K. Rajashekharappa, T. Basavaraj Naik¹ and B.V. Champa²

Department of Entomology, College of Agriculture, Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences (KSNUAHS), Navile, Shivamogga 577204, Karnataka, India.

¹Zonal Agricultural and Horticultural Research Station (ZAHRS), KSNUAHS, Navile, Shivamogga 577204, Karnataka, India.

²Organic Farming Research Centre, ZAHRS, KSNUAHS, Navile, Shivamogga 577204, Karnataka, India.

Email: chetan1091998@gmail.com

ABSTRACT: Experiments were conducted to know the efficacy of different insecticides (Cyantraniliprole 10.26 % OD @ 1.8 ml L⁻¹, Dinotefuran 20 % SG @ 0.3 g L⁻¹, Chlorfenapyr 10 % SC @ 1.5 ml L⁻¹, Imidacloprid 17.8 % SL @ 0.3 ml L⁻¹, Thiamethoxam 25 % WG @ 0.3 g L⁻¹, Lambda cyhalothrin 2.5 % EC @ 1 ml L⁻¹) against cashew leaf miner, *Acrocercops syngamma* Meyrick under field condition and under nursery condition. Among the different insecticides evaluated, under field condition, Lambda cyhalothrin was found to be statistically superior over all other treatments recording 83.39 per cent reduction over control. Imidacloprid was found least effective against leaf miner (46.41% reduction of over control). Under nursery condition, Lambda cyhalothrin recorded 79.13 per cent reduction whereas Imidacloprid recorded only 20.31 per cent. © 2023 Association for Advancement of Entomology

KEY WORDS: Field, nursery, lambda cyhalothrin, imidacloprid, leaf miner control

Leaf miner, *Acrocercops syngamma* Meyrick (Lepidoptera, Gracillariidae) is one of the important pests of cashew during the post-monsoon period all over the country. The larvae, after hatching from the eggs, start mining the epidermal layer on the upper surface of the tender cashew leaves as well as tender shoots. As a result of feeding, the affected area forms blistered patches of grayish white colour. When the infested leaves mature, the damage manifests as big holes. Young plants are observed to be more prone to attack by this pest

and up to 8 and 15 caterpillars have been observed on a single leaf. During the developmental period, leaf miner larvae are dull white and turn pinkish before pupation. After full development, the larvae fall off to the soil where they pupate and emerge after 7-9 days. The adult is a silvery grey moth that lays eggs on tender leaves (Athalye and Patil, 1999). Leaf miner is a defoliating pest of cashew, occurring in almost all the cashew-growing regions of the country as well as the world. In India, it causes serious damage to the tender leaves of cashew

* Author for correspondence

attacking 2-80 per cent of the young leaves (Ayyanna *et al.*, 1985). Upon hatching, the larva makes a silvery sinuous gallery on the upper leaf side feeding below the epidermal layer, causing leaf blisters which later dry up, causing leaf distortion, browning and curling of the leaves. As many as 11 larvae have been observed feeding on a single leaf (Jena *et al.*, 1985). The pest completes its life cycle in a short period of 20-22 days and spreads fast causing leaf blisters over a wide area (Rai, 1984). Over the decades, use of chemical insecticides has become the only line of defense to combat the major pests of cashew (Kar, 2017). Lambda cyhalothrin is a proven insecticide against tea mosquito bug and other associated pests. However, the recent advances in pest management are being directed towards the development of safer and effective insecticides which reduce the pesticide load in the environment.

Efficacy of insecticides under field conditions:

The experiment was conducted in field conditions at Zonal Agricultural and Horticultural Research Station (ZAHRS), Shivamogga, Karnataka in randomized complete block design (RCBD) with six insecticidal treatments (Cyantraniliprole 10.26 % OD @ 1.8ml L⁻¹, Dinotefuran 20 % SG @ 0.3 g L⁻¹, Chlorfenapyr 10 % SC @ 1.5ml L⁻¹, Imidacloprid 17.8 % SL @ 0.3ml L⁻¹, Thiamethoxam 25 % WG @ 0.3g L⁻¹, Lambda cyhalothrin 2.5 % EC @ 1ml L⁻¹) and a control, with three replications (one tree is one replication to evaluate insecticides for the management of cashew leaf miner. In each tree, 20 young leaves were randomly selected representing four directions, and in each directions five leaves three from top and two from middle canopy were selected. Number of leaf miner larvae per leaf were counted and averaged. Pre-treatment count a day before spraying and post treatment counts on three, five, seven and ten days after spraying were recorded. Two sprays were given during the peak incidence in December month. The data were subjected to statistical analysis.

Efficacy of insecticides at Nursery: The experiment was conducted in nursery of one and half year age plants at Agricultural and Horticultural Research Station (AHRS), Bavikere,

Chikkamagaluru, Karnataka with RCBD with seven treatments including control (as above) with three replications to evaluate insecticides for the management of cashew leaf miner. In each replication, there were ten plants. One spray was given during the peak incidence during November month. Five plants were selected randomly from each replication in each treatment. In each plant, two top leaves were selected randomly. Number of leaf miner larvae were counted and averaged. Pre-treatment count a day before spraying and post treatment counts on three, five, seven and ten days after spraying were recorded.

The statistical analysis of the data obtained from insecticidal efficacy trial was done using analysis of variance (ANOVA) using Web Agri Stat Package (WASP-2) developed by Indian Council of Agricultural Research, Research Complex, Goa. The numerical data were subjected to a square root transformation before subjecting to ANOVA. After analysis, data were fitted in the table as per the needs of objectives for interpretation of results and the data were correlated with the weather parameters following the methods of Gomez and Gomez (1984). The interpretation of data was done by using the critical difference value calculated at 0.05 probability level. Further, the per cent reduction of cashew leaf miner over control was calculated by following formula:

$$\text{Per cent reduction} = \frac{C_b - T_a}{C_b} \times 100$$

Where, C_b = number of insects in untreated control before insecticide application

T_a = number of insects in treated plot after insecticide application

Results of field trial revealed that the incidence of leaf miner (larvae per 20 leaves) ranged from 1.90 to 11.44 in all the treatments including control. The least larval population (1.90 larvae) was recorded in Lambda cyhalothrin treatment, followed by Cyantraniliprole (3.48 larvae), Dinotefuran 20 (4.01 larvae), Thiamethoxam (4.65 larvae) and Chlorfenapyr (5.00 larvae). Among the insecticides treated trees, highest mean larval population was

Table 1. Efficacy of insecticides under field condition against Leaf miner, *A. syngamma* on cashew at ZAHRS, Shivamogga during 2021 - 22

No.	Treatments	Dose per lt	Mean number of leaf miner larvae/20 leaves										Reduction (%)
			First Spray					Second Spray					
			1	3	5	7	10	3	5	7	10	Mean	
1	Cyantraliprole 10.26 OD	1.8ml	12.80 (3.64)	6.89 (2.71) ^c	5.91 (2.52) ^{cd}	4.55 (2.22) ^c	3.11 (1.88) ^{cd}	3.01 (1.87) ^d	1.90 (1.54) ^c	1.17 (1.29) ^{de}	1.30 (1.33) ^c	3.48	69.58
2	Dinotefuran 20 SG	0.3g	12.53 (3.61)	6.96 (2.72) ^c	6.50 (2.62) ^{bcd}	7.31 (2.77) ^b	3.51 (1.97) ^c	3.21 (1.92) ^{cd}	2.41 (1.69) ^c	1.73 (1.49) ^{cd}	0.45 (0.97) ^e	4.01	64.94
3	Chlorfenapyr 10 SC	1ml	13.27 (3.71)	8.79 (3.05) ^{bc}	8.35 (2.97) ^{bc}	7.48 (2.81) ^b	4.45 (2.21) ^{bc}	4.10 (2.14) ^{cd}	3.23 (1.91) ^{bc}	2.47 (1.71) ^{bc}	1.12 (1.27) ^{cd}	5.00	56.29
4	Imidacloprid 17.8 SL	0.3ml	12.80 (3.65)	9.33 (3.12) ^b	9.41 (3.14) ^{ab}	8.53 (2.99) ^b	6.32 (2.61) ^b	5.91 (2.52) ^b	4.16 (2.16) ^b	3.56 (1.99) ^b	1.80 (1.51) ^b	6.13	46.41
5	Thiamethoxam 25 WG	0.3g	12.93 (3.66)	8.57 (3.01) ^{bc}	7.23 (2.73) ^{bc}	7.22 (2.77) ^b	4.63 (2.24) ^{bc}	4.23 (2.71) ^c	2.45 (1.71) ^c	1.99 (1.58) ^{cd}	0.85 (1.16) ^d	4.65	59.35
6	Lambda cyhalothrin 2.5 EC	1ml	12.67 (3.62)	4.90 (2.30) ^d	4.12 (2.14) ^d	2.52 (1.71) ^d	1.32 (1.35) ^d	1.14 (1.23) ^e	0.72 (1.10) ^d	0.35 (0.92) ^e	0.10 (0.77) ^f	1.90	83.39
7	Control	-	14.00 (3.81)	13.25 (3.70) ^a	12.85 (3.65) ^a	12.53 (3.60) ^a	10.80 (3.36) ^a	10.91 (3.37) ^a	10.75 (3.35) ^a	10.54 (3.32) ^a	9.89 (3.22) ^a	11.44	
SEm ±			-	0.122	0.179	0.143	0.188	0.096	0.126	0.128	0.054		
CD at 0.05			NS	0.366	0.539	0.431	0.566	0.290	0.380	0.385	0.162		
CV (%)			5.918	6.979	10.727	8.983	14.245	7.499	11.116	12.330	6.213		

Note: DBS - Days Before Spray, DAS - Days After Spray, Figures in parenthesis are $\sqrt{x+0.5}$ transformed values;

Means in the columns followed by the same alphabet do not differ significantly by DMRT (P = 0.05).

recorded in Imidacloprid (6.13) sprayed trees. In untreated control, there were 11.44 larvae per 20 leaves. Leaf miner reduction of over control was maximum in Lambda cyhalothrin (83.39%), followed by Cyantraliprole (69.58%) and Dinotefuran (64.94%), Thiamethoxam (59.35%) and Chlorfenapyr (56.29%). The least was recorded in Imidacloprid (46.41%) among the insecticides (Table 1).

In the nursery experiment, the incidence ranged from 1.88 to 9.01 larvae per 10 leaves. Among the insecticides, the least mean number was recorded in Lambda cyhalothrin treatment (1.88 larvae per 10 leaves), followed by Cyantraliprole (3.11), Dinotefuran (4.55), Thiamethoxam (4.77) and Chlorfenapyr (5.39). The maximum mean number was recorded in Imidacloprid (7.18). In untreated

control it was 9.01. The reduction leaf miner larvae over the control, was highest in Lambda cyhalothrin (79.13%), followed by Cyantraliprole (65.48%), Dinotefuran (49.50 %), Thiamethoxam (47.05%) and Chlorfenapyr (40.17%). The least was recorded in Imidacloprid (20.31%) among the insecticides (Table 2). Overall, among the insecticides evaluated, Lambda cyhalothrin 2.5 EC @ 1 ml L⁻¹ was found most effective in managing the leaf miner, *A. syngamma* in cashew with respect mean number of larvae and percent reduction over the control both in field and nursery conditions.

These results are in line with Kar (2017) and Patel *et al.* (2018) who reported that out of different insecticides evaluated against leaf miner of cashew, Lambda cyhalothrin 5 EC (@ 0.003%) was most efficient which recorded minimum leaf damage in

Table 2. Efficacy of insecticides against Leaf miner, *Acrocercops syngamma* in cashew nursery at AHRS, Bavikere during 2021

No.	Treatments	Dosage per L	Mean no. of leaf miner larvae/10 leaves						Reduction over control %
			1DBS	3DAS	5DAS	7DAS	10DAS	Mean	
1	Cyrantriliprole 10.26 OD	1.8ml	6.99 (2.68)	5.34 (2.38) ^{bc}	3.21 (1.91) ^{de}	2.61 (1.75) ^{cd}	1.28 (1.33) ^d	3.11	65.48
2	Dinotefuran 20 SG	0.3g	9.12 (3.10)	8.59 (3.01) ^a	3.88 (2.07) ^{ede}	3.22 (1.92) ^c	2.52 (1.74) ^c	4.55	49.50
3	Chlorfenapyr 10 SC	1ml	8.09 (2.92)	7.59 (2.83) ^{ab}	5.93 (2.53) ^{bc}	4.16 (2.16) ^{bc}	3.88 (2.06) ^{bc}	5.39	40.17
4	Imidacloprid 17.8 SL	0.3ml	7.59 (2.83)	8.93 (3.06) ^a	8.59 (3.01) ^{ab}	6.29 (2.56) ^{ab}	4.90 (2.32) ^b	7.18	20.31
5	Thiamethoxam 25 WG	0.3g	7.64 (2.85)	6.49 (2.58) ^{abc}	5.32 (2.40) ^{cd}	4.23 (2.17) ^{bc}	3.05 (1.87) ^c	4.77	47.05
6	Lambda cyhalot hrin 2.5 EC	1ml	6.93 (2.72)	4.22 (2.16) ^c	1.87 (1.53) ^e	1.10 (1.26) ^d	0.33 (0.91) ^e	1.88	79.13
7	Control	-	8.01 (2.90)	9.18 (3.10) ^a	9.22 (3.12) ^a	9.08 (3.09) ^a	8.55 (3.00) ^a	9.01	
SEm ±			-	0.193	0.185	0.179	0.128		
CD at 0.05			NS	0.580	0.556	0.538	0.386		
CV (%)			11.68	11.928	13.216	14.198	11.486		

Note: DBS - Days Before Spray; DAS - Days After Spray; Figures in parenthesis are $\sqrt{x+0.5}$ transformed values; Means in the columns followed by the same alphabet do not differ significantly by DMRT ($P = 0.05$).

the field. In nursery also, the experiment results are in line with Kar (2017) and Patel *et al.* (2018) who reported that Lambda cyhalothrin 5 EC (@ 0.003%) was effective in minimizing the leaf damage. Overall, among the insecticides evaluated both in field and nursery conditions, Lambda cyhalothrin 2.5 EC @ 1ml L⁻¹ was found most effective in managing the leaf miner, *A. syngamma* in cashew with respect mean number of larvae and percent reduction over the control.

REFERENCES

- Athalye S.S. and Patil R.S. (1999) Bionomics, seasonal incidence and chemical control of cashew leaf miner. *Journal of Maharashtra Agricultural Universities* 23: 29–23.
- Ayyanna T., Tejkumar S. and Ramadevi (1985) Distribution and status of pests of cashew in coastal districts of Andhra Pradesh. *Cashew Causerie* 7: 4–5.
- Gomez K.A. and Gomez A.A. (1984) Statistical procedures for agricultural research, 2nd addition. An International Rice Research Institute Book, A Wiley - Inter science Publication, New York. pp20–29, 382–387.
- Jena B.C., Patnaik N.C. and Satapathy C.R. (1985) Insect pests of cashew. *Cashew Causerie* 7(3): 10–11.
- Rai P. (1984) Seasonal distribution of cashew in coastal Karnataka. *Indian Cashew Journal* 13(4): 17–19.
- Kar A. (2017) Evaluation of few insecticides against insect pests of cashew. *Journal of Entomology and Zoology Studies* 5(2): 1345–1347.
- Patel R.B., Patel D.R., Makati J.P. and Patel R.R. (2018) Evaluation of insecticides against pest complex of cashew. *International Journal of Economic Plants* 5(1): 36–39.

(Received August 03, 2023; revised ms accepted October 10, 2023; published December 31, 2023)

Biology of shoot and fruit borer, *Earias vittella* Fabricius (Noctuidae, Lepidoptera) on okra

Bikash Subba¹, Nilanjana Chudhuri² and S.K. Senapati³

¹Department of Entomology School of Agriculture, Institute of Technology Management, Gwalior 474001, Madhya Pradesh, India.

²Department of Agricultural Entomology, Regional Research Station, Terai Zone, Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar 736165, West Bengal, India.

³Department of Agricultural Entomology, Uttar Banga Krishi Viswavidyalaya, Pundibari 736165, Cooch, Behar, West Bengal, India.

Email: bsubba097@gmail.com

ABSTRACT: The biology of fruit and shoot borer (*Earias vittella* F.) was studied under the natural conditions in the instructional farm of Uttar Banga Krishi Viswavidyalay in *pre-kharif*, *kharif* and *post-kharif* on okra. The study revealed that the egg incubation in *pre-kharif*, *kharif* and *post-kharif* revealed that the egg incubation were 3.18 ± 0.40 , 3.61 ± 0.70 and 4.25 ± 0.67 and larval period were 11.00 ± 0.88 , 11.00 ± 0.82 and 13.01 ± 1.81 during corresponding seasons respectively. The pupal period in *pre-kharif*, *kharif* and *post-kharif* were 13.01 ± 1.81 ; 11.00 ± 0.82 ; 10.97 ± 2.15 days respectively. The adult male longevities observed were 3.64 ± 0.67 , 5.60 ± 0.70 and 7.33 ± 0.97 during *pre-kharif*, *kharif* and *post-kharif*. On the other hand female longevities observed were about 7.45 ± 1.57 ; 8.87 ± 1.10 ; 9.41 ± 1.51 days in corresponding seasons respectively. The female fecundity during *pre-kharif*, *kharif* and *post-kharif* were 91.55 ± 10.93 , 117.55 ± 10.60 and 132.04 ± 5.83 eggs in their lifetime. The pre-oviposition during *pre-kharif*, *kharif* and *post-kharif* were 1.64 ± 0.50 , 1.52 ± 0.53 and 1.88 ± 0.57 and the oviposition periods in corresponding seasons were 2.73 ± 0.47 , 3.22 ± 0.79 and 4.32 ± 0.52 . While post-oviposition periods in *pre-kharif*, *kharif* and *post-kharif* were 3.09 ± 0.70 , 4.14 ± 0.63 and 3.23 ± 0.92 respectively. Its total life-cycle was completed in 27.45 ± 1.57 , 31.60 ± 1.90 and 41.35 ± 2.55 days during *pre-kharif*, *kharif* and *post-kharif* respectively. Higher fecundity may lead to faster population growth that may surpass the economic threshold level (ETL), even though the shoot and fruit borer's life cycle is longer in *kharif* and *post-kharif*. Thus, it may be said that the *kharif* and *pos-kharif* seasons are the most susceptible to shoot fruit and shoot borer infestation.

© 2023 Association for Advancement of Entomology

KEY WORDS : Life cycle, seasons, susceptibility, population growth

Okra, *Abelmoschus esculentus* (L.) is a native of tropical or sub-tropical Africa and belongs to the Family Malvaceae. In India, okra is cultivated around the year in *pre-kharif*, *kharif* and *post-kharif* seasons. Okra is attacked by several species of insect pests and infected by a few diseases from seedling to harvesting. Economic losses depend on the degree of damage, pest density, environmental

condition, stage of growth and the plant part damaged by the pest. Studying the life cycle of insect pests is vital for effective pest management, sustainable agriculture, environmental protection, and reducing the economic impact of pest damage. It serves as the cornerstone for making well-informed decisions about pest control tactics and aids in balancing the need for pest control with

* Author for correspondence

ecological and economic considerations. Among different insect pests infesting okra in the terai region of West Bengal, fruit borer and jassid are considered key pests since they are causing regular menace to okra in huge amounts. Therefore, an attempt has been made to study the biology of the fruit and shoot borer (*Earias vittella* F.) as influenced by the seasons.

A laboratory culture of *E. vittella* was started with several infested okra pods collected from the field which were transferred into plastic jars covered with muslin cloth and kept in laboratory conditions till pupation and adult emergence. After mating adults were shifted to potted okra plants with young fruit and covered with net for egg laying. The egg masses were examined daily and the incubation period was recorded. On hatching, the larva was monitored until the end of the last instar. The full-grown larva bores a large exit hole, leaves the okra pod, spins a cocoon and pupates on the outer surface of the fruit. The duration of the larval and pupal stages was determined. For studying adult longevity, 10 pairs of newly emerged males and females were maintained in glass vials with 10% sugar solution in absorbent cotton. The time elapsed between the emergence of each moth and its death was recorded as the longevity of such an individual. The duration of each generation was estimated based on the average length of the life cycle. The duration of pre-oviposition, oviposition, and post-oviposition were recorded for the female. The cage was opened daily and the leaves and fruits were observed for oviposition with the help of magnifying lens and this was continued till the last egg was laid. SAS software (ver. 9.2) was used for data analysis. One-way ANOVA was performed for each of the parameters and separation of the means was done using the Least Significant Difference (LSD) test at 5% significant level.

A. Incubation period

The incubation period varied in different crop seasons. The highest incubation period of 4.25 ± 0.67 days was recorded in *post-kharif* followed by 3.61 ± 0.67 days in *kharif* and the shortest duration (3.18 ± 0.40 days) was recorded in *pre-kharif*. This

result is in agreement with Rehman and Ali (1981) in January to March; Singh and Bichoo (1989) in September to October; Raju (2016) in *kharif* and Nishi (2016) in March-April who reported that the incubation period lasted for 3 to 4 days. The incubation period was 3.27 and 3.67 days as reported by Kumar *et al.* (2014) during July-August and October-November supports our study. At par with the present result for the *post-kharif* season, Das and Chaudhuri (2012) recorded 4.50 days of incubation period in October-November. A similar observation of 4.57 days and 5.00 days was also observed by Sundraraj and David (1987) and Suryawanshi *et al.* (2001). Bhat *et al.* (2005) recorded 2.60-3.80 days during June-August and 4.40-4.80 days during September-November which supports the present observation. The results under present studies as well as other reports as discussed above contradict Al-Mehmmady (2000) who found shorter incubation of 2.15 and 2.42 days in August and September. In line with Al-Mehmmady (2000) and Syed *et al.* (2011) also found 2.30 days of incubation period.

B. Larval period

The larva is the most injurious stage and therefore, their growth and development is very important, as it directly affects fruit production. Caterpillar passed through five stages. The duration of different larval instars varied with seasons. Total larval periods of 11.00 ± 0.88 ; 12.25 ± 1.20 and 17.90 ± 1.17 days were recorded in *pre-kharif*, *kharif* and *post-kharif* respectively. In conformity with the present study during *post-kharif* Rukhsana *et al.* (1995), Suryawanshi *et al.* (2001) and Das and Chaudhuri (2012) also recorded larval periods of 18.00 ± 0.88 , 19.00 and 16.66-23.33 days respectively. Other workers reported similar result where the period lasted for 9.30 days (Ambegankar and Billapate, 1984); 12.73 days (Sundraraj and David, 1987); 11.28-11.39 days (Al-Mehmmady, 2000); 10-12 days (David, 2002); 7-15 days (Srivastava, 2003); 9.20-11.20 days (Dhillon and Sharma 2004); 8.20-9.00 days in June-August and 9.80-12.20 days in September to November (Bhat *et al.*, 2005); 9.16 days (Syed *et al.*, 2011); 8.00 days in March-August (Shah *et al.*, 2012); 11.79 days in July-

Table 1: Duration of different developmental stages of *Earias vitella* on okra over seasons

Life stages	Duration in days (Mean±SD)			F	Pr>F	LSD
	<i>Pre-Kharif</i>	<i>Kharif</i>	<i>Post-Kharif</i>			
Mean temperature (Min-max)	25.08°C (20.60-31.55 °C)	28.08 °C (23.65-32.50 °C)	27.37 °C (21.92-32.82 °C)			
Mean RH (Min-Max)	74.28% (72.97-79.71 %)	84.26% (78.69-87.83 %)	78.45% (74.73-82.16%)			
Incubationperiod	3.18±0.40b	3.61±0.70b	4.25±0.67a	3.68±0.57	8.58	0.0020 0.539
Larval period						
1st instar	1.09±0.30c	1.60±0.52b	2.15±0.42a	1.61±0.56	17.69	<.0001 0.370
2nd instar	2.00±0.09b	2.03±0.20b	3.04±0.57a	2.36±0.64	30.07	<.0001 0.318
3rd instar	2.18±0.40b	2.34±0.48b	3.61±0.67 a	2.71±0.85	21.84	<.0001 0.494
4th instar	2.36±0.50c	2.85±0.42b	4.76±0.32a	3.32±1.36	68.11	<.0001 0.454
5th instar	3.36±0.50b	3.43±0.70b	4.35±0.52a	3.71±0.59	17.45	<.0001 0.387
Total Larva	11.00±0.88b	12.25±1.20b	17.90±1.17a	13.80±3.94	51.74	<.0001 1.508
Pupal Period	8.91±0.54c	11.00±0.82b	13.01±1.81a	10.97±2.15	32.63	<.0001 1.059
Adult longevity						
Male	3.64±0.67c	5.60±0.70b	7.33±0.97a	5.52±1.91	58.27	<.0001 0.714
Female	7.45±1.57b	8.87±1.10a	9.41±1.51a	8.58±2.05	7.10	0.0047 1.231
Life cycle	27.45±1.57a	31.60±1.90b	41.35±2.55a	33.46±1.70	67.10	<.0001 2.568
Fecundity	91.55±10.93c	117.55b±10.60b	132.04±5.83a	113.72±21.33	54.71	<.0001 8.183
Pre-oviposition	1.64±0.50a	1.52±0.53a	1.88±0.57a	1.68±0.42	1.42	0.2648 0.458
Oviposition	2.73±0.47	3.22±0.79	4.32±0.52	3.42±0.56	20.47	<.0001 0.528
Post-oviposition	3.09±0.70b	4.14±0.63a	3.23±0.92a	3.48±0.19	5.51	0.0124 0.717

* Within row means followed by the same letter(s) are not significantly different at 5% level

August and 11.15 days in October-November (Kumar *et al.*, 2014) and 11.50±1.08 days in March-April (Nishi, 2016). The newly hatched larvae wandered about for few hours before boring into the fruit. The period of different larval instars were 1.09±0.30, 2.00±0.09, 2.18±0.40, 2.36±0.50 and 3.36±0.50 days in *pre-kharif*; 1.60±0.52, 2.03±0.20, 2.34±0.48, 2.85±0.42 and 3.43±0.70 days in *kharif*; 2.15±0.42, 3.04±0.57, 3.61±0.67, 4.76±0.32 and 4.35±0.52 days in *post-kharif* season respectively. The average duration of different larval stages in the three seasons were 1.61±0.56, 2.36±0.64, 2.71±0.85, 3.32±1.36 and 3.71±0.59 days. This finding was at par with Sewak (2016) where the

average duration of 1st, 2nd, 3rd, 4th and 5th larval instars was 1.60±0.52, 2.00±0.00, 2.50±0.53, 2.50±0.53 and 3.00±0.00 days respectively. In support of the present finding during *post-kharif*, Das and Chaudhuri (2012) recorded 3.33, 3.00, 3.00, 3.33 and 4.00 days of development period of different larval instars during October-November.

C. Pupal period

At the end of its development, the larva leaves the fruits and settles down to spin their cocoons at different places outside the okra fruits. Pupation occurred in a dirty white boat-shaped cocoon. The average pupal period was 10.97±2.15 days in three

seasons. The longest period was observed in *post-kharif* (13.01 ± 1.81 days) followed by *kharif* (11.00 ± 0.82 days) and the shortest (8.91 ± 0.54 days) was in *pre-kharif*. In support of the present findings other workers recorded a period of 6-14 days (Rehman and Ali, 1981 and Singh and Bichoo, 1989); 11.16 days (Sundraraj and David, 1987); 6.45 and 7.78 days during August-October (Al-Mehmmady, 2000); 10.00 days (Suryawanshi *et al.*, 2001); 7-10 days (David, 2002); 7.8-8.6 days during June-August and 9.8-10.2 days during September-November (Bhat *et al.*, 2005); 10.00 in July and 11.80 days in September (Syed *et al.*, 2011) and 8.0 ± 0.82 days (Nishi, 2016). The pupal period lasted for 8.50-9.50 days in October-November as reported by Das and Chaudhuri (2012) deviates from the present work, particularly for *post-kharif* crop.

D. Adult stage

The duration of different stages of *Earias* adult life is presented (Table 1). The adult longevity was highest in *post-kharif* (7.33 ± 0.97 days for males and 9.41 ± 1.51 days for females) and it was supported by the findings of Suryawanshi *et al.* (2001) where it was 10 days and Das and Chaudhuri (2012) with 9.16 days of adult longevity. The male and female adults during *kharif* lived for 5.60 ± 0.70 and 8.87 ± 1.10 days and it was 3.64 ± 0.67 and 7.45 ± 1.57 days in *pre-kharif* confirms the report of Shah *et al.* (2012) and (Nishi, 2016) who recorded the period as 6-12 days and 9-14 days and 4.2 days and 9.5 days for male and female respectively. The average life span of male and female moths in three generations was 5.52 ± 1.91 days and 8.52 ± 2.05 days respectively. However, longer male and female adult longevity was recorded by different workers and the duration was 9.25 and 13.91 days (Rehman and Ali, 1981); 10.76 and 14.60 days (Sundraraj and David, 1987); 12.45 and 14.00 days during August and 13.36 and 14.20 days during October (Al-Mehmmady, 2000) and 13.9 and 14.2 days in July, 11.66 and 13.3 days in September and 8.0 and 11.8 days in October (Syed *et al.*, 2011). This might be due to variations in regional climatic conditions influencing the activity of the insect.

The pre-oviposition, oviposition and post-oviposition period were 1.64 ± 0.50 , 2.73 ± 0.47 and 3.09 ± 0.70 days in *pre-kharif*; 1.52 ± 0.53 , 3.22 ± 0.79 and 4.14 ± 0.63 days in *kharif* and 1.88 ± 0.57 , 4.32 ± 0.52 and 3.23 ± 0.92 days in *post-kharif* with an average of 1.68 ± 0.42 , 3.42 ± 0.56 and 3.48 ± 0.19 days respectively. Das and Chaudhuri (2012) also recorded the 1.00-2.00 days of pre-oviposition, 5.00-5.15 days of oviposition and 2.00-2.05 days of post-oviposition period in October-November. The earlier work by Bhat *et al.* (2005) revealed that the pre-oviposition period varied from 1.8 to 2.44 days in August to 3.6 ± 0.54 days in November, whereas the oviposition and post-oviposition period ranged from 2.2 ± 0.44 (August) to 4.2 ± 0.44 in November and 8.4 ± 0.54 in June to 12.2 ± 0.83 in November. However, Suryawanshi *et al.* (2001) recorded a longer oviposition period of 7 days and the post-oviposition period was only 1.00 days. In support of the *pre-kharif* result, Nishi (2016) also obtained 1.8 ± 0.78 , 2.8 ± 0.79 and 4.5 ± 0.53 days of pre-oviposition, oviposition and post-oviposition period during March-April while Rehman and Ali (1981) reported 3.5, 5.83 and 4.75 days respectively. The corresponding values were 1.5, 7.0 and 10.0 in July, 0.66, 5.0 and 3.0 in September and 1.75, 4.5 and 2.0 in October respectively as recorded by Syed *et al.* (2011). However, Shah *et al.* (2012) reported 6-8 days of egg laying period.

E. Life cycle

The duration of the life cycle varied with the seasons. In *post-kharif*, the duration was longest (41.35 ± 2.55 days) followed by *kharif* (31.60 ± 1.90 days) and shortest (27.45 ± 1.57 days) in *pre-kharif*. The present findings are in close conformity to Al-Mehmmady (2000) who reported life cycle of 39.00 to 49.40 days, 31.82 to 36.59 days and 35.40-39.64 days during February-May, June-August and September to December. Sharma *et al.* (1985) recorded the duration ranged from 29 to 49 days which supports the present result. The period of 32.50 ± 3.24 days as reported by Nishi (2016) during March-April also confirms the present result. Rehman and Ali (1981) reported a total life span of 24-45 days at 36.7°C . However, a shorter period was recorded by Nayar *et al.* (1976) (20-22 days);

Butani and Jotwani (1984) (22 to 25 days) and Sundraraj and David (1987) (26.9 days). Sharma *et al.* (1985) found that dry and cold weather prolongs the duration of the different stages resulting in prolonged duration of generation, whereas humid and warm weather is considered favorable for the growth and development of this insect as it completes its generation in a shorter time. It could be concluded that temperature influences the development of the moth which is inversely proportional to it (Table 1).

F. Fecundity

The longevity of adult female is an important factor in the realization of its oviposition potential. The fecundity was highest in *post-kharif* 132.45±5.83 eggs/female followed by *kharif* (117.55±10.60 eggs/female) and *pre-kharif* (91.55±10.93 eggs). In support of the present result Das and Chaudhuri (2012) reported that the adult female of the October-November generation laid 135 eggs. However, a higher fecundity of 200-400 eggs/female was recorded by Atwal and Dhaliwal (2005); Panwar (2002) and Srivastava (2003). David (2002) also recorded higher fecundity of 385 eggs/female. According to Bhat *et al.* (2005) highest fecundity was recorded during September (206.40 eggs/female) and the lowest in June (187.4 eggs/female). In support of Bhat *et al.* (2005), Kumar *et al.* (2014) also revealed that the fecundity was 196 eggs during July-August as compared to 203 eggs during October-November. Nishi (2016) found that the fecundity was 199±34.3 eggs in March-April. Each female laid about 277 eggs and 150-250 eggs singly as reported by Syed *et al.* (2011) and Shah *et al.* (2012). Although, the fecundity of fruit borer in aforesaid works is not in conformity with the present investigation trends in seasonal variation are at par with the results present study.

Shoot and Fruit borer life cycles were shorter during *pre-kharif* (27.45 days), indicating that there were more numbers of generations, but because of the low fecundity (91.00 eggs/female), population expansion may have been slower. The Fruit and Shoot Borer took 31.60 days in *kharif*, which is longer than in *pre-kharif*, but its higher fecundity

(117.55 eggs/female) may lead to a large population expansion. The longest life cycle was recorded in *post-kharif* season (44.10 days) but higher fecundity (133.00 days), may lead to higher population growth than in the *pre-kharif* and *kharif* seasons. It can be concluded *post-kharif* seasons is most vulnerable to okra fruit and shoot bore followed by *kharif* and *pre-kharif*.

ACKNOWLEDGEMENT

First Author is grateful to Professor SK Senapati and Dr. Nilanjana Chudhuri for proper guidance during the investigation.

REFERENCES

- Al-Mehmmady Roqaya M. (2000) Biological Studies on the okra moth, *Earias vittella* (F.) (Lepidoptera: Noctuidae) in Jeddah, Saudi Arabia. Research Bulletin No. 96, Research Centre, College of Agriculture, King Saud University. pp5–8.
- Ambegankar, J.K. and Billapate G.G. (1984) Growth development and biometrics of *Earias vittella* (Fab) on cotton and okra. Journal of Maharashtra Agricultural University 9: 254–256.
- Atwal S. and Dhaliwal G.S. (2005) Agricultural pests of South Asia. Kalyan publisher, New Delhi India.
- Bhat O.K., Bhagat K.C., Koul V.K. and Rajinder B. (2005) Studies on the biology of spotted bollworm *Earias vittella* (Fab.) on okra. Annals of Plant Protection Sciences 13(1): 65–68.
- Butani D.K. and Jotwani M.G. (1984) Insect in vegetable. Periodical expert Book Agency. Vivek-Vihar, Delhi, India. pp45–66.
- Das M. and Chaudhuri N. (2012) Life system analysis of *Earias vittella* for formulation of pest model. Research on Crops 13(2): 608–613.
- Dhillon M.K. and Sharma P.D. (2004) Studies on biology and behavior of *Earias vittella* (Lepidoptera: Noctuidae) for mechanisms of resistance in different cotton genotypes. Crop Protection 23(3): 235–241.
- Kumar D., Raghuraman M., Singh R.N., Santeswari and Singh J. (2014) Effect of environmental factors on the population of spider mite, *Tetranychus urticae* Koch on okra in Varanasi region. The Ecoscan 6: 231–235.

- Nayar K.K., Ananthakrishnan T.N. and David B.V. (1976) General and Applied Entomology. Tata Mc Graw Hill Publishing Co. Ltd. New Delhi, India. pp589.
- Nishi S. (2016) Biology of spotted bollworm, *Earias vittella* (Fab.) on okra. Journal of Global Research, Computer Science and Technology 4(2): 62–71.
- Raju A. (2016). Ecology and management of okra shoot and fruit borer (*Earias vittella* Fabricius): M.Sc. thesis submitted to College of Agriculture Rajendranagar, Professor Jayashankar Telangana State Agricultural University, Hyderabad, India.
- Rehman M.H. and Ali H. (1981) Biology of spotted bollworm of cotton, *Earias vittella* (F.). Pakistan Journal of Zoology 13: 105–110.
- Rukhsana G.I., Khan, M.F. and Naqvi H. (1995) Comparative study of *Abelmoschus esculentus* (L.) Moench Meth, okra fruit and cotton leaves suitability for rearing and biology of *Earias fabia* Stoll. Pakistan Journal of Scientific and Industrial Research 38(1): 22–24.
- Shah M.A., Memon N., Manan A., Shah N.A. (2012) Effect of different temperatures on the development of spotted bollworm, *Earias vittella* (Fab.) (Lepidoptera: Noctuidae) in the laboratory. Sindh Universities Research Journal (Sci. Ser.) 44 (3): 487–490.
- Sharma I. N., Lall B.S. Sinha R. P. and Singh B. N. (1985) Biology of spotted bollworm, *Earias vittella* Fab. Journal of Bulletin of Entomology 26 (1): 38–41.
- Singh Y. and Bichoo S.I. (1989) Some biological and bionomical observations on *Earias vittella* Fabia Stoll. Bulletin of Entomology, New Delhi 30(1): 84–91.
- Sundararaj R.T. and David B.V. (1987) Influence of biochemical parameters of different hosts of the biology of *Earias vittella* (Fab.) (Noctuidae: Lepidoptera. Proceeding of Indian Academy Science (Animal Science) 96 (3): 329–332.
- Suryawanshi D.S., Pawar V.M. and Borikar P.S. (2001) Life fecundity tables of *Earias vittella* (Fabricius) on okra and cotton. Journal of Entomology 25(4): 249–262.
- Syed T.S., Abro G.H., Khanum A. and Sattar M. (2011) Effect of host plants on the biology of *Earias Vittella* (Fab) (Noctuidae: Lepidoptera) under laboratory conditions. Pakistan Journal of Zoology 43(1): 127–132.

(Received June 01, 2023; revised ms accepted September 14, 2023; published December 31, 2023)

***Lysinibacillus fusiformis*: A novel mosquitocidal bacterium isolated from Western Ghats, Kerala, India**

K. Aneha, Bhagyashree Bora, P. Hemaladkshmi, S. Manikandan, Jibi Lukose, V. Abhisubesh, Kakhuangailu Gangmei, Sahadiya Mandodan, A. Mathivanan, K. Vijayalakshmi and S. Poopathi*

Unit of Microbiology and Immunology, ICMR- Vector Control Research Centre, Ministry of Health and Family Welfare, Puducherry 605006, India.

Email: subbiahpoopathi@rediffmail.com

ABSTRACT: An extensive field survey was carried out from December 2021 to January 2022 for the collection of soil samples in the Western Ghat region of the Wayanad district of Kerala, India, to isolate potent novel mosquitocidal bacteria which could be used as efficient formulations to control mosquito vectors. Several bacterial colonies were isolated and screened, for mosquitocidal activity. Toxicity assay showed that only one bacterium isolated from forest loamy soil had promising larvicidal activity against *Culex quinquefasciatus* the lymphatic filariasis vector (LC_{50} : 0.03mg L⁻¹ and LC_{90} : 0.07mg L⁻¹) and moderate activity was shown against the dengue vector, *Aedes aegypti* (LC_{50} : 1.03 mg L⁻¹ and LC_{90} : 1.8 mg L⁻¹). The bacterium was identified as *Lysinibacillus fusiformis* by a phylogenetic tree constructed using 16S rRNA genome sequencing. This is the first report that *L. fusiformis* isolated from the forest loamy soil of the Western Ghats, Kerala, which showed proficient mosquitocidal activity against disease-transmitting mosquito vectors. © 2023 Association for Advancement of Entomology

KEYWORDS: *Culex quinquefasciatus*, *Aedes aegypti*, soil, genome sequencing, first report, toxicity assays

Mosquito-borne diseases (MBD), such as dengue, malaria, and Zika virus fever, chikungunya, pose severe risks to the public's health (WHO, 2020). Vector control is crucial for reducing the epidemics of several MBDs, limiting disease transmission, and improving the quality of life. The resurgence of vector-borne diseases as a result of favorable environmental factors, such as rapid urbanization, the development of vector resistance to insecticides, and human lifestyle changes encourage the need for the adoption of safer and more efficient vector control techniques (Chediak *et al.*, 2016; Garcia *et*

al., 2018). The WHO promotes integrated vector management (IVM), which emphasizes the use of long-lasting, eco-friendly alternative vector control measures. Biological control measures, including bacterial pesticides, are found to be a better alternative since they are eco-friendly, cost-effective, and target-specific. To control mosquito vectors, *Bacillus thuringiensis israelensis* (*Bti*) and *B. sphaericus* (*Bs*) bacteria are frequently used (Prummongkol *et al.*, 2019; Vimala Devi *et al.*, 2021). However, the long-term usage of these bio-control agents like *Bs* resulted in the occurrence

* Author for correspondence

of resistance in vector mosquitoes (Guidi *et al.*, 2013; Su *et al.*, 2019). This challenging situation prompted the researchers to look for novel mosquitocidal bacteria from natural sources. The Western Ghats in Kerala are renowned for their diverse and unique collection of fauna and flora. Soil is the richest source of microbes and there have been numerous reports of mosquitocidal bacteria isolated from soil in different parts of the world (Nair *et al.*, 2020; Iftikhar *et al.*, 2023). Whereas the soil of the Western Ghat region of Kerala has not been explored for isolation of mosquitocidal bacteria. Therefore, the present study aimed to isolate novel mosquitocidal bacteria from various soil types collected from the Western Ghats, Wayanad, Kerala, India.

The soil samples were collected from tea plantations, coffee plantations, and forests in the Wayanad district, which includes parts of the Western Ghats (11.68°N; 76.13°E). The soil sampling was done from December 2021 to January 2022. In each sampling spot, the surface layer, 5 and 10 centimeters below the soil, were collected and pooled together in sterile vials. Soil types were documented. Samples were brought into the laboratory and processed by serial dilution and spread plate method. A required amount of soil sample was weighed (1g) and serially diluted (10^{-3}) in 10 ml of sterile water, and 0.1 ml of this serially diluted sample was evenly spread on Nutrient Yeast Salt Mineral (NYSM) agar plates (spread plate method) and these plates were incubated for 24 hours at room temperature. Bacterial colonies were inoculated in 10 ml NYSM broth and incubated in an Orbitek shaker at 250 rpm for 72 hours.

The preliminary toxicity assay (bioassay) was carried out as recommended by WHO guidelines

(WHO, 2005) using 72 hours of bacterial culture to screen for mosquitocidal activity. The bioassays were conducted in paper cups (wax-coated) having 100 ml of tap water (chlorine-free) and 25 late third instar larvae of *Culex quinquefasciatus* and *Aedes aegypti* were released into the cups and acclimatized. A dose of 1 µl of 72 hours of bacterial culture was used to treat the larvae in the cups, while controls without the addition of the bacterial culture were maintained. After 24 hours of exposure, the bioassay results were recorded by scoring the live larvae in respective cups. The bacterial isolate with 100 percent mortality of larvae was considered potential bacteria, and stored in a deep freezer as glycerol stock (30%) until further use. The potential mosquitocidal bacterium was subjected to extensive bioassay /detailed toxicity assay to determine the lethal concentration values (LC_{50} and LC_{90}). Bacterial glycerol stocks were inoculated in 10 ml of NYSM broth, kept for incubation overnight at 250 rpm in an Orbitek shaker, and sub-cultured in 100 ml of NYSM broth for 72 hours. After complete sporulation, the bacterial cell pellets were separated by centrifugation (10000g, 20 min) (Hitachi, Japan) and then lyophilized (Christ ALPHA 1-2 LD plus, Germany). A homogenized bacterial stock solution was prepared (5mg/10 ml) using lyophilized powder. Seven different doses of this stock solution were used for conducting the extensive bioassay. Four replicates for each dose were maintained in every experiment with appropriate controls (WHO, 2005). After 24 hours, the result was scored by counting the live larvae. The experiment was repeated three times to ensure reproducibility. The results were finally analyzed by Probit analysis using SPSS.16.0 software.

Table 1. Mosquitocidal activity of *L. fusiformis*

Mosquito species	LC_{50} (mg l ⁻¹)	LC_{90} (mg l ⁻¹) (LCL*-UCL*)	Slope (LCL*-UCL*)	Intercept	X ²
<i>Culex quinquefasciatus</i>	0.03 (0.025-0.039)	0.07 (0.068-0.081)	0.006	-1.160	89.879
<i>Aedes aegypti</i>	1.02 (0.94-1.1)	1.8 (1.69-1.9)	0.002	-1.714	17.864




Fig. 1 Phylogenetic tree constructed using 16S rRNA genome of the newly isolated mosquitocidal bacterium (Neighbour joining model)




Fig. 2 Colony morphology of *L. fusiformis*




Fig. 3 Vegetative cells of *L. fusiformis* after Gram staining




Fig. 4 Spores of *L. fusiformis* after Safranin-Malachite green endospore staining

The characteristics of bacterial colonies such as the shape, margin, opacity, and color of the bacterial isolate, were observed in the NYSM plate culture. Gram-staining was done to study whether the bacteria were Gram-negative or positive. Spore staining, also known as Schaeffer-Fulton staining, was done to visualize the presence of spores through a microscope (Olympus CX41RF Binocular Microscope, Japan).

For the identification of the mosquitocidal bacterium the genomic DNA was extracted using the GenElute™ Bacterial Genomic DNA Kit (Sigma Aldrich, USA), and polymerase chain reaction (PCR) was done to amplify the genomic DNA with universal forward and reverse primers (16S rRNA). The Qiaquick PCR purification kit (QIAGEN, USA) was employed to purify the amplified products, which then served as a template for forward and reverse sequencing. The BigDye Version 3.1 kit from Applied Biosystems was utilized to do the DNA sequencing on the ABI-PRISM 3730 DNA sequencer and the contigs sequences were assembled by Bio-Edit (Version 7.0.9.0) and examined via Mascot Server 2.4. BLAST program (NCBI), which was used to identify the species. The 16S rRNA genome sequence was combined with the closely related sequence, and a next-generation phylogenetic tree was constructed using

MEGA 5 using the K2P model and 1000 bootstrapping.

The Western Ghats in the states of Kerala were granted a “heritage tag” from UNESCO as they are the “gene pool” that harbors millions of Millions of species. The soil in this region is an excellent source of numerous unique microorganisms that might be investigated for potential applications. There has been limited research on the isolation of mosquitocidal bacteria from the soils of the Western Ghats region of Kerala. (Nampoothiri *et al.*, 2013). In the present study, explorative research was carried out by collecting various soil types from the Western Ghats of Wayanadu District of Kerala for isolating novel mosquitocidal bacteria. A total of 180 soil samples were collected, and several bacteria were isolated. Among these 12 bacterial strains showed mosquitocidal activity and among these only one bacterium isolated from forest loamy soil had promising larvicidal activity against *Culex quinquefasciatus* the lymphatic filariasis vector (LC_{50} : 0.03mg L⁻¹ and LC_{90} : 0.07mg L⁻¹) and moderate activity was shown against the dengue vector, *Aedes aegypti* (LC_{50} : 1.03mg L⁻¹ and LC_{90} : 1.8mg L⁻¹) (Table 2). This potential bacterial strain was identified as *Lysnibacillus fusiformis* by constructing the phylogenetic tree using the 16S rRNA genome sequence (Fig. 1). The colony

morphology of *L. fusiformis* was circular, dry, flat, and rough, with a dull white color (Fig. 2). Microscopic studies on the vegetative stage of the bacterium found it to be rod-shaped and Gram-positive (Fig. 3). The strain was endospore-forming with terminal spherical spores (Fig. 4).

Ramalakshmi and Udayasuriyan (2010) reported different of *Bt* strains from the soil samples of the Western Ghats extension in Tamil Nadu. Similarly, Ganesan and his co-workers (2018) reported *Streptomyces enissocaesilis* (S12-17) from soils of the Western Ghats of Tamil Nadu with effective mosquito larvicidal, ovicidal, and repellent activity (Ganesan *et al.*, 2018). It is the first report of *L. fusiformis* from the soil of Western Ghats, Kerala having mosquitocidal activity. It is a naturally occurring bacterium in the family Bacillaceae and was first discovered in 1901 (Pinheiro *et al.*, 2022). *L. sphaericus* was widely used for mosquito control whereas; the larvicidal activity of *L. fusiformis* has not been reported so far. The underlying mechanism *i.e.*, the toxin (s) responsible for mosquito toxic effect and its mode of action are yet to be studied. In conclusion, this study reports a novel mosquitocidal bacterium, namely, *L. fusiformis* for the first time. It is suggested that the strain may be useful for the control of disease-transmitting mosquito vectors in the current scenario of resistance to some bio-pesticides.

ACKNOWLEDGMENTS

The first author acknowledges the University Grant Commission (UGC), Ministry of Human Resource and Development, Government of India, for the award of a Senior Research Fellowship (Ref: UGC/191620033726). The authors acknowledge the Director, ICMR-Vector Control Research Centre, Puducherry for proving the facilities and RIU approval. This is part of PhD thesis (Ref: PhD/PU/20201129001, 12/04/2021) for the first author under the guidance of the corresponding author (SP).

REFERENCES

- Chediak M., Pimenta F.G.Jr., Coelho G.E., Braga I.A., Lima J.B.P., Cavalcante K.R.L., De Sousa L.C., De Melo Santos M.A.V., Macoris M.L.G., De Araújo A.P., Ayres C.F.J., Andrighetti M.T.M., Gomes R.G.A., Campos K.B. and Guedes R.N.C. (2016) Spatial and temporal country-wide survey of temephos resistance in Brazilian populations of *Aedes aegypti*. *Memórias do Instituto Oswaldo Cruz* 111(5): 311–321.
- Ganesan P., Anand S., Sivanandhan S., David R.H.A., Paulraj M.G., Al-Dhabi N.A. and Ignacimuthu S. (2018) Larvicidal, ovicidal and repellent activities of *Streptomyces enissocaesilis* (S12-17) isolated from Western Ghats of Tamil Nadu, India. *Journal of Entomology and Zoology Studies* 6(2): 1828–1835.
- Garcia G.D. A., David M.R., Martins A.D.J., Maciel-de-Freitas R., Linss J.G.B., Araújo, S.C. and Valle D. (2018) The impact of insecticide applications on the dynamics of resistance: The case of four *Aedes aegypti* populations from different Brazilian regions. *PLoS neglected tropical diseases* 12(2): e0006227.
- Guidi V., Lehner A., Lüthy P. and Tonolla M. (2013) Dynamics of *Bacillus thuringiensis* var. *israelensis* and *Lysinibacillus sphaericus* spores in urban catch basins after simultaneous application against mosquito larvae. *PloS one* 8(2): e55658.
- Iftikhar S.R., Majeed M.Z., Afzal M., Ali A., Saadia M., Ali Z. and Ahmed S. (2023) Isolation, characterization and larvicidal potential of indigenous soil inhabiting bacteria against larvae of southern house mosquito (*Culex quinquefasciatus* Say). *International Journal of Tropical Insect Science* 43:781–791
- Nair K., Al-Thani R., Ginibre C., Chandre F., Alsafran M. and Jaoua S (2020) *Bacillus thuringiensis* strains isolated from Qatari soil, synthesizing ä-endotoxins highly active against the disease vector insect *Aedes aegypti* Bora Bora. *Heliyon* 6(10): e05003.
- Nampoothiri K.M., Ramkumar B. and Pandey A. (2013) Western Ghats of India: Rich source of microbial biodiversity. *Journal of Scientific & Industrial Research* 72: 617–623.
- Pinheiro L.R.S., Gradissimio D.G., Xavier L.P. and Santos A.V. (2022) Degradation of azo dyes: bacterial potential for bioremediation. *Sustainability* 14(3): 1510.
- Prummongkol S., Komalamisra N., Phanphoo Wong T., Chauvatcharin S., Panbangret W. and Srisawat

- R. (2019) Efficacy of liquid bacterial formulation of *Bacillus thuringiensis* subsp. *israelensis* and *Bacillus sphaericus* 1593 mixture in controlling mosquito larvae in Bangkok. *Southeast Asian Journal of Tropical Medicine and Public Health* 50(5): 808–815.
- Ramalakshmi A. and Udayasuriyan V. (2010). Diversity of *Bacillus thuringiensis* Isolated from Western Ghats of Tamil Nadu State, India. *Current Science* 61(1): 13–18.
- Su T., Thieme J., White G. S., Lura T., Mayerle N., Faraji A. and Brown M. Q. (2019) High resistance to *Bacillus sphaericus* and susceptibility to other common pesticides in *Culex pipiens* (Diptera: Culicidae) from Salt Lake City, UT. *Journal of Medical Entomology* 56(2): 506–513.
- Vimala Devi P.S., Duraimurugan P., Chandrika K. S.V.P and Vineela V. (2021) Development of a water dispersible granule (WDG) formulation of *Bacillus thuringiensis* for the management of *Spodoptera litura* (Fab.). *Biocontrol Science and Technology* 31(8): 850–864.
- WHO (2005) Guidelines for laboratory and field testing of mosquito larvicides. World Health Organization, Available from <https://apps.who.int/iris/handle/10665/69101>
- WHO (2020) Vector Borne Diseases. World Health Organization, Available from <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>

(Received June 25, 2023; revised ms accepted September 26, 2023; published December 31, 2023)

New faunistic records of two species of pygmy backswimmers (Hemiptera, Heteroptera, Pleidae) from Kerala, India

K. Jyothylakshmi and S. Nandakumar*

P.G and Research department of Zoology, N.S.S. College, Pandalam 689501, Kerala, India

Email:nandakumar78@gmail.com

ABSTRACT: Members of the family Pleidae are commonly referred to as pygmy backswimmers. They are small predatory bugs; usually reside in stagnant or slow moving aquatic habitats with plenty of vegetation. Two species of aquatic insects *Paraplea frontalis* (Fieber, 1844) and *P. liturata* (Fieber, 1844) in the family Pleidae are reported for the first time from Kerala. Both species are relatively common and widespread in India. However, the present faunistic records shall add to the database of geographical distributional range of these species. © 2023 Association for Advancement of Entomology

KEYWORDS: Aquatic bugs, *Paraplea frontalis*, *P. liturata*, nepomorpha, distribution

Pygmy backswimmers or pleids are small globular bugs generally seen in vegetated areas of lentic water bodies (Andersen and Weir, 2004; Chen *et al.*, 2005). As the name indicates, they swim in the inverted or back position. World fauna of the family Pleidae is represented by 37 species under three genera namely, *Plea* Leach, 1815, *Neoplea* Esaki and China 1928 and *Paraplea* Esaki and China 1928. Morphotaxonomic identification of different genera of the family Pleidae is primarily based on the number of tarsal segments. The genus *Paraplea* has two tarsal segments on their fore tarsi and hind tarsi (Cook, 2017; Cook *et al.*, 2020). *Paraplea* is the biggest and extensively distributed genus among pleids (Schuh and Slater, 1995). It is generally worldwide in distribution, but it has not been reported yet from the Palearctic and Antarctic regions (Cook, 2017). Since information on the systematics and detailed bioecology of aquatic bugs

from the state of Kerala is sparse, there is scope for future research regarding the taxonomy and bionomics of true bugs in the family Pleidae. Therefore, an endeavour was made to study water bugs from Sasthamkotta lake, the largest freshwater lake of Kerala. Two species of pygmy backswimmers *Paraplea frontalis* (Fieber, 1844) and *P. liturata* (Fieber, 1844) are hereby reported for the first time from Kerala.

The examined specimens were collected from Sasthamkotta lake, Kollam District, Kerala, by using a D-frame aquatic net with a mesh size of 500µm. The collected samples were sorted, and then preserved in 70 per cent ethanol. The adult specimens were taken for taxonomic study. Observations and measurements were done using Olympus SZ51 stereomicroscope. Photographs of the specimens were taken using Olympus TG- 6

* Author for correspondence

digital camera and later identified with the help of standard literatures and monograph (Bal and Basu, 2000; Nahar, 2004; Thirumalai, 2004; Nieser *et al.*, 2005; Thirumalai and Suresh Kumar, 2005; Jehamalar and Chandra, 2016; Basu *et al.*, 2018; Mitamura *et al.*, 2018; Cook, 2020).

Order Hemiptera Linnaeus, 1758

Suborder Heteroptera Latreille, 1810

Infraorder Nepomorpha Popov, 1968

Superfamily Pleoidea Fieber, 1851

Family Pleidae Fieber, 1851

Genus *Paraplea* Esaki and China, 1928

1. *Paraplea frontalis* (Fieber, 1844)

1844. *Ploa frontalis* Fieber, *Entomologische Monographien, Leipzig*, 18.

1898. *Plea frontalis* Fieber: Kirkaldy, *Wien. Ent. Zeit.*, 17: 141.

1906. *Plea frontalis* (Fieber): Distant, *Fauna of British India*, 3: 48.

1934. *Plea (Paraplea) frontalis* (Fieber): Lundblad, *Archiv für Hydrobiologie Supplement*, 12:138.

1947. *Plea (Paraplea) frontalis* (Fieber): Hafiz & Pradhan, *Rec. Indian Mus.*, 45: 349

1995. *Paraplea frontalis* (Fieber): Polhemus *et al.*, *Cat. Het. Palaerctic region*, 1: 74.

Materials examined: 2 ♂, 1 ♀, Sasthamkotta Lake; Vettolikadavu, Kollam district, Kerala, 8m amsl., 9°1'56.93" N; 76°37'29.72" E, 18.xii.2022, Coll. K. Jyothylakshmi and S. Nandakumar.

Diagnosis: Body length: 1.8- 2mm; colour: golden brown to dark brown; body prominently punctuate, honeycomb pattern; eyes reddish brown; a small dark reddish-brown longitudinal stripe or light brown bar between the eyes; vertex with two pair of dark blotches or spots, it may be indistinct or absent in some specimens; pronotum wider than long; scutellum with distinct dark punctures; Hemelytra with very fine dark punctures; legs yellowish brown;

thoracic keels differently shaped, with spine like projection; male paramere with tuft of setae; female ovipositor, triangular in shape with numerous spines (Figs. 1A- E).

Distribution: India: Andaman and Nicobar Islands, Andhra Pradesh, Arunachal Pradesh, Bihar, Chandigarh, Chhattisgarh, Karnataka, Madhya Pradesh, Maharashtra, Odisha, Pondicherry, Punjab, Tamil Nadu, Uttar Pradesh and West Bengal.

Bionomics: The species is very specific to stagnant or slow moving water bodies with considerable vegetation. They are often seen at the interface between water and vegetation mats, sometimes found associated with the emergent macrophytes. Colour polymorphism can be seen among different individuals of the species. Different colour morphs of *Paraplea frontalis* have been collected. The colour ranges from light brown to dark brown and yellowish brown to golden yellow. This difference in colour might be attributed by the habitat and microhabitat conditions. The species have agile movement and predators that feed primarily on mosquito larvae and other small invertebrates.

Remarks: This is the first record of *Paraplea frontalis* (Fieber, 1844) from Kerala. *Paraplea frontalis* was first named as *Ploa frontalis* in the description made by Fieber (1844). The species is relatively wide spread.

2. *Paraplea liturata* (Fieber, 1844)

1844. *Ploa frontalis* Fieber, *Entomologische Monographien, Leipzig*, 19.

1904. *Plea frontalis* Kirkaldy, *Wien. Ent. Zeit.* 23: 129

1906. *Plea liturata* (Fieber), Distant, *Fauna British India*, 3: 47.

1910. *Plea metiadura* Distant, *Fauna British India*, 5: 337.

1934. *Paraplea liturata* (Fieber): Lunblad. *Arch. Hydrobiol. Suppl.*, 12: 129.




Fig. 1 A - E. *Paraplea frontalis* (Fieber, 1844) A. Dorsal view; B. Frontal view; C. Ventral view; D. Lateral view; E. Hemelytra

1994. *Paraplea liturata* (Fieber): Agarker *et al.*, *Bionature* 14: 78.

Materials examined: 2 ♂, Sasthamkotta Lake; Punnakkadu Kadavu, Kollam district, Kerala, 8m amsl., 9°2' 41.27"N; 76°37' 44.17" E, 24.ii.2022, Coll. K. Jyothylakshmi and S. Nandakumar.

Diagnosis: Body size: 1.2- 1.6mm; colour: yellowish brown to golden brown; dark bands on hemelytra; eyes red coloured; antenna three-segmented, hidden below; punctures spread all over the body; five characteristic dark black spots on the pronotum; legs with numerous setae and tiny spines; thoracic regions clearly separate, anterior thoracic keel




Fig. 2 A- D. *Paraplea liturata* (Fieber, 1844) A. Dorsal view; B. Head and Pronotum; C. Ventral view; D. Lateral view

rounded; two posterior thoracic segments serrated; abdominal keel with anterior two segments conjoined; male subgenital plate almost pentagon shaped, longer than wide, surface slightly corrugated; female ovipositor rectangle shaped with spines (Figs. 2A- D).

Distribution: India: Bihar, Delhi, Madhya Pradesh, Tripura and West Bengal

Bionomics: They were found crawling on the lake vegetation. Like other pleids, they prey on mosquito larvae and small invertebrates. The area in which the species was collected had a substantial amount of aquatic vegetation. The species is usually overlooked and rarely collected due to its minute size and swift movement. More comprehensive observations are needed to reveal their bioecology in detail, since only limited specimens have been

obtained during the present investigation.

Remarks: This is the first record of *Paraplea liturata* (Fieber, 1844) from Kerala. It can be easily separated from the closely resembling species, *Paraplea lateromaculata* Cook, 2020 by the presence of five distinct dark spots on pronotum.

A total of four species under the genus *Paraplea* such as *Paraplea buenoi* Kirkaldy, 1904, *Paraplea frontalis* (Fieber, 1844), *Paraplea indistinguenda* (Matsumura, 1905) and *Paraplea liturata* (Fieber, 1844) have been so far reported from India (Thirumalai, 2007). *Paraplea indistinguenda* (Matsumura, 1905) has been the only known species of *Paraplea* reported from Kerala so far (Thirumalai, 2007). The current record of *Paraplea frontalis* (Fieber, 1844) and *Paraplea liturata* (Fieber, 1844) is pivotal, since it is the pioneer report from Kerala and the records of the same from a wetland of international significance. The present faunistic records of the pygmy backswimmers are from Sasthamkotta lake, the largest freshwater lake of Kerala, and one among the top Ramsar sites of India. Macrophytes influence the diversity, abundance and distribution pattern of pleids by providing diverse ecological niches. Sasthamkotta Lake is infested with several native and non native macrophyte species. Therefore, it is imperative to conduct ample research activities on the taxonomy and faunistic of pygmy backswimmers of the lake, so that knowledge regarding this group can be utilized as baseline data for further research and conservation planning. *P. frontalis* (Fieber, 1844) was earlier reported from other states of India by Thirumalai (2004), Nieser *et al.* (2005), Thirumalai and Suresh Kumar (2005), Jehamalar and Chandra (2016) and Basu *et al.* (2018). Previous reports of *Paraplea liturata* (Fieber, 1844) were also made by Bal and Basu (2000) and Nahar (2004). Hence, keeping in view the scarcity of knowledge regarding the pygmy backswimmers of the State, intensive surveys in the unexplored ecosystems of Kerala are required. The present work could direct attention towards the necessity of faunistic study of pygmy backswimmers from other wetlands of Kerala to reinforce knowledge on its distribution.

ACKNOWLEDGEMENTS

The first author thanks University of Kerala for the Junior Research Fellowship and grateful to The Principal, N.S.S College, Pandalam for providing facility in the PG & Research Department of Zoology to complete the work. Authors are thankful to Dr. E. Eyarin Jehamalar, Zoological survey of India, Kolkata, for constant encouragement. Authorse are beholden to Dr. A. Biju Kumar, Professor and Head, Department of Aquatic biology and Fisheries, University of Kerala, Kariavattom, for the facilities and encouragement.

REFERENCES

- Andersen N.M. and Weir T.A. (2004) Australian water bugs, their biology and identification (He-miptera-Heteroptera, Gerromorpha & Nepomorpha): 21. Family Pleidae, pygmy back-swimmers. Entomonograph 14: 304–308.
- Bal A. and Basu R.C. (2000) Insecta: Hemiptera: Water Bugs. Zoological Survey of India, Fauna of Tripura, State Fauna Series 7 (Pal 1-2): 427–443.
- Basu S., Chandra K., Subramanian K.A. and Saha G.K. (2018) Water bugs (Insecta: Hemiptera: Heteroptera) of Himalayan and sub-Himalayan regions of West Bengal, India. Journal of Threatened Taxa 10 (12): 12619–12714.
- Chen P.P., Nieser N. and Zettel H. (2005) The Aquatic and Semi-aquatic Bugs (Heteroptera: Nepomorpha & Gerromorpha) of Malesia. Brill, Leiden, The Netherlands. 546pp.
- Cook J.L. (2017) A Second Species of Heteroplea Cook (Hemiptera: Pleidae). Proceedings of the Entomological Society of Washington 119(4): 545–550. doi: 10.4289/0013-8797.119.4.545
- Cook J.L., Sites R.W. and Vitheepradit A. (2020) The Pleidae (Hemiptera, Heteroptera) of Thailand, with the descriptions of two new species and a discussion of species from Southeast Asia. ZooKeys 973: 35.
- Jehamalar E.E. and Chandra K. (2016) Additional records of aquatic and semi-aquatic Heteroptera (Insecta: Hemiptera) from Chhattisgarh, India. Records of the Zoological Survey of India 116 (2): 99–109.
- Lundblad O. (1934) Zur kenntnis der aquatilen und semiaquatilen Hemipteren von Sumatra, Java und

- Bali. Archiv für Hydrobiologie - Supplement 12: 1–95, 263–489.
- Mitamura T., Liu H-C., Hirasawa K. and Yoshii S. (2018) New distributional record of *Paraplea liturata* (Fieber, 1844) (Hemiptera: Pleidae) from Taiwan. Taiwanese Journal of Entomological Studies 3: 72–74.
- Nahar S.C. (2004) Aquatic Insects. Zoological Survey of India. State Fauna series, 11 : Fauna of Bihar (Including 1 Iza rkhand) Part 1 : 195–213.
- Nieser N., Chen P.P. and Chang Man Yang (2005) A new subgenus and six new species of Nepomorpha (Insecta: Heteroptera) from Yunnan, China. Raffles Bulletin of Zoology 53 (2): 189–209.
- Schuh R.T. and Slater J.A. (1995) True Bugs of the World. Cornell University Press, Ithaca, New York. 336pp.
- Thirumalai G. (2004) A checklist of aquatic and semi-aquatic Hemiptera (Insecta) of Karnataka. Records of Zoological Survey of India 102 (1-2): 57–72.
- Thirumalai G. (2007) A synoptic list of Nepomorpha (Hemiptera: Heteroptera) from India. Records of the Zoological Survey of India Occasional Papers 273: 1–84.
- Thirumalai G. and Suresh Kumar R. (2005) Aquatic and semi-aquatic Hemiptera (Heteroptera) of Karaikal and Pondicherry. Records of Zoological Survey of India 105(1-2): 5–24.

(Received June 29, 2023; revised ms accepted November 11, 2023; published December 31, 2023)

Calamus tenuis Roxb. a new host for *Chilo suppressalis* (Walker) (Lepidoptera, Crambidae)

Selim Mehmud¹, Sasanka Sekhar Ghosh², Himu Roy³, Dip Jyoti Haloi^{4*} and
Pranab Bujarbarua¹

¹Department of Botany, Handique Girls' College, Panbazar, Guwahati 781001, Assam, India.

²Department of Zoology, Cotton University, Panbazar, Guwahati 781001, Assam, India.

³Department of Botany, Cotton University, Panbazar, Guwahati 781001, Assam, India.

⁴Department of Zoology, Handique Girls' College, Panbazar, Guwahati 781001, Assam, India.

Email: diphalo1979@gmail.com

ABSTRACT: *Calamus tenuis* Roxb. belonging to Arecaceae was recorded as a new larval host for *Chilo suppressalis* (Lepidoptera, Crambidae) from Kamrup district of Assam, India. However, other parts like leaf, stem, root and pistillate inflorescence were found unaffected.

© 2023 Association for Advancement of Entomology

KEY WORDS: Arecaceae, rattan, inflorescence, borer, Assam

Rattans are climbing palms of the monocot family Arecaceae (Renuka *et al.*, 2010). The association between different insects and rattans has been reported by many workers in various countries (Chung, 1995; Mattes *et al.*, 1998; Howard *et al.*, 2001; Sunderland, 2004; Merklinger *et al.*, 2014; Shameem and Prathapan, 2014; Liu *et al.*, 2019). However, not much information is available on the insect pests of rattans in India (Renuka *et al.*, 2010). *Calamus tenuis* Roxb. (Fig. 1 A) is one of the common and economically important rattan species of Assam. The staminate inflorescence of this dioecious species is around 3m long with 6-9 partial inflorescence (Mehmud and Roy, 2021). After anthesis (Fig. 1 B), the staminate inflorescences dry up and their remains were found to adhere to the plant for several months. During observations of the staminate flowers, larvae

measuring 2-2.5cm (Fig. 1 C-E), were noticed on the rachis, which were reared (Fig. 1, F-I) and got identified as *Chilo suppressalis* (Walker) belonging to Lepidoptera, Crambidae. However, other parts like leaf, stem, root and pistillate inflorescence were found unaffected.

Khan *et al.* (1991) reported that *C. suppressalis* as an important rice borer in India, Indonesia and East Asia and there are around 41 host plants belonged to 30 genera under six families (four monocot and two dicot), where most of the plants were of the grass family Poaceae. Adult *C. suppressalis* measuring 1.3cm, is pale brown. Although *C. suppressalis* affects every part of the rice plant (Lu *et al.*, 2017; Meng *et al.*, 2019), it is worth mentioning that in rattans, only the staminate inflorescence of *C. tenuis* is infested. Voucher

* Author for correspondence




Fig. 1 *Calamus tenuis*: A. Habit (staminate inflorescence), B. Anthesis of staminate inflorescence. *Chilo suppressalis*: C. Larva inside rachis; D-E. Larva inside partial staminate inflorescence; F. Pupa; G-I. Adult.

specimens were submitted in the Department of Botany and *C. suppressalis* in the Department of Zoology, Handique Girls' College respectively.

ACKNOWLEDGEMENTS

The authors are thankful to the Heads of Department of both Botany and Department of Zoology of Handique Girls' College and Cotton University.

REFERENCES

- Chung A.Y.C. (1995) Insect pests on rattans in Sabal. The Planter, Kuala Lumpur 75: 55–66.
- Howard F.W., Moore D., Giblin-Davis R.M. and Abad R.G. (2001) Insects on palms. CABI Publishing. 400pp.
- Khan Z.R., Litsinger J.A., Barrion A.T., Villanueva F.F.D., Fernandez N.J. and Taylo L.D. (1991) World bibliography of rice stem borers, 1794-1990. International Rice Research Institute, Manila. 415pp.
- Liu K., Mansor A., Nadine R., Lee C.Y., Azman N.M. and Fadzly N. (2019). Rattan litter-collecting structures attract nest-building and defending ants. Plant Signaling & Behavior e1621245-1-8.
- Lu Y.H., Wang G.R., Zhong L.Q., Zhang F.C., Bai Q., Zheng X.S. and Lu Z. (2017) Resistance monitor of *Chilo suppressalis* (Walker) (Lepidoptera: Crambidae) to chlorontranilprole in eight field populations from east and central China. Crop Protection 100: 196–202. doi:10.1016/j.cropro.2017.07.006
- Mattes M., Moog J., Werner M., Fiala B., Naisa J. and Maschwitz U. (1998) The rattan palm *Korthalsia robusta* Bl. and its ant and aphid partners: studies of a myrmecophytic association in the Kinabala Park. Sabah Parks Nature Journal 1: 47–60.
- Mehmud S. and Roy H. (2021) Diversity and distribution of palms (Arecaceae) in Assam, India. Check List 17(1): 69–93. doi:10.15560/17.1.69.
- Meng X.K., Dong F., Qian K., Miao L.J., Yang X.M., Ge H.C., Wu Z. and Wang J. (2019) Transcriptome analysis reveals global gene expression changes of *Chilo suppressalis* in response to sublethal dose of chlorontranilprole. Chemosphere 234: 648–657. doi:10.1016/j.chemosphere.2019.06.129.
- Merklinger F.F., Baker W.J. and Rudall P.J. (2014) Comparative development of the rattan ocrea, a structural innovation that facilitates plant mutualism. Plant Systematics and Evolution 300: 1973–1983.
- Renuka C., Bhat K.V. and Pandalai R.C. (2010) Rattans of India- Taxonomy, Biology and Utilization. Kerala Forest Research Institute, Thrissur, Kerala, India. 339pp.
- Shameem K.M. and Prathapan K.D. (2014) Rattan cane *Calamus travancoricus*- a new host plant record for spindle bug of Arecanut palm, *Mircarvalhoia arecae* (Miller & China). Indian Journal of Entomology 76(3): 252–253.
- Sunderland T.C.H. (2004) Ant and rattan associations in forest of tropical Africa. Ghana Journal of Forestry 15 & 16: 13–19.

(Received June 29, 2023; revised ms accepted September 30, 2023; published December 31, 2023)

Contents of Volume 48

No. 1

- <https://doi.org/10.33307/entomon.v48i1.838> 1
 A new subspecies of *Caltores* Swinhoe, 1893 (Lepidoptera, HesperIIDae)
 from the Malabar Coast, Kerala, India
Kalesh Sadasivan, Dipendra Nath Basu and Krushnamegh Kunte
- <https://doi.org/10.33307/entomon.v48i1.839> 11
 Genetic diversity of cotton *Pectinophora gossypiella* (Saunders)
 (Lepidoptera, Gelechiidae) population inferred from mitochondrial *COI*
 sequences
*I. Padma-Shree, M. Muthuswami, N. Manikanda Boopathi,
 K. Senguttuvan and S. Rajeswari*
- <https://doi.org/10.33307/entomon.v48i1.840> 19
 Dissipation kinetics, effect of household processing, and dietary risk
 assessment of the insecticide chlorantraniliprole residue in bitter gourd
 and soil
*M.M. Mawtham, Bhuvaneswari Kaithamalai, Suganthi Angappan
 and S. Kulanthaisami*
- <https://doi.org/10.33307/entomon.v48i1.841> 33
 Evaluation of the residual activity of newly developed aqueous
 formulation of novel biopesticide *Bacillus cereus* VCRC-641 through
 simulated field trial
*S. Manikandan, V. Abhisubesh, Kakhuangailiu Gangmei, Jibi
 Lukose, K. Aneha, Sahadiya Mandodan, P. Hemaladkshmi,
 Bhagyashree Bora, A. Mathivanan, K. Vijayalakshmi and
 S. Poopathi*
- <https://doi.org/10.33307/entomon.v48i1.842> 39
 Antixenosis of Thodan (AAB), a resistant *Musa* cultivar to *Odoiporus*
longicollis (Olivier) and characterization of larvicide molecules
*T. Ajitha, R. Pradeepkumar, R.V. Ayana Gayathri, J. Anil John,
 V.R. Reshma and D.A. Evans*
- <https://doi.org/10.33307/entomon.v48i1.843> 51
 A checklist of Indian Epipaschiinae (Lepidoptera, Pyraloidea, Pyralidae)
*Venkatesh Tanagal, M. Shankara Murthy and
 P. Maheswara Reddy*

- <https://doi.org/10.33307/entomon.v48i1.844> 63
Morphological and biochemical basis of resistance against the pod borers *Maruca vitrata* F. and *Helicoverpa armigera* (Hübner) in cowpea
D.V. Muchhadiya, J.J. Patel, N.H. Garaniya and D.R. Patel
- <https://doi.org/10.33307/entomon.v48i1.845> 69
Temperature dependent development and population performance of *Tetranychus fijiensis* Hirst (Acari, Tetranychidae) on Papaya
Shivakumar Veerlapati and N. Srinivasa
- Short Communication**
- <https://doi.org/10.33307/entomon.v48i1.846> 77
First record of three exotic whitefly pests (Hemiptera, Aleyrodidae) from Andaman and Nicobar Islands, India
Anil Kumar Dubey
- <https://doi.org/10.33307/entomon.v48i1.847> 83
Studies on external genitalic attributes of two species of genus *Traminda* Saalmüller (Lepidoptera, Geometridae, Sterrhinae) from India
Tarun Goyal, Jagbir Singh Kirti and S. Staklin Stella
- <https://doi.org/10.33307/entomon.v48i1.848> 89
Coptosoma variegatum (Herrich-Schäffer, 1838) (Hemiptera, Plataspidae) infesting mango in Kerala and Karnataka with redescription of the species
Jyothi Sara Jacob, S. Salini and S. Shanasa
- <https://doi.org/10.33307/entomon.v48i1.849> 95
Record of *Onomarchus uninotatus* (Serville, 1838) (Orthoptera, Tettigoniidae) as a pest of jackfruit from Kerala, India
P. Sreeja, Neena Lenin, Haseena Bhaskar, Mani Chellappan and Dhaneesh Bhaskar
- <https://doi.org/10.33307/entomon.v48i1.850> 99
Four new records of dragonflies (Insecta, Odonata) from Amboli region of Western Ghats, Maharashtra, India
Dattaprasad Sawant, Rakesh M. Deulkar, Srujan Ogale and Hemant Ogale

	621
https://doi.org/10.33307/entomon.v48i1.851	107
Salad cucumber, <i>Cucumis sativus</i> L.: A new host record for <i>Apomecyna saltator</i> (Fab.) (Coleoptera, Cerambycidae) <i>P. Sreeja and Sainamol P. Kurian</i>	
https://doi.org/10.33307/entomon.v48i1.852	111
Isolation and characterization of sex specific cuticular hydrocarbons in non- <i>Drosophila</i> , <i>Phorticella striata</i> (Sajjan and Krishnamurthy, 1975) (Diptera, Drosophilidae) <i>B.V. Mahendra, M. Prathibha and R.S. Kavitha</i>	
https://doi.org/10.33307/entomon.v48i1.853	117
First record of vesicant beetles: <i>Paederus nigricornis</i> Bernhauer, 1911 from south India; <i>P. extraneus</i> Wiedemann, 1823 and <i>P. alternans</i> Walker, 1858 (Staphylinidae, Paederinae) from Kerala <i>P.M. Kavyamol, P.J. Vineesh and V.P. Vineetha</i>	
https://doi.org/10.33307/entomon.v48i1.854	123
Oviposition behaviour of <i>Callosobruchus chinensis</i> (L.) (Coleoptera, Chrysomelidae, Bruchinae) on horse gram seed varieties <i>Bhramar Bhattacharyya and Anandamay Barik</i>	
https://doi.org/10.33307/entomon.v48i1.855	129
Trapping analysis of ants (Hymenoptera, Formicidae) in Kurunthamalai, Coimbatore, Tamil Nadu, India <i>J. Sornapriya and M. Lekeshmenaswamy</i>	
https://doi.org/10.33307/entomon.v48i1.856	135
First record of <i>Ahmaditermes emersoni</i> Maiti (Isoptera, Termitidae, Nasutitermitinae) from Meghalaya, India <i>Khirod Sankar Das and Sudipta Choudhury</i>	
https://doi.org/10.33307/entomon.v48i1.857	141
First record of <i>Microcerotermes lahorensis</i> Akhtar (Blattodea, Isoptera, Termitidae) from India <i>Edwin Joseph, B. Albright, Amala Anna Francis, Namitha Jayan and Jobin Mathew</i>	
Book Review	145
Susthira theneechakrishi (Sustainable Bee Keeping) <i>Dr. G. Madhavan Nair</i>	

No. 2

- <https://doi.org/10.33307/entomon.v48i2.886> 149
Description of two new species of *Lisotrigona*
(Hymenoptera, Apidae, Meliponini) from Central India and their nests
Shashidhar Viraktamath, Jagruti Roy, Ashish Kumar Jha and Shubham Rao
- <https://doi.org/10.33307/entomon.v48i2.887> 167
A new species of *Purana* Distant, 1905 (Hemiptera, Cicadidae),
from the Western Ghats, with comments on the erroneous records of
Purana tigrina (Walker, 1850) in south India
Kalesh Sadasivan, Jebine Jose, Bernad M Thampan, P.V. Muralimohan, Baiju Kochunarayanan, Anzil Shereef and Mick Webb
- <https://doi.org/10.33307/entomon.v48i2.888> 185
Efficacy of compounds used in mosquito repellents (DEET, picaridin,
prallethrin and IR3535) against odorant binding protein (OBP20) of
Anopheles gambiae: A molecular docking study
Arunima Choudhury, Innifa Hasan, Pobi Gogoi and Dip Jyoti Haloi
- <https://doi.org/10.33307/entomon.v48i2.889> 195
Species list and pictorial key to the dung beetles
(Coleoptera, Scarabaeidae, Scarabaeinae) in the Chinnar Wildlife Sanctuary
in the south Western Ghats, India
K.A. Sobhana and Thomas K. Sabu
- <https://doi.org/10.33307/entomon.v48i2.890> 209
Compatibility of carbosulfan 25 EC with certain agrochemicals in
brinjal ecosystem
I. Merlin Kamala and S. Chandrasekaran
- <https://doi.org/10.33307/entomon.v48i2.891> 219
Studies on influence of various stages of mulberry leaf in the growth and
cocoonic parameters of silkworm *Bombyx mori* (L.)
J. Nihal Nila and R.D. Stevens Jones
- <https://doi.org/10.33307/entomon.v48i2.892> 229
Age-specific ecological life table of *Spodoptera litura* (F.)
(Lepidoptera, Noctuidae) on groundnut
Sunil Kumaraswamy, Ashok Karuppannasamy, B. Prajna Devaiah, Pramod Katti and Sushila Nadagouda

[https://doi.org/ 10.33307/entomon.v48i2.893](https://doi.org/10.33307/entomon.v48i2.893)

241

Trap crop selection and economic threshold based ecological management of *Spilarctia obliqua* (Walker) (Lepidoptera, Erebidae) for sesame

Souren Dutta, Neha Chowdhury, Bratatee Sardar and Nayan Roy

[https://doi.org/ 10.33307/entomon.v48i2.894](https://doi.org/10.33307/entomon.v48i2.894)

253

A review of *Macromia* Rambur, 1842 (Odonata, Macromiidae) of Western Ghats, with taxonomic notes on *Macromia miniata* Fraser, 1924 and *M. irata* Fraser, 1924

Kalesh Sadasivan, Vinayan P. Nair and K. Abraham Samuel

Short Communication

[https://doi.org/ 10.33307/entomon.v48i2.895](https://doi.org/10.33307/entomon.v48i2.895)

287

Courtship and mating behaviour of pupal parasitoid, *Xanthopimpla flavolineata* Cameron (Hymenoptera, Ichneumonidae, Pimplinae)

C. Kathirvelu, S. Manickavasagam and L. Gopianand

[https://doi.org/ 10.33307/entomon.v48i2.896](https://doi.org/10.33307/entomon.v48i2.896)

293

Termite sampling methods: A comparative study in four habitats of north Kerala

A.V. Anushya, P.R. Swaran and Anjusha Bhasker

[https://doi.org/ 10.33307/entomon.v48i2.897](https://doi.org/10.33307/entomon.v48i2.897)

299

New distributional records of egg parasitoids (Hymenoptera, Chalcidoidea, Trichogrammatidae) from Chhattisgarh, India

Mohsin Ikram, Manendra Kaneria, Mohd Faisal and Waseem Ahmad

[https://doi.org/ 10.33307/entomon.v48i2.898](https://doi.org/10.33307/entomon.v48i2.898)

303

Report of *Nonartha birmanicum* (Jacoby) (Coleoptera, Chrysomelidae) on mango inflorescence

J. Nayanathara and R. Narayana

[https://doi.org/ 10.33307/entomon.v48i2.899](https://doi.org/10.33307/entomon.v48i2.899)

305

Evergestis forficalis (L.) (Lepidoptera, Crambidae), a pest of cruciferous crops in the UT of Jammu and Kashmir, India

Deen Mohd Bhat

[https://doi.org/ 10.33307/entomon.v48i2.900](https://doi.org/10.33307/entomon.v48i2.900)

309

Evaluation of cauliflower genotypes and eco-friendly molecules for management of Diamond back moth, *Plutella xylostella* (Linnaeus) (Lepidoptera, Yponomeutidae) and their safety to natural enemies

L.S. Bhushan and Jayakumar Pathma

No. 3

- <https://doi.org/10.33307/entomon/v48i3/934>** 317
 First record of a rare masked bee *Hylaeus (Indialaeus) strenuus* (Cameron, 1897) from south India with an updated checklist of *Hylaeus* species (Hymenoptera, Colletidae)
Arati Pannure, K.L. Manjunatha, Vidya Mulimani and Vasuki V. Belavadi
- <https://doi.org/10.33307/entomon/v48i3/935>** 323
 Indian record of the old world psyllid *Heterotrioza chenopodii* (Reuter) (Hemiptera, Psylloidea, Triozidae) on quinoa *Chenopodium quinoa* (Amaranthaceae)
Aishwarya Palanisamy, Murugan Marimuthu, Chitra Narayanasamy and Balasubramani Venkatasamy
- <https://doi.org/10.33307/entomon/v48i3/936>** 331
 Grasshoppers, crickets and katydids of Kerala, an updated checklist for the order Orthoptera
E.S. Thasnim, C. Bijoy and Dhaneesh Bhaskar
- <https://doi.org/10.33307/entomon/v48i3/937>** 357
 Growth dilution and its effect on pesticide dynamics in okra
G.T. Pradeepkumar, V. Vijayasree and K.P. Subhash Chandran
- <https://doi.org/10.33307/entomon/v48i3/938>** 365
 Studies on biology and preference of *Helicoverpa armigera* (Hubner) on different hosts and evaluation of botanicals for its management
B. Deepti and J. Pathma
- <https://doi.org/10.33307/entomon/v48i3/939>** 371
 Randomized detection of *kdr* allele frequencies in wild populations of *Aedes aegypti* (Diptera, Culicidae) in Colombo District, Sri Lanka
Chimasha Induni, B.A.N Mendis, H.S.D. Fernando and B.G.D. Nissanka K. De Silva
- <https://doi.org/10.33307/entomon/v48i3/940>** 383
 Effect of heat shock on embryonic development and its impact on commercial traits of silkworm *Bombyx mori* L.
Prashant N. Bavachikar, Hothur Lakshmi, P.J. Raju, Gianluca Tettamanti, Amr Mohamed and Anitha Mamillapalli

<https://doi.org/10.33307/entomon/v48i3/941>

397

Diversity and distribution of true flies (Diptera) of
Kuldiha Wildlife Sanctuary, Odisha, India:
Functional roles based on ecological guilds

*Sankarsan Roy, Santanu Ghara, Santu Paria,
Prakash Chandra Mardaraj and
Susanta Kumar Chakraborty*

<https://doi.org/10.33307/entomon/v48i3/942>

411

A checklist of robber flies (Diptera, Asilidae) of Kerala, India

*Kavya G. Pillai, C. Bijoy, Chris M. Cohen and
Gigi Poulouse*

Short Commuication

<https://doi.org/10.33307/entomon/v48i3/943>

421

Seasonal population dynamics of *Aceria erineus* (Nalepa)
(Acari, Eriophyidae) on walnut trees in Kashmir, India

Sheema Tabasum and A.A. Buhroo

<https://doi.org/10.33307/entomon/v48i3/944>

427

Effect of sowing dates and cultivars on the incidence of *Spodoptera*
frugiperda (J.E. Smith) on maize (*Zea mays* L.)
in Nagaland, India

B. Longkumer, Pankaj Neog, Waluniba and H.S. Devi

<https://doi.org/10.33307/entomon/v48i3/945>

433

A new species of *Chalcis* Fabricius (Hymenoptera, Chalcididae)
from south India

A.P. Ranjith and Dharma Rajan Priyadarsanan

<https://doi.org/10.33307/entomon/v48i3/946>

439

Invitro screening of leaf extracts of selected plants from Lamiaceae,
Asteraceae and Fabaceae for mortality and repellency to *Odoiporus*
longicollis (Olivier)

J.B. Hyzil, U. Gayathri Elayidam and K.V. Baiju

<https://doi.org/10.33307/entomon/v48i3/947>

445

Population dynamics of southern birdwing
(*Troides minos* Cramer, 1779) in Sirumalai Reserve Forest, Eastern
Ghats, India

*B. Archana, E. Joy Sharmila, K. Rangesh, V. Pandi,
S. Susaritha and S. Kalaiselvi¹*

- <https://doi.org/10.33307/entomon/v48i3/948> 449
 A check list of blow fly fauna (Diptera, Calliphoridae) of Kerala including
 forensically significant species
M.P. Rejith Paul and C.F. Binoy
- <https://doi.org/10.33307/entomon/v48i3/949> 455
 Acarine species associated with subterranean termites (Blattodea,
 Termitidae)
Anjitha K Bhadrar and N. Ramani
- <https://doi.org/10.33307/entomon/v48i3/950> 461
 New distributional records of six Aphodiinae species (Coleoptera,
 Scarabaeidae) from south India
*K.A. Sobhana, P. Girish Kumar, Sabu K. Thomas and
 V.D. Hegde*
- <https://doi.org/10.33307/entomon/v48i3/951> 467
 Distributional records of *Onthophagus germanus* Gillet, 1927 and *O.
 orissanus* Arrow, 1931 (Coleoptera, Scarabaeidae, Scarabaeinae) from south
 India
K.A. Sobhana, T.K. Viswanath and V.D. Hegde
- <https://doi.org/10.33307/entomon/v48i3/952> 471
 Diversity of edible insects in Tuensang District,
 Nagaland, India
*Chaueichongla Phom and Jeyaparvarthi
 Somasundaram*
- <https://doi.org/10.33307/entomon/v48i3/953> 475
 Impact of seed dressing insecticides on natural enemies of
Bt cotton ecosystem
P.R. Parmar, G.R. Bhandari, R.D. Patel and H.R. Desai
- Obituary - Professor Rajan Asari 481

No. 4

- <https://doi.org/10.33307/entomon/v48i4/982> 485
 Bees of the genera *Hoplonomia* and *Crocisaspidia* (Halictidae, Nomiinae) of India, with key to species
K.T. Manjusha and T. Jobiraj
- <https://doi.org/10.33307/entomon/v48i4/983> 497
 Taxonomical studies of dragonfly nymphs (Odonata, Libellulidae) using their exuviae
Shaun Paul Adambukulam and Francy K. Kakkassery
- <https://doi.org/10.33307/entomon/v48i4/984> 511
 Screening of wild *Ipomoea* genotypes for resistance against sweet potato weevil *Cylas formicarius* F. based on multiple choice bioassay and phytochemical constituents
B.G. Sangeetha, C.A. Jayaprakas, T. Makesh kumar, T. Santhosh kumar, Shirley Raichal Anil and L.S. Rajeshwari
- <https://doi.org/10.33307/entomon/v48i4/986> 519
 Synthesis and characterization of silver nanoparticles using *Datura metel* L. (Solanaceae) leaf extract and its larvicidal activity on *Epilachna vigintioctopunctata* F.
C. Sundareswari and D.N.P. Sudarmani
- <https://doi.org/10.33307/entomon/v48i4/987> 525
 Redescription of *Carbula indica* (Westwood, 1837) (Hemiptera, Heteroptera, Pentatomidae) from West Bengal, India with a key to the Indian species of the genus
Amartya Pal, Swetapadma Dash, Devanshu Gupta and P.C. Saha
- <https://doi.org/10.33307/entomon/v48i4/988> 531
 Evaluation of the insecticidal action of polyphenolic compounds from *Streblus asper* (Lour.) on the red cotton bug, *Dysdercus cingulatus* (Fab.) (Hemiptera, Pyrrhocoridae)
L. Anila and M.S. Hashim

- <https://doi.org/10.33307/entomon/v48i4/989> 537
 Bioefficacy of *Tagetes minuta* L. against
Aphis craccivora Koch (Hemiptera, Aphididae)
V.V. Ashish, Haseena Bhaskar, U. Sreelatha
and K.B. Deepthy

- <https://doi.org/10.33307/entomon/v48i4/990> 545
 Exploration of natural enemy fauna of aphids and associated ant
 species from eastern dry zone of Karnataka, India
V.S. Chinnu, Vidya Mulimani, S. Shanas
and N. Sumithramma

- <https://doi.org/10.33307/entomon/v48i4/991> 553
 A new species of *Cigaritis* Donzel, 1847
 (Lycanidae, Aphnaeinae) from the southern Western Ghats of
 Peninsular India
S. Ramasamy Kamaya Naicker, Sujitha Prabhakaran
Chandrika, Baiju Kochunarayanan, Jebine Jose,
Manoj Kripakaran, Rajkumar Chidambaram Palaniappan,
Vinayan Padmanabhan Nair and Kalesh Sadasivan

- <https://doi.org/10.33307/entomon/v48i4/992> 567
 Redescription of *Cyphochilus niveosquamosus* (Blanchard, 1851)
 (Coleoptera, Scarabaeidae, Melolonthinae) with notes on two
 species of the genus from India
K.V. Prakash and H.M. Yeshwanth

- <https://doi.org/10.33307/entomon/v48i4/993> 575
 Evaluation of pongamia oil soap against red spider mites,
Tetranychus urticae Koch and its effect on mite predators in
 brinjal ecosystem
P. Jeevitha, K.M. Sreekumar and Mukul Kumar Deka

Short Communication

- <https://doi.org/10.33307/entomon/v48i4/994> 583
 Eri silkworm pupae as fish meal replacement in common carp
 (*Cyprinus carpio* L.) feed in indoor aquaria
R.K. Gokulakrishnaa and Selvamuthukumaran
Thirunavukkarasu

- 629
- <https://doi.org/10.33307/entomon/v48i4/995> 591
- Record of *Anoplocnemis phasianus* (Fabricius, 1781)
(Hemiptera, Heteroptera, Coreidae) from Goa, India
*Vijaykumar S. Gadekar, Ayesha E. Shetkar and
Aishwarya S. Naik*
- <https://doi.org/10.33307/entomon/v48i4/996> 595
- Efficacy of insecticides against cashew leaf miner,
Acrocercops syngramma Meyrick (Lepidoptera, Gracillariidae)
*O. Chethana, Jayalaxmi Narayan Hegde,
K. Rajashekharappa, T. Basavaraj Naik and B.V. Champa*
- <https://doi.org/10.33307/entomon/v48i4/997> 599
- Biology of shoot and fruit borer, *Earias vittella* Fabricius
(Noctuidae, Lepidoptera) on okra
Bikash Subba, Nilanjana Chudhuri and S.K. Senapati
- <https://doi.org/10.33307/entomon/v48i4/998> 605
- Lysinibacillus fusiformis*: A novel mosquitocidal bacterium isolated
from Western Ghats, Kerala, India
*K. Aneha, Bhagyashree Bora, P. Hemaladkshmi,
S. Manikandan, Jibi Lukose, V. Abhisubesh,
Kakhuangailiu Gangmei, Sahadiya Mandodan,
A. Mathivanan, K. Vijayalakshmi and S. Poopathi*
- <https://doi.org/10.33307/entomon/v48i4/999> 611
- New faunistic records of two species of pygmy backswimmers
(Hemiptera, Heteroptera, Pleidae) from Kerala, India
K. Jyothylakshmi and S. Nandakumar
- <https://doi.org/10.33307/entomon/v48i4/1000> 617
- Calamus tenuis* Roxb. a new host for
Chilo suppressalis (Walker) (Lepidoptera, Crambidae)
*Selim Mehmud, Sasanka Sekhar Ghosh, Himu Roy,
Dip Jyoti Haloi and Pranab Bujarbarua*

ACKNOWLEDGEMENTS

The EDITORIAL BOARD of ENTOMON profoundly express profuse and sincere thanks and gratitude to the following peer reviewers for sparing their invaluable time in critically going through the manuscripts and suggesting constructive comments and views on the articles published in the ENTOMON volume 48, issues 1 to 4, during 2023.

Abdulraheem Mukhtar Iderawumi, Department of Agricultural Science Education, Oyo State College of Education, P.M.B. 001, Lanlate, Oyo State, Nigeria.

Abraham Verghese, Former Director, ICAR-National Bureau of Agricultural Insects Resources, Bengaluru 560024, Karnataka, India.

Aijaz A Wachkoo, Assistant Professor, Department of Zoology, Government Degree College, Shopian, Gagren Shopian 192303, Jammu and Kashmir, India.

Alejandro Estrada, Estación de Biología Chamela, Instituto de Biología, UNAM, Apartado Postal 21, 48980. San Patricio, Jalisco, México.

Anil Kumar Dubey, Zoological Survey of India, Andaman and Nicobar Region Centre, Port Blair 744102, Andaman and Nicobar Islands, India.

Anil Kumar Sethy, Institute of Wood Science & Technology, Institutes of Indian Council of Forestry Research & Education, Bengaluru 560003, Karnataka, India.

Ankita Gupta, ICAR-National Bureau of Agricultural Insect Resources, H.A. Farm Post, Bellary Road, Hebbal, Bengaluru 560024, Karnataka, India.

Aparna S. Kalawate, Scientist-E, Zoological Survey of India, Western Regional Centre, Ministry of Environment, Forest & Climate Change, Vidhya Nagar, Akurdi, Pune 411044, Maharashtra, India.

Arunachalam N., Scientist G (Rtd.), Indian Council for Medical Research - VCRC, Madurai, Tamil Nadu, India.

Ashish D. Tiple, Head, Department of Zoology, Vidya Bharati college, Seloo, Wardha 442105, Maharashtra, India

Avtar Kaur Sidhu, High Altitude Regional Centre, Zoological Survey of India, Saproon, Solan 173211, Himachal Pradesh, India.

Balasubramanian, R., ICMR-National Institute of Virology-Kerala Unit, Dept. of Health Research, Ministry of Health & Family Welfare, Government of India, Govt. T. D. Medical College Hospital, Vandanam, Alappuzha 688005, Kerala, India.

Baskaran, V., Assistant Professor, Acarology, Agricultural Entomology Faculty, Centre for Plant Protection Studies, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India.

Basker Parasuram, Consultant Entomologist, National Centre for Disease Control, Directorate General of Health Services, Ministry of Health & Family Welfare, Government of India, Delhi, India.

Belavadi, V.V., Former Emeritus Scientist, Department of Entomology, University of Agricultural Sciences, GKVK, Bengaluru 560065, Karnataka, India.

Bhuvaneswari, K., Professor, Department of Agricultural Entomology, Centre for Plant Protection Studies, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India.

- Bijoy, C., Assistant Professor and Research Supervisor, Department of Zoology, Christ College (Autonomous), Irinjalakuda, Thrissur 680125, Kerala, India.
- Binoy, C.F., Dean of Sciences, Associate Professor, HoD, Research & PG Dept. of Zoology, St. Thomas' College (Autonomous), Thrissur 680001, Kerala, India.
- Chinna Babu Naik, V., ICAR-Central Institute for Cotton Research. Nagpur, Shankar Nagar, Nagpur 440010, Maharashtra, India.
- Chitra Narayanasamy, Professor & Curator, TNAU Insect Museum, Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India.
- Christoph Hörweg, Head of the Zoology (Invertebrates), Curator Collection Arachnoidea, Natural History Museum Vienna, Burgring 7, A-1010 Vienna, Austria.
- Debjani Dey, Principal Scientist, Division of Entomology, ICAR - Indian Agricultural Research Institute, New Delhi 110012, India.
- Devanshu Gupta, Officer-in-Charge & Scientist-C, Coleoptera Section, Zoological Survey of India, M-Block, New Alipore, Kolkata 700053, West Bengal, India.
- Devasahayam, S., Principal Scientist (Rtd), ICAR - Indian Institute of Spices Research, Vellimadukunnu, Kozhikode 673012, Kerala, India.
- Dinendra Raychaudhuri, Arachnologist, Entomology Laboratory, Department of Zoology, University of Calcutta, 35, Ballygunge Circular Road, Kolkata 700019, West Bengal, India.
- Enas Moustafa Yousef Mohammed, Senior Researcher, Plant Protection Research Institute, Agriculture Research Center, Egypt.
- Evans D. Asirvadam, Associate Professor (Rtd), Department of Zoology, University College, Thiruvananthapuram 695034, Kerala, India.
- George Mathew, Scientist G and Programme Coordinator, Forest Health Division (Rtd.), International Forest Insect Pest Specialist (ADB Project), Kerala Forest Research Institute, Peechi 680653, Kerala, India.
- Gregory D. Edgecombe, Specialist in Arthropods, Zoology, Monophyly, Evolutionary biology and Sister group, Natural History Museum, London.
- Guido Sabatinelli, Membre correspondant du Muséum, Secteur Invertébrés, Muséum d'histoire naturelle (MHN), Département de la culture et de la transition numérique, Route de Malagnou 1, 1208 Geneve.
- Guy Tomasovic, Université de Liège, Gembloux Agro-Bio Tech, Passage des Déportés, 25030 Gembloux – Belgique, France.
- Harry Brailovsky, Instituto de Biología UNAM, Depto. de Zoología, Apod Postal 04510, Ciudad de Mexico.
- Hassan, M.E., Scientist – E, Zoological Survey of India, Gangetic Plains regional Centre, Sector -8, Vijay Nagar, Bahadurpur Housing Colony, Patna 800026, Bihar, India
- Hegde, V.D., Scientist-E, Zoological Survey of India, Western Ghat Regional Centre, S.V. Colony, Balan K.Nair Road, Kozhikode 673006, Kerala, India.

- Hemant V. Ghate, Professor, Department of Zoology, Modern College, Shivaji Nagar, Pune 411005, Maharashtra, India.
- Hideyuki Chiba, Research Associate, Bernice Pauahi Bishop Museum, 1525 Bernice Street, Honolulu, Hawaii 96817-2704, USA.
- Himender Bharti, Professor, Ant Systematics and Molecular Biology Lab., Department of Zoology and Environmental Sciences, Punjabi University, Patiala 147002, Punjab, India.
- Irinel E. Popescu, Associate Professor, Department of Biology, Faculty of Biology -Entomology, Al. I. Cuza University, Bd. Carol I nr. 20A, 700506 Iasi, Romania.
- Jagdish Saini, Scientist, Zoological Survey of India, M-Block, New Alipore, Kolkata 700053, West Bengal, India.
- Jayita Sengupta, Senior Zoological Assistant, Zoological Survey of India, M Block, New Alipore, Kolkata 700053, West Bengal, India.
- Jerry L. Cook, Director, Sam Houston Natural History Collections, Department of Biological Sciences; Sam Houston State University; Huntsville, TX 77341-2116; U.S.A.
- Jobiraj, T., Assistant professor and Research Guide, PG Department of Zoology, Government college Kodanchery, Kozhikode 673580, Kerala, India.
- John Caleb, Entomology division, Loyola College, Nelson Manickam Rd, Nungambakkam, Chennai 600034, Tamil Nadu, India.
- Jorge Ari Noriega, Specialist, Department of Biogeography and Global change, Cra. 6b No. 113-51, Bogotá, Colombia – Calle General Pardiñas 22, Madrid, Spain.
- Josephraj Kumar, A., ICAR - Central Plantation Crops Research Institute, Regional Station, Kayamkulam 690502, Kerala, India.
- Kaomud Tyagi, Scientist, Centre for DNA Taxonomy, Molecular Systematics Division, Zoological Survey of India, Kolkata 700053, West Bengal, India.
- Kathirvelu, C., Associate Professor, Department of Entomology, Faculty of Agriculture, Annamalai University, Annamalainagar 608002, Chidambaram, Tamil Nadu, India.
- Kesavan Subaharan, ICAR - National Bureau of Agricultural Insect Resources, Bengaluru 560064, Karnataka, India.
- Khirod Sankar Das, Department of Zoology, North-Eastern Hill University, Shillong 793022, Meghalaya, India.
- Korada Rajasekhara Rao, Head, Division of Crop Management, ICAR- Central Tobacco Research Institute, Rajahmundry 533105, Andhra Pradesh, India.
- Kuldeep Srivastava, Principal Scientist (Agricultural Entomology), Division of Crop Protection, ICAR - Indian Institute of Vegetable Research, Jakkhini, Varanasi 221305, Uttar Pradesh, India.
- Kumar Ghorpade, Emeritus Scientist, University of Agricultural Sciences, Dharwad 580005, Karnataka, India.
- Manjunath Gowda, Professor of Sericulture, Silkworm Genetics and Breeding, Silkworm Physiology and Silkworm Pathology, University of Agricultural Sciences, Bengaluru 560065, Karnataka, India.

- Mark S. Harvey, Head, Terrestrial Zoology, Curator of Arachnids & Myriapods, Western Australian Museum, Locked Bag 49, Welshpool DC, Western Australia 6986, Australia.
- Mathew M.J., Department of Zoology, Sacred Heart College, Thevara, Kochi 682013, Kerala, India.
- Meenakshi Bharti, Professor, Department of Zoology & Environmental Sciences, Punjabi University, Patiala 147002, Punjab, India.
- Mingzhi Zhao, Department of Entomology, College of Plant Protection, South China Agricultural University, 483 Wushan Road, Guangzhou 510642, China.
- Mohankumar, S., Professor, Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India.
- Mohd Yousuf, Dean School of Life Sciences, Central University of Kashmir, Nuner, Ganderbal 191201, Jammu and Kashmir, India.
- Muhammed Jafer Palot, Zoological Survey of India, Western Ghat Regional Centre, Jafer Khan Colony, Erannhipalam, Kozhikode, Kerala 673006, India.
- Murugan, M., Professor, Department of Agricultural Entomology, Centre for Plant Protection Studies, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India.
- Muzafar Riyaz, Senior Research Fellow, Entomology division, Entomology Research Institute, Loyola College, Nungambakkam, Chennai 600034, Tamil Nadu, India.
- Nasser, M., Professor, Department of Zoology, University of Calicut, Calicut 673635, Kerala, India
- Noppadon Makbun, 211/5 Moo 4, Takhli, Nakhon Sawan 60140, Thailand.
- Omprakash Navik, Division of Germplasm Collection and Characterisation, ICAR-National Bureau of Agricultural Insects Resources, Bengaluru 560024, Karnataka, India.
- Padmanaban, B., Former Principal Scientist (Agril.Entomology), Agricultural Research Service of ICAR. ICAR-National Research Centre for Banana, Tiruchirapalli 620006, Tamil Nadu, India.
- Pandey, J.P., Scientist, Silkworm Biotechnology Laboratory, Central Tasar Research & Training Institute, Central Silk Board, Patrachauli, Ranchi 835303, Jharkhand, India.
- Panicker K.N., Professor Emeritus, Amrita Institute of Medical Sciences, Kochi 682041, Kerala, India [Former Sr. Scientist ICMR, WHO / TDR Consultant Geneva. Rockfeller Foundation Consultant, Honduras. Executive Director and Chairman SEUF (Dutch/World Bank/UNICEF/DFID)].
- Pankaj Koparde, School of Sustainable Development, Faculty of Sustainability Studies, Dr. Vishwanath Karad MIT World Peace University, Pune, Maharashtra, India.
- Paramasivan R, Scientist 'F' & Officer in Charge, ICMR-Vector Control Research Centre, Indian Council of Medical Research, Department of Health Research, Ministry of Health & Family welfare, Govt of India, Chinna Chokkikulam, Madurai 625002, Tamil Nadu, India.
- Pawan U. Gajbe, Associate Professor, Department of Zoology, Shri Mathuradas Mohota College of Science, Sakardara, Chowk, Umred Rd, Nagpur 440009, Maharashtra, India.
- Phan Quoc Toan, Director, Center for Entomology & Parasitology Research Institute of Research & Training of Medicine, Biology & Pharmacy, Duy Tan University, Hoang Minh Thao Street, Lien Chieu District, Da Nang, Vietnam.

- Prakash, K.V., Assistant Entomologist, Department of Agricultural Entomology, GKVK, University of Agricultural Sciences, Bengaluru 560065, Karnataka, India.
- Prasad, G., Professor and Head of Department of Zoology, University of Kerala, Kariavattom, Tiruvananthapuram 695 581, Kerala, India.
- Prasada Rao, G.M.V., Acharya N.G. Ranga Agricultural University, Regional Agricultural Research Station, Guntur 522 034, Andhra Pradesh, India
- Premalatha, K., Assistant Professor, Termite taxonomy and management; Agricultural Entomology Faculty, Centre for Plant Protection Studies, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India.
- Priyadarsanan Dharma Rajan, Senior Fellow, Ashoka Trust for Research in Ecology and the Environment (ATREE), Royal Enclave, Srirampura, Jakkur, Bengaluru 560 064, Karnataka, India.
- Rahul Joshi, Scientist 'D' (Lepidoptera Section), CPIO, Zoological Survey of India, M-Block, New Alipore Kolkata 700053, West Bengal, India.
- Rajmohana, K., Scientist-E, Officer in charge Miscellaneous Insect Order Section and Isoptera Section, Coordinator, ENVIS Centre on Faunal Diversity, Zoological Survey of India, New Alipore, Kolkata 700053, West Bengal, India.
- Raman, A., Professor, CSIRO (Health and Biosecurity), Underwood Avenue, Floreat Park, WA 6014 & Charles Sturt University, PO Box 883, Orange, NSW 2800, Australia.
- Ramaraju, K., Professor of Agricultural Entomology (Rtd.), Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India.
- Ramasubramanian, T., Principal Scientist, Division of Crop Protection, ICAR - Sugarcane Breeding Institute, Coimbatore 641007, Tamil Nadu, India.
- Rameash, K. Principal Scientist -Agricultural Entomology, ICAR - Central Institute for Cotton Research, Regional Station, Coimbatore 641 003, Tamil Nadu, India
- Rameshkumar, A., Scientist - D, Hymenoptera Section, Zoological Survey of India, Ministry of Environment, Forest & Climate Change, Kolkata 700 053, West Bengal, India.
- Reghu Ravindran, Assistant Professor, Department of Veterinary Parasitology, College of Veterinary and Animal Sciences, Pookode, Kerala Veterinary And Animal Sciences University, Lakkidi 673576, Wayanad, Kerala, India.
- Sabu K. Thomas, Professor & Director, Malaviya Mission Teacher Training Centre, University Of Calicut, Malappuram 673 635, Kerala, India.
- Sajan, K.C., Pokhara, Kaski 33700, Gandaki Province, Nepal; Eastern New Mexico University, Portales, NM, United States.
- Samiran Chakrabarti, Formerly: UGC Emeritus and Professor & Head, Department of Zoology, University of Kalyani, Kalyani 741235, India. Correspondence: 32/B. Green Park, Belghoria, Kolkata 700 056, India.
- Sampathkumar, M., Senior Scientist (Entomology), Division of Germplasm Collection and Characterization, ICAR - National Bureau of Agricultural Insect Resources, H.A. Farm post, Bengaluru 560024, Karnataka, India.

- Sanborn, A., Department of Biology, Barry University, 11300 NE Second Avenue, Miami Shores, Florida, 10 33161-6695, USA.
- Sandeep Singh, Principal Scientist - Entomology, All India Coordinated Research Project on Fruits, Department of Fruit Science, Punjab Agricultural University, Ludhiana 141004, Punjab, India.
- Sanjay Sondhi, Indian Foundation for Butterflies. C-703, Alpine Pyramid, Rajiv Gandhi Nagar, Bengaluru, Karnataka 560097, India.
- Santhosh Shreevihar, Asst. Professor, Department of Zoology, Malabar Christian College, Kozhikode 673001, Kerala, India.
- Selva Narayanan, V., Professor, Department of Entomology, Faculty of Agriculture, Annamalai University, Annamalainagar 608 002, Chithambaram, Tamil Nadu, India.
- Selvanarayanan, V., Professor, Department of Entomology, Faculty of Agriculture, Annamalai University, Annamalainagar 608 002, Chithambaram, Tamil Nadu, India.
- Shashidhar Viraktamath, Professor, Biosystematic Laboratory, Department of Entomology, University of Agricultural Sciences, GKVK, Bengaluru 560065, Karnataka, India.
- Shelley Acharya, Scientist, Zoological Survey of India, New Alipore, Kolkata 700053, West Bengal, India.
- Singh, B.N., BHU Distinguished Professor, Department of Zoology, Institute of Science, Banaras Hindu University, Varanasi-221005, India.
- Smriti Sharma, Sr. Scientist (Residue Analysis), Department of Entomology, Punjab Agricultural University, Ferozepur Rd, Ludhiana 141027, Punjab, India.
- Souvik Sen, Scientist-D, Officer-in-Charge, Arachnida Section, Zoological Survey of India. New Alipore, Kolkata 700053, West Bengal, India.
- Sreekumar, K.M., Professor and Head, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Padnekkad, Nileswarem 671314, Kasaragod, Kerala, India.
- Srikanth, J., Principal Scientist (Rtd), Section of Entomology, Division of Crop Protection, ICAR - Sugarcane Breeding Institute, Coimbatore 641007, Tamil Nadu, India.
- Srinivasa, N., All India Network Project on Agricultural Acarology, Department of Agricultural Entomology, University of Agricultural Sciences, GKVK, Bengaluru 560065, Karnataka, India.
- Srivastava, G.K., 1/56, Vibhav Khand, Gomtinagar, Lucknow 226010, Uttar Pradesh, India.
- Subbiah Poopathi, Scientist-G, Head of the Unit of Microbiology and Molecular Biology, Vector Control Research Centre, Indian Council of Medical Research, Department of Health Research, Ministry of Health & Family Welfare, Govt. of India, Medical Complex, Indira Nagar, Pondicherry 605 006, India.
- Subhankar Kumar Sarkar, Entomology Laboratory, Department of Zoology, University of Kalyani, Kalyani 741235, West Bengal, India.
- Subramanian, K.A., Head, Southern Regional Centre, Zoological Survey of India, Chennai 600028, Tamil Nadu, India.
- Sudhir Singh, Professor, Systematic Laboratory, Forest Entomology Division, Forest Research Institute, Dehradun 248006, Uttarakhand, India.

- Sundararaj, R., Scientist- G and Head, Forest and Wood Protection Division, Institute of Wood Science and Technology, Bangalore., India.
- Suprakash Pal, Professor in Agricultural Entomology, Entomology Lab, Regional Research Station (Terai Zone), Directorate of Research, Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar 736165, West Bengal, India.
- Suresh, S., Professor of Agricultural Entomology (Retired), Tamil Nadu Agricultural University, A1, TNHB, S.N. Palayam, Coimbatore 641007, Tamil Nadu, India.
- Sureshan, P.M., Scientist, Western Ghats Field Research Centre, Zoological Survey of India, Eraniel, Kozhikode 673006, Kerala, India.
- Susmita Gupta, Professor, Department of Ecology and Environmental Science, Assam University, Silchar 788011, Assam, India.
- Swaminathan Subramanian, Indian Council of Medical Research, Department of Health Research, Ministry of Health & Family Welfare, Govt. of India, Medical Complex, Indira Nagar, Pondicherry 605 006, India.
- Swaminathan, R., Professor and Former Dean (Rtd.), Department of Entomology, Rajasthan College of Agriculture, Maharan Pratap University of Agriculture and Technology, Udaipur 313001, Rajasthan, India.
- Talmale, S.S., Scientist-C, Western Regional Centre, Zoological Survey of India, Ministry of Environment, Forest and Climate Change, Akurdi, Pune 411 044, Maharashtra, India.
- Thomas Pape, Lars Vilhelmsen and Sree Gayathree Selvantharan, Natural History Museum, Copenhagen, Denmark.
- Tosaphol Saetung Keetapithchayakul, Department of Zoology, Faculty of Science, Kasetsart University, 50 Ngam Wong Wan Rd, Lat Yao Chatuchak, Bangkok.
- Tristan McKnight, Department of Entomology, University of Arizona, 410 Forbes Bldg, Tucson, Arizona, 85721 USA.
- Upadhyay R.K., Department of Zoology, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur 273009, Uttar Pradesh, India.
- Vasantharaj David, B., 76/2A Sree Ramulu st., Santhosh Nagar, Madanandapuram, Chennai 600125, Tamil Nadu, India.
- Vasuki Belavadi, V., Emeritus Scientist, Department of Entomology, University of Agricultural Sciences, GKVK, Bengaluru 560065, , Karnataka, India.
- Vijaya Kumar, K.T., Professor, AICRP-Honey Bees & Pollinators, Department of Apiculture, University of Agricultural Sciences, GKVK, Bengaluru 560065, Karnataka, India.
- Viraktamath, C.A., Emeritus Scientist, Department of Entomology, University of Agricultural Sciences, GKVK, Bengaluru 560065, Karnataka, India.
- Virendra Prasad Uniyal, Sr. Professor & Scientist G, Wildlife Institute of India, Chandrabani, Dehradun 248171, Uttarakhand, India.
- Vishlesh Shankar Nagrare, Principal Scientist (Agricultural Entomology), ICAR - Central Institute for Cotton Research, Shankar Nagar, Nagpur 440010, Maharashtra, India.

Vivek Sarkar, WII C2C, Wildlife Institute Rd, Chandrabani, Mehu Wala Mafi, Dehradun 248001, Uttarakhand, India.

Winston M.O.Thompson, P.O. Box 7226, Bellevue, Washington, WA 98008, United States.

Yuvarajan, P., Assistant Professor, Department of Aquaculture, Dr. M.G.R Fisheries College and Research Institute, Tamil Nadu Dr. J. Jayalalithaa Fisheries University, Thalainayeru 614 712, Tamil Nadu, India.

Zsolt Bálint, Lead curator, Lepidoptera collection, Department of Zoology, Hungarian Natural History Museum. 1088-Budapest, Baross utca 13, Hungary.

AUTHOUR INDEX

- | | |
|---------------------------|--------------------------------|
| Abhisubesh V., 33, 605 | Balasubramani Venkatasamy, 323 |
| Abraham Samuel K., 253 | Basavaraj Naik T., 595 |
| Aishwarya Palanisamy, 323 | Bernad M Thampan, 167 |
| Aishwarya S. Naik, 591 | Bhagyashree Bora, 33, 605 |
| Ajitha T., 39 | Bhanderi G.R., 475 |
| Albright B., 141 | Bhramar Bhattacharyya, 123 |
| Amala Anna Francis, 141 | Bhushan L.S., 309 |
| Amartya Pal, 525 | Bhuvaneswari Kaithamalai, 19 |
| Amr Mohamed, 383 | Bijoy C., 331, 441 |
| Anandamay Barik, 123 | Bikash Subba, 599 |
| Aneha K., 33, 605 | Binoy C.F., 449 |
| Anil John J., 39 | Bratatee Sardar, 241 |
| Anil Kumar Dubey, 77 | Buhroo A. A., 421 |
| Anila L., 531 | Champa B.V., 595 |
| Anitha Mamillapalli, 383 | Chandrasekaran S., 209 |
| Anjitha K Bhadrn, 455 | Chaueichongla Phom, 471 |
| Anjusha Bhasker, 293 | Chethana O., 595 |
| Anushya A.V., 293 | Chimasha Induni, 371 |
| Anzil Shereef, 167 | Chinnu V.S., 545 |
| Arati Pannure, 317 | Chitra Narayanasamy, 323 |
| Archana B., 445 | Chris M. Cohen, 411 |
| Arunima Choudhury, 185 | Dattaprasad Sawant, 99 |
| Ashish Kumar Jha, 149 | Deen Mohd Bhat, 303 |
| Ashish V.V., 537 | Deepthy K.B., 537 |
| Ashok Karuppannasamy, 229 | Deepti B., 365 |
| Ayana Gayathri R.V., 39 | Desai H.R., 475 |
| Ayesha E. Shetkar, 591 | Devanshu Gupta, 525 |
| Baiju K.V., 439 | Devi H. S., 427 |
| Baiju Kochunarayanan, 167 | Dhaneesh Bhaskar, 331 |
| Baiju Kochunarayanan, 553 | Dhaneesh Bhaskar, 95 |

- Dharma Rajan Priyadarsanan, 433
 Dip Jyoti Haloi, 185
 Dip Jyoti Haloi, 617
 Dipendra Nath Basu, 1
 Edwin Joseph, 141
 Evans D.A., 39
 Fernando H.S.D., 371
 Francy K. Kakkassery, 497
 Garaniya N.H., 63
 Gayathri Elayidam U., 439
 Gianluca Tettamanti, 383
 Gigi Poulouse, 411
 Girish Kumar P., 461
 Gokulakrishnaa R.K., 583
 Gopianand L., 287
 Haseena Bhaskar, 95, 537
 Hashim M.S., 531
 Hegde V. D., 461, 467
 Hemaladkshmi P., 33, 605
 Hemant Ogale, 99
 Himu Roy, 617
 Hothur Lakshmi, 383
 Hyzil J.B., 439
 Innifa Hasan, 185
 Jagbir Singh Kirti, 83
 Jagruti Roy, 149
 Jayakumar Pathma, 309
 Jayalaxmi Narayan Hegde, 595
 Jayaprakas C.A., 511
 Jebine Jose, 167
 Jebine Jose, 553
 Jeevitha P., 575
 Jeyaparvarthi Somasundaram, 471
 Jibi Lukose, 33
 Jibi Lukose, 605
 Jobin Mathew, 141
 Jobiraj T., 415
 Joy Sharmila E., 445
 Jyothi Sara Jacob, 89
 Jyothylakshmi K., 611
 Kakhuangailiu Gangmei, 33
 Kakhuangailiu Gangmei, 605
 Kalaiselvi S., 445
 Kalesh Sadasivan, 1, 167, 253, 553
 Kathirvelu C., 287
 Kavitha R.S., 111
 Kavya G. Pillai, 411
 Kavyamol P.M., 117
 Khirod Sankar Das, 135
 Krushnamegh Kunte, 1
 Kulanthaisami S., 19
 Lekeshmenaswamy M., 129
 Longkumer B., 427
 Mahendra B.V., 111
 Maheswara Reddy P., 51
 Makesh kumar T., 511
 Manendra Kaneria, 299
 Mani Chellappan, 95
 Manickavasagam S., 287
 Manikanda Boopathi N., 11
 Manikandan S., 33
 Manikandan S., 605
 Manjunatha K. L., 317
 Manjusha K.T., 485

- Manoj Kripakaran, 553
Mathivanan A., 33
Mathivanan A., 605
Mawtham M.M., 19
Mendis B.A.N., 371
Merlin Kamala I., 209
Mick Webb, 167
Mohd Faisal, 299
Mohsin Ikram, 299
Muchhadiya D.V., 63
Mukul Kumar Deka, 575
Muralimohan P.V., 167
Murugan Marimuthu, 323
Muthuswami M., 11
Namitha Jayan, 141
Nandakumar S., 611
Narayana, R., 303
Nayan Roy, 241
Nayan Thara, J., 303
Neena Lenin, 95
Neha Chowdhury, 241
Nihal Nila J., 219
Nilanjana Chudhuri, 599
Nissanka K. De Silva B.G.D., 371
Padma-Shree I., 11
Pandi V., 445
Pankaj Neog, 427
Parmar P.R., 475
Patel D.R., 63
Patel J.J., 63
Patel R.D., 475
Pathma J., 365
Pobi Gogoi, 185
Poopathi S., 33
Poopathi S., 605
Pradeepkumar G.T., 357
Prajna Devaiah B., 229
Prakash Chandra Mardaraj, 397
Prakash K.V., 567
Pramod Katti, 229
Pranab Bujarbarua, 617
Prashant N. Bavachikar, 383
Prathibha M., 111
Prdeepkumar R., 39
Rajashekharappa K., 595
Rajeshwari L.S., 511
Rajeswari S., 11
Rajkumar Chidambaram Palaniappan, 553
Raju P. J., 383
Rakesh M. Deulkar, 99
Ramani N., 455
Ramasamy Kamaya Naicker S., 553
Rangesh K., 445
Ranjith A.P., 433
Reject Paul M. P., 449
Reshma V.R., 39
Sabu K. Thomas, 461
Saha P.C., 525
Sahadiya Mandodan, 33, 605
Sainamol P. Kurian, 107
Salini S., 89
Sangeetha B.G., 511
Sankarsan Roy, 397
Santanu Ghara, 397

- Santhosh kumar T., 511
Santu Paria, 397
Sasanka Shekhar Ghosh, 617
Selim Mehmud, 617
Selvamuthukumaran Thirunavukkarasu, 583
Senapati, 599
Senguttuvan K., 11
Shanas S., 89, 545
Shankara Murthy M., 51
Shashidhar Viraktamath, 149
Shaun Paul Adambukulam, 497
Sheema Tabasum, 421
Shirly Raichal Anil 511
Shivakumar Veerlapati, 69
Shubham Rao, 149
Sobhana K. A., 195, 461, 467
Sornapriya J., 129
Souren Dutta, 241
Sreeja P., 95, 107
Sreekumar K.M., 575
Sreelatha U., 537
Srinivasa N., 69
Srujan Ogale, 99
Staklin Stella S., 83
Stevens Jones R.D., 2019
Subhash Chandran K.P., 357
Sudarmani D.N.P. , 519
Sudipta Choudhury, 135
Suganthi Angappan, 19
Sujitha Prabhakaran Chandrika, 553
Sumithramma N., 545
Sundareswari, 519
Sunil Kumaraswamy, 229
Susanta Kumar Chakraborty, 397
Susaritha S., 445
Sushila Nadagouda, 229
Swaran P.R., 293
Swetapadma Dash, 525
Tarun Goyal, 83
Thasnim E. S., 331
Thomas K. Sabu, 195
Vasuki V. Belavadi, 317
Venkatesh Tanagal, 51
Vidya Mulimani, 317, 545
Vijayalakshmi K., 33, 605
Vijayasree V., 357
Vijaykumar S. Gadekar, 591
Vinayan P. Nair, 253
Vinayan Padmanabhan Nair, 553
Vineesh P.J., 117
Vineetha V.P., 117
Viswanath T.K., 467
Waluniba, 427
Waseem Ahmad, 299
Yeshwanth H.M., 567

Statement of ownership and other particulars of ENTOMON

(Form IV, Rule 8 of Registration of Newspapers (Central) Rules 1956)

1. Place of publication : Trivandrum
2. Periodicity of publication : Quarterly
3. Printer's name, nationality and address : Dr K D Prathapan, Indian,
Secretary,
Association for Advancement of Entomology,
Department of Agricultural Entomology,
College of Agriculture,
Kerala Agricultural University, Vellayani PO,
Thiruvananthapuram 695522, Kerala, India
4. Publisher's name, nationality and address : - do-
5. Editor's name, nationality and address : Dr M S Palaniswami, Indian,
Chief Editor, ENTOMON,
Association for Advancement of Entomology,
Thiruvananthapuram 695522, Kerala, India
6. Name and address of the
Individual who owns the paper : Association for Advancement of Entomology,
Department of Agricultural Entomology,
College of Agriculture,
Kerala Agricultural University, Vellayani PO,
Thiruvananthapuram 695522, Kerala, India

I, Dr K. D. Prathapan, Secretary, Association for Advancement of Entomology, here by declare that the particulars given above are true to the best of my knowledge and belief.

Sd/-

Vellayani PO, Thiruvananthapuram 695522

Dr K. D. Prathapan

31 December 2023

Publisher, ENTOMON



Association for Advancement of Entomology

(Reg. No. 146/ 1975)

Department of Agricultural Entomology, Kerala Agricultural University,
Vellayani, Thiruvananthapuram 695522, Kerala, India. web: www.entomon.in;

Email: aae@kau.in

EXECUTIVE COMMITTEE (2022 – 2025)

President: Professor N. Mohandas, Former Head, Department of Agricultural Entomology & Research Coordinator, Kerala Agricultural University, Thiruvananthapuram, 695522, Kerala, India.

Vice Presidents:

1. Professor S. Devanesan, Former Dean, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram 695522, Kerala, India.
2. Professor G. Prasad, Head of Department of Zoology, University of Kerala, Kariavattom, Thiruvananthapuram 695581, Kerala, India.
3. Dr. R. Rajendran, Consultant, National Centre for Disease Control, Govt of India, Cherthala 688524, Kerala, India.

Secretary: Dr. K. D. Prathapan, Department of Agricultural Entomology, Kerala Agricultural University, Vellayani, Thiruvananthapuram 695522, Kerala, India.

Joint Secretaries:

1. Dr. D. A. Evans, Former Associate Professor, Department of Zoology, University College, Thiruvananthapuram 695 034, Kerala, India.
2. Professor O. P. Reji Rani, Department of Agricultural Entomology College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram 695522, Kerala, India.
3. Dr. S. Shanas, Assistant Professor (Agricultural Entomology), Integrated Farming Systems Research Station, Kerala Agricultural University, Karamana, Thiruvananthapuram 695002, Kerala, India.

Treasurer: Dr. Thania Sara Varghese, Assistant Professor, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram, 695522, Kerala, India.

Members:

1. Professor A. Visalakshi, Former Professor and Head, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram, 695522, Kerala, India.
2. Dr. Hebsibai, Former Professor, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram, 695522, Kerala, India.
3. Professor G. Madhavan Nair, Former Head, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram, 695522, Kerala, India.
4. Dr. C. A. Jayaprakas, Principal Scientist, Division of Crop Protection, ICAR – Central Tuber Crops Research Institute, Sreekariyam, Thiruvananthapuram, 695017, Kerala, India.
5. Dr. T. Sivakumar, Subject Matter Specialist (Entomology), KVK, ICAR- Central Plantation Crops Research Institute, Kayamkulam 690533, Kerala, India.
6. Dr. T. Santhoshkumar, Assistant Professor, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram, 695522, Kerala, India.
7. Dr. S. Poopathi, Scientist-G (Director Grade), Head of Microbiology and Molecular Biology, Vector Control Research Centre, Indian Council of Medical Research, Pondicherry 605 006, India.
8. Dr. Ambili Paul, Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram, 695522, Kerala, India.
9. Dr. E. R. Harish, Scientist, Division of Crop Protection, ICAR – Central Tuber Crops Research Institute, Sreekariyam, Thiruvananthapuram, 695017, Kerala, India.
10. Dr. M. S. Palaniswami, Chief Editor, ENTOMON, AAE.
11. Dr. K. M. Sreekumar, Professor and Head, Department of Agricultural Entomology, Kerala Agricultural University, College of Agriculture, Padnekkad, Nileswaram, Kasaragod, Kerala, India.
12. Dr. C. Bijoy, Assistant Professor and Research Supervisor, Department of Zoology, Christ College (Autonomous), Irinjalakuda, Thrissur 680 125, Kerala, India.



Published by :
Association for Advancement of Entomology
Email : aae@kau.in; web: www.entomon.in